# MONTANA DEPARTMENT OF TRANSPORTATION WETLAND MITIGATION MONITORING REPORT: YEAR 2001

Creston Site Creston, Montana



Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION 2701 Prospect Ave Helena, MT 59620-1001

July 2002

Project No: 130091.007

Prepared by:

LAND & WATER CONSULTING, INC. P.O. Box 8254 Missoula, MT 59807



# MONTANA DEPARTMENT OF TRANSPORTATION

# WETLAND MITIGATION MONITORING REPORT:

# **YEAR 2001**

Creston
Creston, Montana

# Prepared for:

# MONTANA DEPARTMENT OF TRANSPORTATION

2701 Prospect Ave Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING, INC.

P.O. Box 8254 Missoula, MT 59807

July 2002

Project No: 130091.007



# TABLE OF CONTENTS

1.0	INTRODUCTION	.1
2.0	METHODS	.1
	2.1 Monitoring Dates and Activities	. 1
	2.2 Hydrology	. 1
	2.3 Vegetation	.3
	2.4 Soils	.3
	2.5 Wetland Delineation	.3
	2.6 Mammals and Herptiles	.4
	2.7 Birds	.4
	2.8 Macroinvertebrates	.4
	2.9 Functional Assessment	.4
	2.10 Photographs	.4
	2.11 GPS Data	.5
	2.12 Maintenance Needs	.5
3.0	RESULTS	.5
	3.1 Hydrology	.5
	3.2 Vegetation	.5
	3.3 Soils	.7
	3.4 Wetland Delineation	.7
	3.5 Wildlife	.7
	3.6 Macroinvertebrates	.8
	3.7 Functional Assessment	.8
	3.8 Photographs	.8
	3.9 Maintenance Needs/Recommendations	.8
	3.10 Current Credit Summary	.8
4.0	REFERENCES	9



# **TABLES**

Table 1 2001 Creston Vegetation Species List
 Table 2 Fish and Wildlife Species Observed at the Creston Mitigation Site
 Table 3 Summary of 2001 Wetland Function/Value Ratings and Functional Points at the Creston Mitigation Project

# **FIGURES**

Figure 1 Project Site Location Map

## **APPENDICES**

Appendix A: Figures 2 - 3

Appendix B: Completed 2001 Wetland Mitigation Site Monitoring Form

Completed 2001 Bird Survey Forms

Completed 2001 Wetland Delineation Forms

Completed 2001 Field and Full Functional Assessment Forms

Appendix C: Representative Photographs

Appendix D: Bird Survey Protocol

Macroinvertebrate Sampling Protocol

GPS Protocol



#### 1.0 INTRODUCTION

The Creston mitigation site was constructed in 1998 to mitigate wetland impacts associated with three Montana Department of Transportation (MDT) roadway projects; the Flathead River Bridge and Creston North and South projects. The site is located one mile south of the Creston Fish Hatchery adjacent to Highway 35 and Broeder Loop (**Figure 1**). The site consists of 20 acres located in Flathead County within the Flathead River Watershed (No. 4). The site elevation is 2,940 feet above mean sea level (amsl).

The site was designed to mitigate for riparian floodplain habitat, rooted emergent wetland, and ditches associated with previous highway construction. The mitigation goal was to enhance approximately two acres of existing wetland and create four acres of wetland. A formal wetland delineation and functional assessment were not performed prior to construction.

## 2.0 METHODS

# 2.1 Monitoring Dates and Activities

The site was visited on May 8th (spring) and July 25<sup>th</sup> and August 25<sup>th</sup> (mid-season) 2001. The primary purpose of the spring visit was to conduct a bird/general wildlife reconnaissance. The May period was selected for the spring visit because monitoring between mid-May and early June is likely to detect migrant and early nesting activities for a variety of avian species, as well as maximizing the potential for amphibian detection. In Montana, most amphibian larval stages are present by early June.


The mid-season visit was conducted between late July and August to document vegetation, soil, and hydrologic conditions used to map jurisdictional wetlands. All information contained on the Wetland Mitigation Site Monitoring Form (**Appendix B**) was collected at this time. Activities and information conducted/collected included: wetland delineation; wetland/open water boundary mapping; vegetation community mapping; vegetation transects; soils data; hydrology data; bird and general wildlife use; photograph points; GPS data points; functional assessment; and (non-engineering) examination of dike structures.

# 2.2 Hydrology

Hydrologic indicators were evaluated at the site during the mid-season visit. Wetland hydrology indicators were recorded using procedures outlined in the Army Corps (COE) 1987 Wetland Delineation Manual. Hydrology data was recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**).

All additional hydrologic data were recorded on the mitigation site monitoring form (**Appendix B**). The boundary between wetlands and open water (no rooted vegetation) aquatic habitats was mapped on the aerial photograph and an estimate of the average water depth at this boundary was recorded.





Three groundwater-monitoring wells are present on site; however, Land & Water personnel were not able to obtain the correct key to access the wells. Groundwater located within 18 inches of the ground surface (soil pit depth for purposes of delineation), was documented on the routine wetland delineation data form at each data point.

# 2.3 Vegetation

General dominant species-based vegetation community types (e.g., *Elymus repens/Phleum pratense*) were delineated on an aerial photograph during the mid-season visit. Standardized community mapping was not employed as many of these systems are geared towards climax vegetation and may not reflect yearly changes. Estimated percent cover of the dominant species in each community type was listed on the site monitoring form (**Appendix B**).

One 10-foot wide belt transect was established during the mid-season monitoring event to represent the range of current vegetation conditions **Figure 2** (**Appendix A**). Percent cover was estimated for each vegetative species encountered within the "belt" using the following values: + (<1%); 1 (1-5%); 2 (6-10%); 3 (11-20%); 4 (21-50%); and 5 (>50%). The transect will be used to evaluate changes over time, especially the establishment and increase of hydrophytic vegetation. The transect location was marked on the air photo and all data recorded on the mitigation site monitoring form. Transect endpoint locations were recorded with the GPS unit. Photos along each transect were taken from both ends during the mid-season visit.

A comprehensive plant species list for the site was compiled and will be updated as new species are encountered. Ultimately, observations from past years will be compared with new data to document vegetation changes over time. Woody species were planted at this mitigation site. Monitoring relative to the survival of such species was conducted and recorded on the Planted Woody Vegetation Survival Form in **Appendix B**.

## 2.4 Soils

Soils were evaluated during the mid-season visit according to hydric soils determination procedures outlined in the COE 1987 Wetland Delineation Manual. Soil data was recorded for each wetland determination point on the COE Routine Wetland Delineation Data Form (**Appendix B**). The most current terminology used by NRCS was used to describe hydric soils (USDA 1998).

#### 2.5 Wetland Delineation

Wetland delineation was conducted on the 20-acre mitigation site during the mid-season visit according the 1987 COE Wetland Delineation Manual. Wetland and upland areas within the monitoring area were investigated for the presence of wetland hydrology, hydrophytic vegetation and hydric soils. The indicator status of vegetation was derived from the National List of Plant Species that Occur in Wetlands: Northwest Region 9 (Reed 1988). The information was recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). The wetland/upland boundary was delineated on the air photo and recorded with a resource grade GPS unit. The



wetland/upland boundary in combination with the wetland/open water habitat boundary was used to calculate the wetland area developed at each impoundment.

# 2.6 Mammals, Reptiles, and Amphibians

Mammal and herptile species observations and other positive indicators of use, such as vocalizations, were recorded on the wetland monitoring form during each visit. Indirect use indicators, including tracks; scat; burrows; eggshells; skins; bones; etc., were also recorded. Observations were recorded as the observer traversed the site while conducting other required activities. Direct sampling methods, such as snap traps, live traps, and pitfall traps, were not implemented. A comprehensive list of observed species was compiled. Observations from past years will ultimately be compared with new data.

#### **2.7 Birds**

Bird observations were recorded during each visit. No formal census plots, spot mapping, point counts, or strip transects were conducted. During the spring visit, observations were recorded and are shown in **Appendix D**. During the mid-season visit, bird observations were recorded according to the established protocol while conducting the other monitoring activities. During the second visit, observations were categorized by species, activity code, and general habitat association (see field and office data forms in **Appendix B**). Observations from past years will be compared with new data.

#### 2.8 Macroinvertebrates

Macroinvertebrate sampling was not conducted during the 2001 monitoring year. Sampling will be conducted during the spring and summer of 2002 and in future monitoring events, according to the established protocol provided in **Appendix D**.

## 2.9 Functional Assessment

A functional assessment form was completed for the site using the 1999 MDT Montana Wetland Assessment Method. Field data necessary for this assessment were generally collected during each mid-season site visit. An abbreviated field data sheet for the 1999 MDT Montana Wetland Assessment Method was compiled to facilitate rapid collection of field information (**Appendix B**). The remainder of the functional assessment was completed in the office.

# 2.10 Photographs

Photographs were taken during the mid-season visit showing the current land use surrounding the site, the monitored area, and the vegetation transects. Each photograph point location was recorded with a resource grade GPS. The approximate location of photo points is shown on **Figure 2**, **Appendix A**. All photographs were taken using a 50 mm lens. A description and compass direction for each photograph was recorded on the wetland monitoring form.



#### 2.11 GPS Data

During the 2001 monitoring season, point data were collected with a resource grade GPS unit at the vegetation transect beginning and ending locations and at all photograph locations. Wetland boundaries were also recorded with a resource grade GPS unit. The method used to collect these points is described in the GPS protocol in **Appendix D**.

# 2.12 Maintenance Needs

The dike structure was examined during site visits for obvious signs of breaching, damage, or other problems. This did not constitute an engineering-level structural inspection, but rather a cursory examination. No problems were documented. Bird boxes were also inspected and appeared to be in good condition.

#### 3.0 RESULTS

# 3.1 Hydrology

Inundation was present in the two large depressions and was estimated to be 10 to 15% of the mitigation site (see **Figure 3**, **Appendix A**). Emergent vegetation was observed throughout the inundated areas. The water table was depressed relative to previous years due to drought conditions. According to the Western Regional Climate Center, Creston yearly precipitation totals for 2000 (13.91 inches) and 2001 (15.7 inches) were 70 and 79 percent, respectively, of the total annual mean precipitation (19.84 inches) in this area.

The upper pond was virtually dry in mid-July. The artesian well that discharges to the upper pond was flowing but the discharge rate was estimated to be less than one-gallon per minute. Three groundwater wells are located on the site; however, Land & Water was unable to obtain the correct key to access the wells. The wells will be monitored in subsequent monitoring years.

# 3.2 Vegetation

Vegetation species identified on the site are presented in **Table 1** and on the attached data form. Six wetland community types were identified and mapped on the mitigation area (**Figure 3**, **Appendix A**). These included Type 1: *Elymus repens/Phleum pratense*; Type 2: *Typha latifolia*; Type 3: *Typha latifolia* with mixed grasses; Type 4: *Phalaris arundinacea*; Type 5: *Potamogeton pectinatus*; and Type 6: *Alopecurus pratensis*. Dominant species within each of these communities are listed on the attached data form (**Appendix B**).

Type 1 occurs in the upland and consists primarily of *Elymus repens* with an even distribution of *Phleum pratense*, *Agrostis stolonifera*, and *Cirsium arvense*. This community type is weedy and included a trace of *Cynoglossum officinale* (common hound's tongue), which is classified as a noxious weed in Flathead County. Type 2 is present around the pond edges, particularly the upper pond and consists primarily of *Typha latifolia*, *Ceratophyllum demersum* and *Phalaris arundinacea*. Type 3 is present in small depressions with less frequent inundation and consists of *Typha latifolia* mixed with weedy grasses. Type 4 is dominated by *Phalaris arundinacea* and



is present adjacent to the large pond and in some of the small depressions. Type 5 is the emergent vegetation and is dominated by *Potamogeton pectinatus*. Type 6 is a minor upland community that is dominated by *Alopecurus pratensis*. Vegetation transect results are detailed in the attached data form, and are summarized graphically below.

	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,,,,,,,,,,,,,,			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,,,,,,,,	,,,,,,,
VT	Type 1 Upland (195')	Type 2 (80')	Type 3 (63')	Type 4 (100')	Type 5	Total:	VT
M AT	1 ypc 1 Opiana (173 )	1 ypc 2 (00 )	1 ypc 3 (03 )	1 ypt 4 (100 )	Jr	I viai.	4 Y T N
Start				1	(25')	463'	End
South to		~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~					

Table 1: 2001 Creston Vegetation Species List

Species	Region 9 (Northwest) Wetland Indicator
Agrostis stolonifera	FAC+
Alopecurus pratensis	FACW
Barbarea vulgaris	FAC-
Beckmannia syzigachne	OBL
Bromus inermis	
Carex arcta	FACW+
Carex bebbii	OBL
Carex flava	OBL
Ceratophyllum demersum	OBL
Chenopodium album	FAC
Chenopodium rubrum	FACW+
Cirsium arvense	FAC-
Cirsium vulgare	FACU
Cynoglossum officinale	FACU
Dactylis glomerata	FACU
Eleocharis palustris	OBL
Elymus repens	FACU
Epilobium ciliatum	FACW-
Equisetum arvense	FAC
Erigeron lonchophyllus?	FACW
Gnaphalium palustre	FAC+
Juncus articulatus	OBL
Juncus regelii	FACW
Juncus tenuis	FAC
Lactuca serriola	FACU
Linum perenne	
Medicago lupulina	FAC
Melilotus alba	FACU
Melilotus officinale	FACU
Myosotis laxa	OBL
Phalaris arundinacea	FACW
Phleum pratense	FAC-
Plantago lanceolatum	FACU+
Plantago major	FAC+
Poa pratens is	FAC
Polygonum convolvulus	FACU-
Populus balsamifera	FAC
Potamogeton natans	OBL
Potamogeton pectinatus	OBL
Potentilla anserina	OBL
Ranunculus aquatilis	OBL
Ranunculus sceleratus	OBL
Rumex crispus	FACW
Silene latifolia	
Sparganium emersum	OBL
Taraxacum officinale	FACU
Thlaspi arvense	NI
Tragopogon dubius	UPL
Trifolium hybridum	FACU+
Trifolium pratense	FACU
Typha latifolia	OBL
	1 *
Verbascum thapsus	UPL



#### 3.3 Soils

According to the Upper Flathead Valley Area soil survey (Soil Conservation Service 1960), soils in the mitigation site are classified as poorly drained alluvial land and (Aa) and the Swims silt loam (So). The poorly drained alluvial land soil has poor surface and internal drainage, mottling in the subsurface and typically consists of loam or silty loam. The Swims soil consists of silt loam and tends to occupy low terraces occupying the Flathead River.

These characteristics were generally confirmed during monitoring. Three test pits were excavated and described using the ACE routine wetland monitoring form. The TP1 located adjacent to the pond consisted of 16-inches of organic detritus overlying a mottled silt loam. Hydric soil characteristics are well developed including a histic epipedon. TP2 is classified as a poorly developed hydric soil. A thin (1-inch) layer of organic detritus was present. A low-chroma (7.5 YR 2.5/2) A-horizon is present from 1 to 9-inches and mottles were observed below 9-inches. These soil characteristics indicate an oxygen-depleted environment with a fluctuating water table. TP3 is a loam representative of the upland soil, which did not exhibit hydric characteristics in the A horizon (7.5 YR 2.5/2) or B horizon (7.5 YR 4/3).

#### 3.4 Wetland Delineation

Delineated wetland boundaries are illustrated on **Figure 3**. Completed wetland delineation forms are included in **Appendix B**. Soils, vegetation, and hydrology are discussed in preceding sections. Delineation results indicate a total of 5.2 acres of wetland.

The original mitigation goal was to enhance two acres of existing wetland and create four acres for a total of six acres. It is likely that area within the Type 3 Community and within the ditches will develop hydric soil characteristics with continued inundation.

## 3.5 Wildlife

Wildlife species, or evidence of wildlife, observed on the site during 2001 monitoring efforts are listed in **Table 2**. Specific evidence observed and activity codes pertaining to birds are provided on the completed monitoring form in **Appendix B**. Two mammal and numerous bird species were noted using the mitigation site.



7

Table 2: Fish and Wildlife Species Observed on the Creston Mitigation Site

<u> </u>	
FISH	
none	
AMPHIBIANS	
None observed	
REPTILES	
None observed	
BIRDS	Pintail (Anas acuta)
American Robin (Turdus migratorius)	Red-winged Blackbird (Agelaius phoeniceus)
Canada Goose (Branta Canadensis)	Ring-necked duck (Aythya collaris)
Cinnamon Teal (Anas cyanoptera)	Ring-necked Pheasant (Phasianus colchicus)
Common Goldeneye (Bucephala clangula)	Spotted Sandpiper (Actitis macularia)
Killdeer (Charadrius vociferous)	Tree Swallow (Tachycineta bicolor)
Mallard (Anas platyrhynchos)	Wood Duck (Aix sponsa)
Northern Shoveler (Anas clypeata)	Osprey (Pandion haliaetus)
MAMMALS	
White-tailed Deer (Odocoileus virginianus)	
Meadow vole ( <i>Microtus pennsylvanicus</i> )	

#### 3.6 Macroinvertebrates

Macroinvertebrate sampling was not conducted due to lack of sampling equipment, but will be conducted in May and July of 2002 and in subsequent monitoring years.

## 3.7 Functional Assessment

Completed functional assessment forms are presented in **Appendix B**. Functional assessment results are summarized in **Table 3**. The site was evaluated as a single assessment area and rated as a Category II wetland. Wildlife habitat and groundwater discharge are the primary functions of the site. The site provides a total of 35.4 functional units and achieved 76% of possible points.

A functional assessment was not conducted prior to site construction and therefore cannot be used for comparison. Future assessments will be compared to the 2001 functional assessment.

# 3.8 Photographs

Representative photos taken from photo-points and transect ends are provided in **Appendix C**.

#### 3.9 Maintenance Needs/Recommendations

The berm was in good condition during the spring and mid-season visits. We have no recommendations at this time. The bird boxes also appeared to be in good condition.

# 3.10 Current Credit Summary

Approximately 5.2 acres of wetlands are currently present on the mitigation site. Based on preconstruction goals two acres were to be enhanced and four acres created for a total of 6 acres.



The existing acreage is close to the goal. Based on site conditions it is expected that additional wetland acres will develop in the future with continued inundation.

Table 3: Summary of 2001 Wetland Function/Value Ratings and Functional Points <sup>1</sup> at the Creston Mitigation Project

Function and Value Parameters From the 1999 MDT Montana Wetland Assessment Method	Wetland Numbers
Listed/Proposed T&E Species Habitat	Mod (0.7)
MNHP Species Habitat	Low (0.1)
General Wildlife Habitat	High (0.9)
General Fish/Aquatic Habitat	NA
Flood Attenuation	NA
Short and Long Term Surface Water Storage	High (0.8)
Sediment, Nutrient, Toxicant Removal	Mod (0.7)
Sediment/Shoreline Stabilization	NA
Production Export/Food Chain Support	High (0.9)
Groundwater Discharge/Recharge	High (1)
Uniqueness	Mod (0.6)
Recreation/Education Potential	High (1)
Actual Points/Possible Points	6.8 / 9
% of Possible Score Achieved	76%
Overall Category	II
Total Acreage of Assessed Wetlands within Easement	5.2 ac (calculated)
Functional Units (acreage x actual points)	35.4 fu
Net Acreage Gain	NA
Net Functional Unit Gain	NA
Total Functional Unit "Gain"	NA

# 4.0 REFERENCES

- Carlson, J. Program Zoologist, Montana Natural Heritage Program. Helena, MT. April 2001 conversation with Jeff Berglund.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. US Army Corps of Engineers. Washington, DC.
- Ralph, C.J., Geupel, G.R., Pyle, P., Martin, T.E., and D.F. DeSante. 1993. *Handbook of field methods for monitoring landbirds*. Gen. Tech. Rep. PSW-GTR-144. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Dept. of Agriculture. 41 p.



- Reed, P.B. 1988. National list of plant species that occur in wetlands: North West (Region 9). Biological Report 88(26.9), May 1988. U.S. Fish and Wildlife Service. Washington, D.C.
- Soil Conservation Service. 1960. Soil survey of Upper Flathead Valley Area, Montana. Series 1946, No. 4.
- USDA Natural Resources Conservation Service. 1998. *Field Indicators of Hydric Soils in the United States*, Version 4. G. Hurt, P. Whited and R. Pringle (eds.). USDA, NRCS Fort Worth, TX.
- Werner, K. Herpetologist, Salish-Kootenai Community College. Pablo, MT. May 1998 instructional presentation (per Jeff Berglund).





# Appendix A

# FIGURES 2 - 3

MDT Wetland Mitigation Monitoring Creston Creston, Montana







# Appendix B

COMPLETED 2001 WETLAND MITIGATION SITE MONITORING FORM
COMPLETED 2001 BIRD SURVEY FORMS
COMPLETED 2001 WETLAND DELINEATION FORMS
COMPLETED 2001 FIELD AND FULL FUNCTIONAL
ASSESSMENT FORMS

MDT Wetland Mitigation Monitoring Creston Creston, Montana



# DRAFT - MDT WETLAND MITIGATION SITE MONITORING FORM

Project Name: MDT Month ing Project Number: 130091.00% Assessment Date: 7 /25	101							
Location: Creston MDT District: Milepost:								
Legal description: T R Section Time of Day: Atendo								
Weather Conditions: clear 80°, Smph to cast Person(s) conducting the assessment: A Kahle								
Initial Evaluation Date: 5 /8 /01 Visit #: 2 Monitoring Year: 2001								
Size of evaluation area: 20 acres Land use surrounding wetland: agriculture + wibon								
THE POLOCIAL								
HYDROLOGY								
Surface Water								
Inundation: Present X Absent Average depths: ft Range of depths: - ft								
Assessment area under inundation: 10-19%								
Depth at emergent vegetation-open water boundary:> ¬ ft								
If assessment area is not inundated are the soils saturated w/in 12" of surface: Yes No								
Other evidence of hydrology on site (drift lines, erosion, stained vegetation etc.):								
Outer evidence of hydrology on site (drift lines, crosion, stanted vegetation etc.).								
Groundwater								
Monitoring wells: Present X Absent - D.d not have key as of 7/25/01								
Record depth of water below ground surface								
Well #   Depth   Well #   Depth   Well #   Depth								
Well # Deput Well # Deput Well # Deput								
4.3.31d								
Additional Activities Checklist:								
Map emergent vegetation-open water boundary on air photo								
Observe extent of surface water during each site visit and look for evidence of past surface water								
elevations (drift lines, erosion, vegetation staining etc.)								
GPS survey groundwater monitoring wells locations if present								
COMMENTS/PROBLEMS: Obtained key from lary Whom and visited site aga	1							
on 8-25-01. Ky did not fit. I wrote down lock the and bu	<u> </u>							
will attempt to didir a key								

VECET.	ATION	COMN	<b>MUNITIE</b>	S



Dominant Species	% Cover	Dominant Species	% Cove
LUREP		Leaume	
HLPRA	•	DACGLO	
DAPRA		TAROFF	
INPER		MEDLUP	
RARV			
COMMENTS/PROBLEMS:			
Dominant Species	le (main species):	-	% Cov
TYPLAT		CEPDEM (inwater)	507
PHARRU	30%	SPA (inwater)	17
AWPRA .	5%	JUND	17
ELEPAL		EPICIL	T
AGRSTO	5%		
COMMENTS/PROBLEMS:			
		depressions	
			% Cov
community No.: 3 Community Tit	le (main species):		% Cov
Community No.: 3 Community Tit	le (main species):		% Cov
Community No.: 3 Community Tit	le (main species):		% Cov
Community No.: 3 Community Tit	le (main species):		% Cov
Community No.: 3 Community Tit	le (main species):		% Cov
Community No.: 3 Community Tit  Dominant Species  YPLAT	le (main species):	Dominant Species	% Cov
Community No.: 3 Community Tit  Dominant Species  YPLAT	le (main species):	Dominant Species	% Cov
Community No.:_3_ Community Tit  Dominant Species  YPLAT	le (main species):	Dominant Species	% Cov
Community No.:_3_ Community Tit  Dominant Species  YPLAT	le (main species):	Dominant Species	% Cov
Community No.: 3 Community Tit	le (main species):	Dominant Species	% Cov



# **VEGETATION COMMUNITIES - CRESTON**

Dominant Species	% Cover	Dominant Species	% Cover
Elymus repens	20%	Linum perenne	5%
Phleum pratense	10%	Trifolium hybridum	5%
Agrostis stolonifera	10%	Taraxacum officinale	5%
Cirsium arvense	10%	Medicago lupulina	5%
Yellow and purple legumes combined	10%	Poa pratensis	2%
COMMENTS/PROBLEMS:			
Community No.: 2 Community Title	(main species)	: Typha latifolia – pond edges	-
Dominant Species	% Cover	Dominant Species	% Cover
Typha latifolia	50%	Juneus articulatus	1%
Phalaris arundinacea	30%	Enilohium ciliatum	Т
	3070	Ephobium chiatum	Trace
	10%	Epilobium ciliatum  Ceratophyllum demersum (in water)	50%
Eleocharis palustris	10%	Ceratophyllum demersum (in water)	
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera	10% 5% 5%	Ceratophyllum demersum (in water) Sparganium emersum (in water)	50%
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:	10% 5% 5%	Ceratophyllum demersum (in water) Sparganium emersum (in water)	50%
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:  Community No.: 3 Community Title	10% 5% 5% (main species)	Ceratophyllum demersum (in water)  Sparganium emersum (in water)  :_Depressions: mixed Typha latifolia and	weedy grasse
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:  Community No.:3 Community Title  Dominant Species	10% 5% 5% (main species)	Ceratophyllum demersum (in water)  Sparganium emersum (in water)  : Depressions: mixed Typha latifolia and  Dominant Species	weedy grasse
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:  Community No.:3 Community Title  Dominant Species Typha latifolia	10% 5% 5% (main species)	Ceratophyllum demersum (in water)  Sparganium emersum (in water)  :_Depressions: mixed Typha latifolia and	weedy grasse
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:  Community No.:3 Community Title  Dominant Species Typha latifolia Phalaris arundinacea	10% 5% 5% 5% (main species) % Cover 15%	Ceratophyllum demersum (in water)  Sparganium emersum (in water)  :Depressions: mixed Typha latifolia and  Dominant Species  Medicago lupulina	weedy grasse  % Cover 3%
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:  Community No.:3 Community Title  Dominant Species Typha latifolia Phalaris arundinacea Agrostis stolonifera	10% 5% 5% 5% (main species) % Cover 15% 10% 10%	Ceratophyllum demersum (in water)  Sparganium emersum (in water)  :Depressions: mixed Typha latifolia and  Dominant Species  Medicago lupulina  Populus balsamifera	50%   1%
Eleocharis palustris Alopecurus pratensis Agrostis stolonifera  COMMENTS/PROBLEMS:  Community No.:3 Community Title	10% 5% 5% 5% (main species) % Cover 15% 10%	Ceratophyllum demersum (in water)  Sparganium emersum (in water)  : Depressions: mixed Typha latifolia and  Dominant Species  Medicago lupulina  Populus balsamifera  Plantago major	50%   1%

Additional Activities Checklist:
\_\_\_\_\_\_\_ Record and map vegetative communities on air photo

# **VEGETATION COMMUNITIES - CRESTON**



Dominant Species	% Cover	Dominant Species	% Cover
Phalaris arundinacea	95%	Juneus articulatus	trace
Agrostis stolonifera	trace	Chenopodium rubrum	trace
Equisetum arvense	trace	Cirsium arvense	trace
Carex bebbii	trace	Veronica americana	trace
Eleocharis palustris	1%	Plantago major	trace
COMMENTS/PROBLEMS:			
Community No.:5 Community Ti			
Dominant Species	% Cover	A 14 H 10 T 10	% Cover
Potamageton pectinatus	98%	Plantago major	trace
Ranunculus aquatilis	trace	Chenopodium rubrum	1%
Sparganium emersum	trace	Phalaris arundinacea	5%
Barbarea vulgaris	trace		
Eleocharis palustris	trace	<u> </u>	
COMMENTS/PROBLEMS:			
Community No.: 6 Community Ti	itle (main species)	: Alopecurus pratensis	
Community No.: 6 Community Ti	% Cover	Dominant Species	
Dominant Species Alopecurus pratensis	% Cover 70%	Dominant Species Agrostis stolonifera	% Cover
Dominant Species Alopecurus pratensis Phalaris arundinacea	% Cover 70% 10%	Dominant Species Agrostis stolonifera Taraxacum officinale	
Dominant Species Alopecurus pratensis Phalaris arundinacea Cirsium arvense	% Cover 70% 10% 1%	Dominant Species Agrostis stolonifera	trace
Dominant Species Alopecurus pratensis Phalaris arundinacea Cirsium arvense Medicago lupulina	% Cover 70% 10%	Dominant Species Agrostis stolonifera Taraxacum officinale	trace trace
	% Cover 70% 10% 1%	Dominant Species Agrostis stolonifera Taraxacum officinale	trace trace

**Additional Activities Checklist:** 

Record and map vegetative communities on air photo

# PLANTED WOODY VEGETATION SURVIVAL AND & WATER B.5



Species brown dots	Number Originally Planted	Number Observed	Mortality Causes
Silver Burny 8=	27	15	unknown competition as weeks
=9 partland sturberry	6	27	unknow! competition if weeks lack
Villa collin OI	27	23.13	save as about
=15 wilow silvabing	153	₹ 80	braveing + thistle competition
rose	2	00	alousing a missia contention
seinted haf and chakether			
4 11	24	23	severe browsing
# 12+13 Chokochemy of patrons	184	141	boowsing weed composition
+14 37	29		esouthistic howk or re
26 chokechemy			CAN TOTAL TO TOWN ST TO
#1 Apple trees	16	14	booksing
# Z Apply trees	20	6	possi, vo
# 3 Words 1088	SS	45	wied competition
# 4 Woods rose	25	23	weed competting browsing
# 6 Woods nose	113	113	none - weeds are competho
1-7 Silver berry	- 85	78	MRUMU - MERRY WE ERROLLING
T STOCK BOLLY	1 23	10	MICHON,
			<del> </del>
		is .	
		**	
	-		
OMMENTS/DDODLEMS.			-
OMMENTS/PROBLEMS:			
Houndstooth : nov. Dis	3.0-1		
- 10 Kennel 1040 . 110 K 1412 .	300 4	·	



## **BIRD SURVEY - FIELD DATA SHEET**

SITE: Custon

Page | of | Date: 5-8-01 + 5-9-01

Survey Time: 3:40 pm

~ Jam

Bird Species	Behavior	Habitat Type	Bird Species	Behavior	Habitat Type
Wood Duck	Pair				
Cinnamon Teal	Pair				
Common Goldeneye	Pair		20.00 ON 10.00 ON 10.00 ON		
Hortlein Shoulor	2 Pair	11 (190) V			
Canada Goose	RP				
true smallows		FO			
American Robin					
starlings				l	
Killderr	200				
mallards					
Goldeneye	Park				
Red wired black bods					
		-			
OSpecey					
cinnon real					
Shrasen+ (runner)					
				ļ	
				ļ	
				<u> </u>	, , , , , , , , , , , , , , , , , , ,
		$\vdash$			
	100				

OTES: Mark + And	nka visited site on afternoon of 5.8-01
Mark returned +	o site on morning of 5-9-01.
- 4	,
This form was	not yet developed and thurdre observations
were morded	in field natebooks.
	<del></del>

**Behavior:** BP - one of a breeding pair; BD-breeding display; F - foraging; FO - flyover; L - loafing; N - nesting **Habitat:** AB - aquatic Bed; FO - forested; I - Island; MA - marsh; MF: Mud Flat; OW - open water; SS - scrub-shrub; UP - upland buffer; WM - wet meadow



## **BIRD SURVEY - FIELD DATA SHEET**

SITE: Creston

Page | of | Date: 7-25-0 | Survey Time: ALLINOON

Bird Species	Behavior	Habitat Type	Bird	Species	Behavior	Habitat Type
Kildeer red-winged blackbird Swellow (ip)	F	M M OW/MA				
red-winded blackbard	_	AM WO				
Swallow (:e)	FO	WP	200000000000000000000000000000000000000			
1.()						
				90 909		7999
					201	(a.c.)
	era angre					
					.'.	
				1000		
			N N N N N N N N N N N N N N N N N N N	400 - 50 - 5000 500		
					1,000	
						100
			1 1/10/2009 11/10/10/2009			
			201727F3 68			100
						100
				59		
			1 1 10 10 10			
				12.00		
NOTES:				NAMES OF THE PARTY		

Behavior: BP - one of a breeding pair; BD-breeding display; F - foraging; FO - flyover; L - loafing; N - nesting Habitat: AB - aquatic Bed; FO - forested; I - Island; MA - marsh; MF: Mud Flat; OW - open water; SS - scrub-shrub; UP - upland buffer; WM - wet meadow



# CRESTON WETLAND MITIGATION SITE MONITORING INFORMATION

# Bird Species observed since 1997: (B) Nest observed on site (BB) broods and/or young observed

Bird Species	Pre - Monitor	Monitor 1998	Monitor 1999	Monitor
Observed:	1997			2000
American Robin	X	X	X	X
Common snipe	X	X (B)	X (B)	X(B)
Magpie	X	X	X	X
Barn swallow	X	X	X	X
Killdeer	X	X(BB)	X(BB)	X
Vesper sparrow	X	X	X(B)	X
Song sparrow	X	X	X	
Eastern kingbird	X	X	X	X(B)
Goldfinch	X	X	X	
Wood duck	X	X(BB)	X(BB)	X(BB)
Grt. horned owl	X			
Mallard	X	X(BB)	X(BB)	X(BB)
RW blackbird	X	X(B)	X(B)	X(B)
Sol. sandpiper		X(B)	X(B)	X(B)
Spot. sandpiper	2602000	X(BB)	X(BB)	X(B)
YH blackbird			X(B)	X(B)
Bank swallows		X	X	X
Goldeneye		X	X	X(B)
Bufflehead		X(BB)	X(BB)	X(BB)
Redhead duck			X	X
Wls phalarope		X(B)	X	X(BB)
Sav. Sparrow	100000	X(B)		X
ViGr.swallows		X	X	
Gr. winged teal			X	X
Cinnamon teal			X(BB)	X
Blue-winged teal			X	X
Canada goose			X(BB)	X(BB)
Pintails	10.000		X	X
Wigeon			X	
Marsh wrens			X(B)	X(B)
Ruddy duck			X	X
Sora rail			X	X
American coot			X(BB)	X(BB)
AM. Shovellers	1000 00 00 Mg (1000 000		X	X
Brw. blackbirds			X	X
Bobolink				X
Turkey				X

		-	_
wii	m	16	L
w			r



				F	BIRDS					
Species	Number Observed	Nesting or Breeding Activity	Likely Breeding Resident	Likely Migrating	Sp	ecies	Number Observed	Nesting or Breeding Activity	Likely Breeding Resident	Likely Migrating
				Ļ	_					
					4					
					<b>√</b>					+
				<del> </del>	<u> </u>					
				<del> </del>	11					-
				-	+			<b></b>		1
					┨├ <del>──</del>	AN				1
					┨ ├───	142				+
	_			<del>                                     </del>	-	_				
					1					
			· · · · · · · · · · · · · · · · · · ·		1					
							1			
Were man ma structures bein	ing dulinzed:	105					•			
	Spec	iae	MAN	MALS	AND HERI Number	TILES	Indirect	indication	of use	
	Spec	103		- 1	Observed	Tracks	Sca		rows	Other
deer						X				
dog - co	Lava					X				
3	7									
								_		
							_	-		
Additional Activities Checklist:										
Macroinvertebrate sampling (if required)										
common and the second second and the second										
COMMENTS/PROBLEMS: D.d not how D.not to sample macro-invertebrates										
	-									

# **PHOTOGRAPHS**

Using a camera with a 50 mm lenses and color film take photographs of the following permanent reference points listed in the checklist below. Record the direction of the photograph using a compass. (The first time at each site establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3' above ground, survey the location with a resource grade GPS and mark the location on the air photo.)

Checklist:

	One photo for each of the 4 cardinal directions surrounding wetland
	At least one photo showing upland use surrounding wetland - if more than one
	upland use exists, take additional photos
$\times$	At least one photo showing buffer surrounding wetland
X	One photo from each end of vegetation transect showing transect

Location	Photo Frame #	Photograph Description	Compass Reading
AI	15.16	Phatopoint 1	N + S
B 2	17.18	2	SW, NE
€ 3	19 20	, 3	E, W
<b>D</b> 4	21,22	· 4	NS.
E S	1,12,13 14	. 5	N.S.E. W
P L	9,10	'' b	W, E
Ø	, , , , , , , , , , , , , , , , , , , ,		
H			

COMMENTS/PROBLEM	<b>1</b> S:	
I'm transit F.	Ks soling tact	
)	X laoken west	
	x1 - 10 420 0x =	
	98- booking east	
	2	

# GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points with the GPS unit set at 5 second recording rate. Record file numbers fore site in designated GPS field notebook

# Checklist:

Y Jurisdictional wetland boundary	
* 4-6 landmarks recognizable on the air photo	mints
Start and end points of vegetation dansect(s)	Qu
Y Photo reference points	
— Groundwater monitoring well locations	
COMMENTS/PROBLEMS:	



# WETLAND DELINEATION

At each site conduct the items on the checklist below:						
Delineate wetlands according to the 1987 Army Corps manual.						
X_ Survey wetland-upland boundary with a resource grade GPS survey						
COMMENTS/PROBLEMS:						
FUNCTIONAL ASSESSMENT						
Collect information to complete MDT Function and Values Assessment in the office.						
See forms						
See forms						
COMMENTS/PROBLEMS:						
MAINTENANCE						
Were man-made nesting structures installed at this site? YESX NO						
If yes, do they need to be repaired? YESNO						
If yes, describe problems below and indicate if any actions were taken to remedy the problems.						
Were man-made structures build or installed to impound water or control water flow into or out of the wetland?						
YES_X_NO If yes, are the structures working properly and in good working order? YES_X_NO						
If no, describe the problems below.						
COMMENTS/PROBLEMS:						
COMMENTS/PROBLEMS:						



# DATA FORM ROUTINE WETLAND DETERMINATION (1987 COE Wetlands Delineation Manual)

Project/Site:		Date: 8-24-01 County: Flatherd State: MT			
Do Normal Circumstances exist on the site? Is the site significantly disturbed (Atypical Situals the area a potential Problem Area? (If needed, explain on reverse.)	Community ID: 5 Transect ID: Plot ID:				
VEGETATION					
Dominant Plant Species  1. Potamogeton pectrals # DBL  2.  3.  4.  5.  6.  7.  8.  Percent of Dominant Species that are OBL, FACW or FAC (excluding FAC-).  Remarks: Dead tree observed in standing	9	Stratum Indicator			
HYDROLOGY	<b>Y</b>				
Recorded Data (Describe in Remarks):Stream, Lake, or Tide GaugeAerial PhotographsOther	Water Mar Drift Lines Sediment (	in Upper 12 Inches ks			
Depth of Surface Water:  Depth to Free Water in Pit:  Depth to Saturated Soil:  (in.)	Secondary Indicators  Oxidized R  Water-Stai  Local Soil : FAC-Neutr	s (2 or more required): loot Channels in Upper 12 Inches ined Leaves Survey Data			
Remarks;					



# SOILS

Map Unit Name (Series and Phase):	L + Peet 11960	Field Obse			
(Inches)   Horizon (Muns	x Color Mottle Colors (Munsell Moist)  R 25/1 NA  5/R 6/1 7.5/8 /3	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.  Muck  B.H. Loan		
Hydric Soil Indicators:					
This area is typically indudated 100 water year has decreated surface water elevation					

# WETLAND DETERMINATION

Hydrophytic Vegetation Present? Wetland Hydrology Present? Hydric Soils Present?	No (Circle) (68) No (768) No	Is this Sampling Point Within a Wetland?	(Circle)
Romarka: Soil pit is located to read-congry gra	along vigitation	on transect immediately	y adjacent

Approved by HQUSACE 3/92



# DATA FORM ROUTINE WETLAND DETERMINATION [1987 COE Wetlands Delineation Manual]

Project/Site: Custon Applicant/Owner: MOT Investigator: A. Kuhlu				Date: 8-24-01 County: Flathead State: MT			
Do Normal Circumstances e Is the site significantly distu Is the area a potential Proble (If needed, explain on rev	Community ID: 4 Transect ID: Plot ID: 2						
VEGETATION							
Dominant Plant Species	Stratum	Indicator	Dominant Plant Species	Stratum Indicator			
1. reed concry grass							
2			ı				
3			11				
4			1				
5			13				
6,			14				
7			15				
8		·	16				
Percent of Dominant Species that are OBL, FACW or FAC (excluding FAC-).							
Remarks:							
HYDROLOGY							
<ul> <li>Recorded Data (Describe in Remarks):</li> <li> Stream, Lake, or Tide Gauge</li> <li> Aerial Photographs</li> <li> Other</li> <li> No Recorded Data Available</li> </ul>			Wetland Hydrology Indicators: Primary Indicators: Inundated Saturated in Upper 12 Inches Water Marks Drift Lines				
Field Observations:				Patterns in Wetlands (2 or more required):			
Depth of Surface Water:	0	(in.)	X Oxidized R	oot Channels in Upper 12 inches			
Depth to Free Water in Pit:		(in.)	Local Soil	ned Leaves Survey Date			
Depth to Saturated Soil:	>14	(in.)	FAC-Neutr Other (Exp	al Test Isin in Remarks)			
Remarks:							



#### SOILS

	Name d Phase): (Subgroup):	ta (poorly due	ined Allewial			
Profile Des Depth (inches)	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.	
1-9 9216	A B	7.54R2.5	NA 2548 5/3	N/A Common/do	loam t. silt loam	
Hydric Soil Indicators:  Histosol Histic Epipedon Sulfidic Odor Aquic Moisture Regime Reducing Conditions Gleyed or Low-Chroma Colors  Concretions High Organic Content in Surface Layer in Sandy Soils Organic Streaking in Sandy Soils Listed on Local Hydric Soils List Listed on National Hydric Soils List Other (Explain in Remarks)						
Remarks:	Hplr.c	en zi hoz	t wall de	ua lagre mi		

#### WETLAND DETERMINATION

Hydrophytic Vegetation Present? Wetland Hydrology Present? Hydric Soils Present?	Yes No		is this Sampling Paint Within a Wetland?	(Circle)
Remarks:		4		

\* Soil in low area by sparse regetation is similar soil however, lacking organic accumulation

# DATA FORM **ROUTINE WETLAND DETERMINATION**



(1987 COE Wetlands Delineation Manual)

Project/Site:	Date: 8-24-01					
(If needed, explain on reverse.)						
VEGETATION  Dominant Plant Species Stratum Indicator Dominant Plant Species Stratum Indicator						
1 Elymus Regards H FACU 9 2 Phillum pratence H FAC- 10 3 Agnostic sobjected H FACH 11 4 Circum evence H FACH 12 5 Legumus (sp?) H ? 13 6 15 8 16  Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). 25% Remarks: Ligume species (yellow) are planted and not identified.						
HYDROLOGY						
Recorded Data (Describe in Remarks):	Wetland Hydrology Indicators:					
Stream, Lake, or Tide Gauge	Primary Indicators:					
Aerial Photographs	Inundated					
Other	Saturated in Upper 12 Inches					
No Recorded Data Available	Water Marks					
	Drift Lines					
Field Observations:	Sediment Deposits					
Depth of Surface Water: (in.)  Depth to Free Water in Pit: (in.)	Drainage Patterns in Wetlands Secondary Indicators (2 or more required):  Oxidized Root Channels in Upper 12 Inches Water-Stained Leaves					
Depth to Saturated Soil: (in.)	Local Soil Survey Data FAC-Neutral Test Other (Explain in Remarks)					
Remarks:						



## SOILS

Map Unit Name (Series and Phase): Swims sitted Taxonomy (Subgroup): Confirm Mapped Type? Yes X No							
			Mottle Colors	0.00.00		Texture, Concretions,	
inches	Horizon	(Munsell Moist)	(Munsell Mois	st) Abu	ndance/Contrast	Structure, etc.	
0-1	0					<u></u>	
1-5	A	7.5/R 2.5/2	NA	N	/A	loam	
5-116	₹,	7.51R 43	N/A	- 1	1/4	loam	
	i .						
Histic Epipedon Sulfidic Odor Aquic Moisture Regime Reducing Conditions Gleyed or Low-Chroma Colors  Remarks: Soil IS Juy Juy Shows Signs of axidation button ~ 10'							
		RMINATION					
Wetland	Hydrophytic Vegetation Present?						
Remark	ks:		1				

Approved by HQUSACE 2/92

Date: 8/24/01 By: 16/1

Field Data Sheet for 1999 MDT Wetland Assessment Form Site: Crestor Estimated AA Size (Circle Ac.): <1 1-5 >5 **Brief Description:** HGM Class (CIRCLE) Cowardin Class Est. % Predominant Water Regime (CIRCLE) of AA Mineral Soil Flats Emergent Perm Flood Int Exp Sem Perm Flood Seas Flood Tem Flood Int Flood Organic Soil Flats Tem Flood Int Flood Aquatic Bed Perm Flood (Int Exp ) Sem Perm Flood Seas Flood Sat Riverine (nonperennial) Riverine (upper perennial) Moss-Lichen Perm Flood Int Exp Sem Perm Flood Seas Flood Sat Tem Flood Int Flood Riverine (lower perennial) Scrub-Shrub Sat Perm Flood Int Exp Sem Perm Flood Seas Flood Tem Flood Int Flood Lacustrine Fringe Depression (closed) Int Flood Forested Perm Flood Int Exp Sem Perm Flood Sat Tem Flood Seas Flood Depression (open, groundwater) Unconsolidated Bottom Perm Flood Int Exp Sem Perm Flood Seas Flood Sat Tem Flood Int Flood Depression (open, surface water) Other: Perm Flood Int Exp Sem Perm Flood Seas Flood Tem Flood Int Flood Slope Organic Soil Flats Total Estimated % Vegetated RELATIVE ABUNDANCE: rare com. (abun.) DISTURBANCE is: High Moderate Low HYDROLOGY: Max. acre-ft surf. water at wetlands in AA subject to inundation: <1 1-5 >5 (if no flooding/ponding, go to groundwater\* section) N If outlet present, is it restricted (subsurface will always be "yes")? Y Ν Does AA contain surface or subsurface outlet? Longest duration of surface water: Surface Water Duration and other attributes (circle) at any wetlands within AA Perm/Peren Seas / Intermit Temp / Ephem Perm Deren Seas / Intermit Temp / Ephem in at least 10% of AA (both wetlands and nonwetlands [deepwater, streambed...] Where fish are or historically were present (circle NA) if not applicable) Perm / Peren Seas / Intermit Temp / Ephem >25% 10-25% <10% % of waterbody containing cover objects €50% >75% 50-74% % bank or shore with riparian or wetland shrub or forested communities Temp / Ephem adjacent to rooted wetland vegetation along a defined watercourse or shoreline subject to wave Perm / Peren Seas / Intermit action (circle NA)f not applicable) (35-64% <35% % cover of wetland bank or shore by sp. with binding rootmasses >65% Flood Attenuation: Do any wetlands on site flood as a result of in-channel or overbank flow? Y N) (if no, go to groundwater\* section below) 2-10 <2 Estimated wetland area subject to periodic flooding (acres): ≥10 <25 Estimated % of flooded wetland classified SS, FO or both: ≥75 25-74 \*Evidence of groundwater discharge or recharge?Y List: HABITAT Habitat for Listed or Proposed Threatened, Endangered, or Montana Natural Heritage Program S1, S2, or S3 Plants or Animals: AA is Documented (D) or Suspected (S) to contain (circle based on defiltions contained in instructions): Primary or critical habitat (list species) D S T/E: D S MNHP: D S MNHP: D S Secondary habitat (list species) T/E: incidental habitat (list species) D S T/E: D S MNHP: No usable habitat D S T/E: D S MNHP: Wildlife observations? Fish observations? OTHERS Do wetlands have potential to receive excess sediments, nutrients, or toxicants Y From: On TMDL List? low to moderate levels high levels Potential to receive: Does site contain bog, fen, warm springs, >80 year-old forested wetland, or MNHP "S1" or "S2" plant association? List: Is AA a known recreation / education site? N Type: Does AA offer strong potential for use as recreation + education site? N Type:



MDT Montana Wetland Assessment Form (revised 5/25/1999) 2. Project #: 130091.007 3. Evaluation Date: Mo. 8 Day 24 Yr. 01 4. Evaluator(s): Kwhle 5. Wetlands/Site #(s) 6. Wetland Location(s): I. Legal: T \_\_\_ N or S; R \_\_\_\_\_ E or W; S \_\_\_\_\_\_; T N or S; R \_\_\_ E or W; S \_\_\_ ii. Approx. Stationing or Mileposts: iii. Watershed: Flathuad - 04 GPS Reference No. (if applies):\_\_\_ Other Location Information: 7. a. Evaluating Agency: LWC/MDT 8. Wetland size: (total acres) \_\_\_\_ (visually estimated) b. Purpose of Evaluation: (measured, e.g. by GPS [if applies]) Wetlands potentially affected by MDT project Mitigation wetlands; pre-construction 2. Assessment area: (AA, tot., ac., (visually estimated) Mitigation wetlands: post-construction see instructions on determining AA) (measured, e.g. by GPS [if applies]) Other 10. Classification of Wetland and Aquatic Habitats in AA (HGM according to Brinson, first col.; USFWS according to Cowardin [1979], remaining cols.) **HGM Class** System Subsystem Class Water Modifier % of AA Regime Palustrine AB LOOK OFFICE TOO 6 EW (Abbreviations: System Palustrine(P) Subsyst.: none/ Classes: Rock Bottom (RB ), Unconsolidated bottom (UB ), Aquatic Bed (AB), Unconsolidated Shore (US ), Moss-lichen Wetland (ML), Emergent Welland (EM), Scrub-Shrub Welland (SS), Forested Welland (FO) System: Lacustrine (L), Subsyst.: Limnetic (2)/ Classes: RB, UB, AB/ Subsystem: Littoral (4)/ Classes: RB, UB, AB, US, EM/ System: Flivering (B) Subsyst.: Lower Perennial (2) Classes: RB, UB, AB, US, EM Subsystem: Upper Perennial (3) Classes: RB, UB, AB, US Water Regimes; Permanently Flooded (H), Intermittently Exposed (G), Semipermanently Flooded (F), Seasonally Flooded (C), Saturated (B), Temporarily Flooded (A), Intermittently Flooded (J) Modifiers: Excavated (E), Impounded (I), Diked (D), Partly Drained (PD), Farmed (F), Artificial (A) HGM Classes: Riverine, Depressional, Slope, Mineral Soil Flats, Organic Soil Flats, Lacustrine Fringe 11. Estimated relative abundance: (of similarly classified sites within the same Major Montana Watershed Basin, see definitions) (Circle one) Unknown Rare Abundant Common Comments: 12. General condition of AA: i. Regarding disturbance: (use matrix below to determine [circle] appropriate response) Conditions within AA Predominant conditions adjacent to (within 500 feet of) AA Land not cultivated, but moderately Land managed in predominantly Land cultivated or heavily grazed or looged: natural state; is not grazed, haved, grazed or hayed or selectively logged; subject to substantial fill placement, grading, logged, or otherwise converted: or has been subject to minor clearing; clearing, or hydrological atteration; high road contains few roads or buildings. does not contain roads or buildings or building density. AA occurs and is managed in predominantly natural state; is not low disturbance low disturbance moderate disturbance grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings. AA not cultivated, but moderately grazed or hayed or selectively moderate disturbance moderate disturbance high disturbance logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings. AA cultivated or heavily grazed or logged; subject to relatively high disturbance high disturbance high disturbance substantial fill placement, grading, clearing, or hydrological alteration; high road, or building density. Comments: (types of disturbance, intensity, season, etc.): most plants are introduced, disturbance is now missing ii. Prominent weedy, alien, & introduced species (including those not domesticated, feral); (list) ... 2 open whiter areas. Adjacent lii. Provide brief descriptive summary of AA and surrounding land use/habitat: ક્લાયન નિરાદ્રે છે to 2-love that, our age transfer sto and agricult 13. Structural Diversity: (based on number of "Cowardin" vegetated classes present (do not include unvegetated classes), see #10 above). # of "Cowardin" vegetated classes present in AA (see #10) 2 vegetated classes ≥ 3 vegetated classes (or ≤ 1 vegetated class ≥ 2 if one is forested) (or 1 if forested) Rating (circle) Moderate Low Comments:



		SECTION PERTAINING to FUNCTIONS & VALUES ASSESSMENT																		
14A. Habitat for Federally  AA is Documented (D)  Primary or critical habitat (list incidental habitat (list No usable habitat	or Su tat (lis t spec	specte it spec ies)	d (S) to		n (circle S S	one ba		n defini				nstruc	ions):							
I. Rating (use the conclustris function)	sions f	rom i a	above an	d the	matrix be	low to	arrive	at (circ	le) the	functio	nal p	oints a	nd ratir	ng (H =	high	, M = m	oderat	e, or L	= lov	v] fo
Highest Habitat Level		doc./	primary	Sus	/primary	ه	oc./se	condary	su	s./secor	ndan	dc	c./incid	ental	Lsu	s./incid	ental	Non	9	
Functional Points and Ra	tina	1 (H)		.90	н)	ا .	(M)		.7	(M)		.5	(L)		1/3			low		
		.g. observations, records, etc):																		
Primary or critical habi Secondary habitat (list Incidental habitat (list No usable habitat II. Rating (use the conclu- this function)	t speci	ies) es)	5. (0)	D D D od the		les le	arrive	at [circ	le] the	function	nal p	-//			-	n, Б .м=п			= lo	w] fo
Highest Habital Level		doc./	primary	ary sus/primary doc/secondary sus/secondary doc/incidental sus/incidental None																
Functional Points and Ra Sources for documented us					H)		(M)		6	(M)			<u> </u>		<u>Li</u>	a)		0 (1)	_	1
i. Evidence of overall will  Substantial (based on any observations of abundant wildlife sign presence of extremely interviews with local b	dlife u of the lant wil such a limitin	se in t e follow Idlife # as scat ng habi	the AA ( ving (che 's or hig t, tracks, itat featu	eck]): h spec nest : ires no	ies diver structures st availab	sity (d	uring a	any peri	od)	— fe	w or	Low ( no will no will adjac	based dlife ob dlife si ent upl	on any servat gn and fo	ions ( od so	e follow during p surces with kno	eak us	e perlo		
common occurrence of adequate adjacent up	red wildli and fo	ed wildlife groups or individuals or relatively few species during peak periods wildlife sign such as scat, tracks, nest structures, game trails, etc.																		
<ul><li>(L) rating. Structural diver- of their percent composition</li></ul>	sity is n of the	(working from top to bottom, circle appropriate AA attributes in matrix to arrive at exceptional (E), high (H), moderate (M), or love y is from #13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in term of the AA (see #10). Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = emporary/ephemeral; and A = absent (see instructions for further definitions of these terms).)																		
Structural diversity				High								erate					Lov	W		
(see #13) Class cover distribution		Eve	en	$\neg$	Uni	even		<del>                                     </del>	Ev	en		<u> </u>	Une	ven		<del> </del>	Eve	n	-	
(all vegetated classes)				4		_		<u> </u>			_	<u> </u>				<u></u>				
Duration of surface	P/	S/I	T/E	/ P	IP SI	T/	E   /	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	<b> </b>	

Structural diversity (see #13)				н	ligh						1	Mode	erate					Lov	٧	
Class cover distribution (all vegetated classes)		Ev	en			Unev	/en			Eve	n			Unev	ren			Eve	n	
Duration of surface water in > 10% of AA	P/ P	S/I	T/E	1	P/P	\$/I	T/E	1	P/P	S/I	T/E	A	P/P	S/I	T/E	^	P/P	S/I	T/E	,
Low disturbance at AA (see #12i)	E	E	E	ŀ	Œ	0	н	١	E	н	н	М	E	н	м	M	Ε	н	М	ľ
Moderate disturbance at AA (see #12i)	н	Н	н	1	Н	н	н	•	Н	н	М	М	н	М	М	L	Н	M	L	ı
High disturbance at AA	М	М	М	ı	M	M	L	ı	М	M	L	L	М	L	L	L	L	L	L	L

iii. Rating (use the conclusions from i and ii above and the matrix below to arrive at [circle] the functional points and rating (E = exceptional, H = high, M = moderate, or L = low) for this function)

Evidence of wildlife use (i)	Wildlife habitat features rating (ii)									
· ·	Exceptional	High	Moderate	Low						
Substantial	1(E)	.9 (H)	.8 (H)	.7 (M)						
Moderate	(H) (H)	.7 (M)	,5 (M)	.3 (L)						
Minimal	.6 (M)	.4 (M)	.2 (L)	.1 (L)						

Comments:

(see #12i)



14D. General Fish/Aquatic Habitat Rating: (Assess this function if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [i.e., fish use is predided by perched culvert or other barrier, etc.]. If the AA is not or was not historically used by fish due to lack of habitat, excessive gradient, etc., circle NA here and proceed to the next function. If fish use occurs in the AA but is not desired from a resource management perspective [such as fish use within an irrigation canal], then Habitat Quality [i below] should be marked as "Low", applied accordingly in ii below, and noted in the comments.)

Habitat Quality (circle appropriate AA attributes in matrix to arrive at exceptional (F), high (H), moderate (M), or low (L) quality rating

ii Habitat Guality (circle appropriate AA attributes in maura to	allits at	6XC6D00118	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	THE PARTY	CI ALC HYIL O	- C-	GOILL ION	14			
Duration of surface water in AA	Perm	ermanent / Perennial Seasonal / Intermittent						porary / Ephemeral			
Cover - % of waterbody in AA containing cover objects such as submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation, etc.	>25%	10–25%	<10%	>25%	10–25%	<10%	>25%	10–25%	<10%		
Shading - >75% of streambank or shoreline within AA contains riparian or wetland scrub-shrub or forested communities.	E	E	Н	Ŧ	н	М	М	М	М		
Shading - 50 to 75% of streambank or shoreline within AA contains rip, or wetland scrub-shrub or forested communities	Н	Н	М	М	М	М	М	L	L		
Shading - < 50% of streambank or shoreline within AA contains rip, or wetland scrub-shrub or forested communities	н	М	М	М	L	Г	L	L	L		

Modified Habitat Quality (Circle the appropriate response to the following question. If answer is Y, then reduce rating in i above by one level [E = H, H = M, M = L, L = L]). Is fish use of the AA precluded or significantly reduced by a culvert, dike, or other man-made structure or activity or is the waterbody included on the MDEQ list of waterbodies in need of TMDL development with listed "Probable Impaired Uses" including cold or warm water fishery or aquatic life support? Modified habitat quality rating = (circle) E

iii. Rating (use the conclusions from i and ii above and the matrix below to arrive at [circle] the functional points and rating [E = exceptional, H = high, M =

Types of fish known or	Iditosiony	Modified Habitat Quality (ii)										
suspected within AA	Exceptional	High	Moderate	Low								
Native game fish	1 (E)	.9 (H)	.7 (M)	.5 (M)								
Introduced game fish	.9 (H)	.8 (H)	.6 (M)	.4 (M)								
Non-game fish	.7 (M)	.6 (M)	.5 (M)	.3 (L)								
No fish	.5 (M)	.3 (L)	.2 (L)	.1 (L)								

#### Comments:

14E. Flood Attenuation: (applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA are not flooded from in-channel or overbank flow, circle NA here and proceed to next function.)

i. Rating (working from top to bottom, use the matrix below to arrive at (circle) the functional points and rating [H = high, M = moderate, or L = low] for this

Estimated wetland area in AA subject to periodic flooding		> 10 acres	w 190 - J. S		<10. >2 acre	s		<2 acres					
% of flooded wetland classified as forested, scrub/shrub, or both	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%				
AA contains no outlet or restricted outlet	1(H)	.9(H)	.6(M)	.8(H)	.7(H)	.5(M)	.4(M)	.3(L)	.2(L)				
AA contains unrestricted outlet	.9(H)	.8(H)	.5(M)	.7(H)	.6(M)	.4(M)	.3(L)	.2(L)	.1(L)				

ii. Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA (circle)? Y N Comments:

14F. Short and Long Term Surface Water Storage: (Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow. If no wetlands in the AA are subject to flooding or ponding, circle NA here and proceed with the evaluation.)

i. Rating (working from top to bottom, use the matrix below to arrive at (circle) the functional points and rating [H = high, M = moderate, or L = low] for this function. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral

[see instructions for further definitions of these terms].)									
Estimated maximum acre feet of water contained in wetlands	. :	>5 acre fee	t	<5	>1 acre fe	eet		≤1 acre fo	ot
within the AA that are subject to periodic flooding or ponding									
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	PÆ	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond > 5 out of 10 years	1(H)	9(H)	.8(H)	(BiHD)	.6(M)	.5(M)	.4(M)	.3(L)	,2(L)
Welfands in AA flood or pond < 5 out of 10 years	O(H)	8(H)	7/8/0	777.1	5(M)	4(M)	3(1)	2(1)	1/1.)

#### Comments:

14G. Sediment/Nutrient/Toxicant Retention and Removal: (Applies to wetlands with potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, circle NA here and proceed with the evaluation.)

i. Rating (working from top to bottom, use the matrix below to arrive at [circle] the functional points and rating [H = high, M = moderate, or L = low] for this function.

Sediment, nutrient, and toxicant input levels within AA	deliver l or co substan	eives or surre ow to moder empounds su ntially impaire rients or toxi	A receives or ver high level such that oth ajor sediment	"related to sediment, es or surrounding land levels of sediments, it other functions are mentation, sources of utrophication present.				
% cover of wetland vegetation in AA	>	70%		< 70%	> 7	0%		< 70%
Evidence of flooding or ponding in AA	Yes	No	Yes	No .	Yes	No	Yes	No
AA contains no or restricted outlet	1 (H)	.8 (H)	(7 (M))	.5 (M)	.5 (M)	.4 (M)	.3 (L)	.2 (L)
AA contains unrestricted outlet	9 (H)	7 (M)	6 (M)	.4 (M)	.4 (M)	.3 (L)	.2 (L)	.1 (L)

1 (L)

14H Sediment/Shoreline Stabilization: (applies only if AA occurs on or within the banks or a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action. If does not apply, circle NA here and proceed to next function)

i. Rating (working from top to bottom, use the matrix below to arrive at [circle] the functional points and rating [E = exceptional, H = high, M = moderate, or L

= low] for this function. % Cover of wetland streambank	Duration	of surface water adjacent to rooted ve	getation
or shoreline by species with deep, binding rootmasses	permanent / perennial	seasonal / intermittent	Temporary / ephemeral
> 65%	1 (H)	.9.(H)	.7 (M)
35.6A%	7 (M)	6 (M)	.5 (M)

3 (L)

< 35% Comments:

14I. Production Export	/Food Chain Support:
------------------------	----------------------

i. Rating (working from top to bottom, use the matrix below to arrive at [circle] the functional points and rating [H = high, M = moderate, or L = low] for this function. Factor A = acreage of vegetated component in the AA; Factor B = structural diversity rating from #13; Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to duration of surface water in the AA, where P/P = permanent/perennial; S/I = seasonal/intermittent; T/F./A= temporary/ephemeral or absent (see instructions for further definitions of these terms!)

2 (L)

1/E /A=	tempo	rarv/eon	emerai c	or absen	t isee in	STRUCTION	S TOT TUE	mer den	HUOTIS O	i mese i	emisj.,										
Α		Vegeta	ted com	ponent :	5 acres	3		Vegetat	ed comp	onent 1	<ul><li>5 acres</li></ul>			Vegeta	ted com	ponent ·	<1 acre				
В	Н	iah	Mode	erate	L	OW	Hi	ah .	Mode	erate	Lo	w	Hi	ah	Mode	erate		W			
C	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No			
P/P	(H)	.9H	.9H	.BH	.8H	.7M	.9H	.8H	.8H	.7M	.7M	.6M_	.7M	.6M	.6M	.4M	.4M	_3L_			
S/I	.9H	.8H	.8H	.7M	.7M	.6M	.8H	.7M	7M	.6M	.6M	.5M_	.6M	5M	.5M	.3L	3L	2L			
T/E/	.8H	.7M	.7M	.6M	.6M	.5M	.7M	.6M	.6M	.5M	.5M	.4M	.5M	.4M	.4M	.2L	.2L	.1L			
I A			9.0								1					1.0		( I			

#### Comments:

14J. Groundwater Discharge/Recharge: (Check the indicators in i & ii bek	ow that apply to the AA)
i. Discharge Indicators	ii. Recharge Indicators
	ermeable substrate present without underlying impeding layer
	etland contains inlet but no outlet
Wetland occurs at the toe of a natural slopeO	ther
Seeps are present at the wetland edge  AA permanently flooded during drought periods - a work or wetland contains an outlet, but no inlet	as invadamen decreased
★AA permanently flooded during drought periods - ARA SUPA ARE  ARE  ARE  ARE  ARE  ARE  ARE  ARE	A 0
Wetland contains an outlet, but no inlet	
Other	
<ol> <li>Rating: Use the information from i and ii above and the table below to a</li> </ol>	rrive at [circle] the functional points and rating [H = high, L = low] for this function
Criteria	Functional Points and Rating
AA is known Discharge/Recharge area or one or more indicators of D/R pro	esent (1 (H))
No Discharge/Recharge indicators present	.1 (L)
Available Discharge/Recharge information inadequate to rate AA D/R potential	ntial N/A (Unknown)

#### Comments:

14K. Uniqueness:

i. Rating (working from top to bottom, use the matrix below to arrive at [circle] the functional points and rating [H = high, M = moderate, or L = low] for this function

Replacement potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MNHP			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MNHP			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate		
Estimated relative abundance (#11)	rare	common	abundant	rare	common	abundant	rare	common	abundant
Low disturbance at AA (#12i)	1 (H)	.9 (H)	.8 (H)	.8 (H)	(6 (M)	.5 (M)	.5 (M)	.4 (M)	.3 (L)
Moderate disturbance at AA (#12i)	.9 (H)	.8 (H)	.7 (M)	.7 (M)	.5 (M)	.4.(M)	4 (M)	.3 (L)	.2 (L)
High disturbance at AA (#12i)	.8 (H)	.7 (M)	.6 (M)	.6 (M)	.4 (M)	.3 (L)	.3 (L)	.2 (L)	.1 (L)

#### Comments:

14L. Recreation/Education Potential: i. Is the AA a known rec./ed. site: (circle Y)N (If yes, rate as [circle] (ligh [1] and go to ii; if no go to iii)
ii. Check categories that apply to the AA: \_\_\_\_ Educational/scientific study; \_\_\_ Consumptive rec.; \_\_\_ Non-consumptive rec.; \_\_\_ Other

iii. Based on the location, diversity, size, and other site attributes, is there strong potential for rec./ed. use? Y N (If yes, go to ii, then proceed to iv; if no, then rate as [circle] Low [0.1])

iv. Rating (use the matrix below to arrive at [circle] the functional points and rating [H = high, M = moderate, or L = low] for this function.

Ownership	Disturbance at AA (#12i)					
	Jow.	moderate	high			
public ownership	(1 (Hz)	.5 (M)	,2 (L)			
private ownership	.7 (M)	.3 (L)	.1 (L)			

#### Comments:

LAND & WATER B-23

FUNCTION & VALUE SUMMARY & OVERALL RATING

Function & Value Variables	Rating	Actual Functional Points	Possible Function al Points	Functional Units; (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	Ł	.7	1	
B. MT Natural Heritage Program Species Habitat	L	. 1	1	
C. General Wildlife Habitat	++	.9	1	
D. General Fish/Aquatic Habitat	N/A		0	
E. Flood Attenuation	N/A		0	
F. Short and Long Term Surface Water Storage	H	. 8	1	
G. Sediment/Nutrient/Toxicant Removal	m	. 7	1	
H. Sediment/Shoreline Stabilization	N/A		0	
I. Production Export/Food Chain Support	H	1	1	
J. Groundwater Discharge/Recharge	H	1	1	
K. Uniqueness	W	. 6	1	
L. Recreation/Education Potential	++	1	1	
Totals:		6.8	9	128 (71%)

OVERALL ANALYSIS AREA (AA) RATING: (Circle appropriate category based on the criteria outlined below) I III IV
Category I Wetland: (Must satisfy one of the following criteria; if does not meet criteria, go to Category II)  Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; or  Score of 1 functional point for Uniqueness; or  Score of 1 functional point for Flood Attenuation and answer to Question 14E.II is "yes"; or  Total actual functional points > 80% (round to nearest whole #) of total possible functional points.
Category II Wetland: (Criteria for Category I not satisfied and meets any one of the following criteria; if not satisfied, go to Category IV)  Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; or Score of .9 or 1 functional point for General Wildlife Habitat; or  Score of .9 or 1 functional point for General Fish/Aquatic Habitat; or  "High" to "Exceptional" ratings for both General Wildlife Habitat and General Fish/Aquatic Habitat; or Score of .9 functional point for Uniqueness; or Total Actual Functional Points > 65% (round to nearest whole #) of total possible functional points.
Category III Wetland: (Criteria for Categories I, II or IV not satisfied)
Category IV Wetland: (Criteria for Categories I or II are not satisfied and all of the following criteria are met; if does not satisfy criteria go to Category III)  "Low" rating for Uniqueness; and "Low" rating for Production Export/Food Chain Support; and Total actual functional points < 30% (round to nearest whole #) of total possible functional points

# MDT WETLAND MONITORING - VEGETATION TRANSECT

LAND	& WATER	B-24
	maliba	

Site:	Creston	Date:	8/13/01	Examiner:	J. Asebrook	Transect #	1
-------	---------	-------	---------	-----------	-------------	------------	---

Approx. transect length: 465 feet Compass Direction from Start (Upland):

Vegetation type 1:	Wee	dy upland (see attached Ta	able 1)
Length of transect in the	his typ	be: 195 (3 sections)	feet
Elymus repens	4	Cirsium vulgare	+
yellow legume	4	Verbascum thapsus	+
Agrostis stolonifera	2	Phalaris arundinacea	+
Cirsium arvense	2	Stipa nelsonii	+
Medicago lupulina	1	Trifolium hybridum	+
Poa spp.	1		9001/02/02/03
Phleum pratense	1		
Poa pratensis	+		
Alopecurus pratensis	+		
Taraxacum officinale	+		
Rumex crispus	+		
Linum perenne	+		
Total Vegetative Cove	er:	75%	

Vegetation type 3:	type 3: Depressions: mixed TYPLAT/grasses				
Length of transect in this type:		63	feet		
Typha latifolia	3	Erigeron lonchophyllu	ıs? +		
Agrostis stolonifera	1	Medicago lupulina	+		
Eleocharis palustris	1	Taraxacum officinale	+		
Juncus tenuis	1	Epilobium ciliatum	+		
Juncus regelii	1	Potentilla anserina	+		
Juneus articulatus	1				
Alopecurus pratensis	1		95 (5)		
Melilotus officinale	1				
Cirsium arvense	+				
Equisetum arvense	+				
Phalaris arundinacea	+				
Plantago major	+				
Total Vegetative Cover	r: 409	%			

Vegetation type 2:	Typha latif	olia	
Length of transect in t	his type:	80	feet
Typha latifolia	5	Water's edge:	
Phalaris arundinacea	4	Ceratophyllum demersum	5
Eleocharis palustris	2	Sparganium emersum	1
Alopecurus pratensis	1	Eleocharis palustris	1
Agrostis stolonifera	1	Juneus articulatus	+
Rumex crispus	+	Beckmannia syzigachne	+
Epilobium ciliatum	+	Ranunculus scleratus	+
Plantago major	+		
Juneus articulatus	+		
Cirsium arvense	+		
Juncus tenuis	+		
Total Vegetative Cove	er: 80%		

Vegetation type 4:   Pha	alaris arundinacea	20
Length of transect in this	type: 100	feet
Phalaris arundinacea	5	
Eleocharis palustris	+	
Equisetum arvense	+	
Agrostis stolonifera	+	
Plantago major	+	9275
Lactuca serriola	+	
Myosotis laxa	+	
Carex bebbii	+	
Juncus tenuis	+	
Juneus articulatus	+	
Total Vegetative Cover:	98%	

MDT V	VETLAND MONITOR	ING – VEGETATION TRANSECT	LAND & WATER B-25
Site: Creston	Date: 8/13/2001	Examiner: J. Asebrook	Transect # 1
Approx. transect length: 465 feet	Compass Direc	tion from Start (Upland):	
Vegetation type 5: Potamageton pectina	tus	Vegetation type 6:	
Length of transect in this type: 25	feet	Length of transect in this type:	feet
Phalaris arundinacea	1		
Potamageton pectinatus	5		
Barbarea vulgaris	+		
Epilobium ciliatum	+		
Total Vegetative Cover: 75%		Total Vegetative Cover:	
Vegetation type 7:		Vegetation type 8:	
Length of transect in this type:	feet	Length of transect in this type:	feet
			200 E 200 E
	12 to 1		
Total Vegetative Cover:		Total Vegetative Cover:	

A-

# COMPREHENSIVE VEGETATION LIST



Species	Vegetation	Species	Vegetation
-	Community		Community
	Number(s)	1900	Number(s)
Elymus repens	1,3	Juneus articulatus	2,3,4
Yellow legume	1,3	Juncus regelii	3
Linum perenne	1	Ranunculus scleratus	2
Poa pratensis	1	Beckmannia syzigachne	2
Rumex crispus	1,2,3	Ceratophyllum demersum	2
Cirsium arvense	1,2,3,4,6	Carex bebbii	3,4
Taraxacum officinale	1,3,6	Erigeron lonchophyllus?	3
Phleum pratense	1	Gnaphalium palustre	3
Dactylis glomerata	1	Populus balsamifera	3
Polygonum convolvulus	1	Equisetum arvense	3,4
Alopecurus pratensis	1,2,3,6	Chenopodium rubrum	4,5
Silene latifolia	1	Veronica americana	4
Melilotus alba	1	Myosotis laxa	4
Melilotus officinale	1,3	Carex flava	3,6
Agrostis stolonifera	1,2,3,4,6	Ranunculus aquatilis	5
Poa spp.	1	Barbarea vulgaris	5
Medicago lupulina	1,3,6	Sparganium emersum	5
Trifolium hybridum	1,3	Potamageton pectinatus	5
Lactuca serriola	1,3,4,6	Potentilla anserina	3
Trifolium pratense	1,3,6	Carex arcta	3
Verbascum thapsus	1	Potamageton natans	5
Tragopogon dubius	1		40.0000.000
Bromus inermis	1		
Cynoglossum officinale	1		
Thlaspi arvense	1		
Cirsium vulgare	1,3		
Chenopodium album	1		
Plantago major	1,2,3,4,5		
Purple legume (Astragalus?)	1		
Phalaris arundinacea	1,2,3,4,5,6		
Epilobium ciliatum	1,2,3		900000
Typha latifolia	2,3		
Eleocharis palustris	2,3,4,5		
Juncus tenuis	2,3,4		
Plantago lanceolatum	1		

OMMENTS/PROBLEMS:			
	 		110-22-22-2
	 		-

# MDT WETLAND MONITORING - VEGETATION TRANSECT (back of form)



			LAND & WATER B-27
Cover Estim	ate	Indicator Class:	Source:
+=<1%	3 = 11-20%	+ = Obligate	P = Planted
1 = 1-5%	4 = 21-50%	- = Facultative/Wet	V = Volunteer
2 = 6-10%	100	0 = Facultative	, , , , , , , , , , , , , , , , , , , ,
2 - 0-1070	3-23070	o – I acuitati ve	
Percent of per	rimeter 100% % dev	eloping wetland vegetation – exclu	ding dam/berm structures.
this location v	with a standard metal fencepos	t. Extend the imaginary transect lin	transect should begin in the upland area. Permanently mark ne towards the center of the wetland, ending at the 3 food depth Mark this location with another metal fencepost.
			num, establish a transect at the windward and leeward sides of ventory, representative portions of the wetland site.
Notes:			
2000			
4			
7.5 10.04.0000			
<u> </u>			

# **Appendix C**

# REPRESENTATIVE PHOTOGRAPHS

MDT Wetland Mitigation Monitoring Creston Creston, Montana





Photo Point No. 1: View looking north; the Flathead County green bins are located in the distance.



Photo Point No. 2: View looking northeast; Highway 35 is visible in the background.



Photo Point No. 3: View looking east. The photo is taken near the north perimeter of the impoundment.



Photo Point No. 4: View looking south across the mitigation site.



Photo Point No. 6: View looking west; the shallow pond is present in the background.



Photo Point No. 5: View looking south and taken from the center of the mitigation site.





Photo Point No. 6: View looking west; the large pond is present in the background.



Photo Point No. 6: View looking north.



East end of vegetation transect looking east.



East end of vegetation transect looking west along the transect



West end of vegetation transect looking east along the transect line.



West end of vegetation transect looking west. Transect stake is present in the foreground.

# **Appendix D**

BIRD SURVEY PROTOCOL
MACROINVERTEBRATE SAMPLING PROTOCOL
GPS PROTOCOL

MDT Wetland Mitigation Monitoring Creston Creston, Montana



## **BIRD SURVEY PROTOCOL**

The following is an outline of the MDT Wetland Mitigation Site Monitoring Bird Survey Protocol. Though each site is vastly different, the bird survey data collection methods must be standardized to a certain degree to increase repeatability. An Area Search within a restricted time frame will be used to collect the following data: a bird species list, density, behavior, and habitat-type use. There will be some decisions that team members must make to fit the protocol to their particular site. Each of the following sections and the desired result describes the protocol established to reflect bird species use over time.

## Species Use within the Mitigation Wetland: Survey Method

Result: To conduct a bird survey of the wetland mitigation site within a restricted period of time and the budget allotment.

### Sites that can be circumambulated or walked throughout.

These types of sites will include ponds, enhanced historic river channels, wet meadows, and any area that can be surveyed from the entirety of its perimeter or walked throughout. If the wetland is not uncomfortably inundated, conduct several "meandering" transects through the site in an orderly fashion (record the number and approximate location/direction of the transects in the field notebook; they do not have to be formalized or staked). If a very small portion of the site cannot be crossed due to inundation, this method will also apply. Though the sizes of the site vary, each site will require surveying to the fullest extent possible within a set time limit. The optimum times to conduct the survey are in the morning hours. Conduct the survey from sunrise to no later than 11:00 AM. (Note: some sites may have to be surveyed in the late afternoon or evening due to time constraints or weather; if this is the case, record the time of day and include this information in your report discussion.) If the survey is completed before 11:00 AM and no additions are being made to the list, then the task is complete. The overall limiting factor regarding the number of hours that are spent conducting this survey is the number of budgeted hours; this determination must be made by site by each individual.

In many cases, binoculars will be the only instrument that is needed to identify and count the birds using the wetland. If the wetland includes deep water habitat that can not be assessed with binoculars, then a scope and tripod are necessary. If this is the case, establish as many lookout posts as necessary from key vantage points to collect the data. Depending on the size of the open water, more time may be spent viewing the mitigation area from these vantage points than is spent walking the peripheries of more shallow-water wetlands.

# Sites that cannot be circumambulated.

These types of sites will include large-bodied waters, such as reservoirs, particularly those with deep water habitat (>6 ft) close to the shore and no wetland development in that area of the shoreline. If one area of the reservoir was graded in such a way to create or enhance the development of a wetland, then that will be the area in which the ambulatory bird survey is conducted. The team member must then determine the length of the shoreline that will be surveyed during each visit.



As stated above in the ambulatory site section, these large sites most likely will have to be surveyed from established vantage points.

## Species Use within the Mitigation Wetland: Data Recording

Result: A complete list of bird species using the site, an estimate of bird densities and associated behaviors, and identification of habitat use.

# 1. Bird Species List

Record the bird species on the Bird Survey - Field Data Sheet using the appropriate 4-letter code of the common name. The coding uses the first two letters of the first two words of the birds' common name or if one name, the first four (4) letters. For example, mourning dove is coded MODO and mallard is MALL. If an unknown individual is observed, use the following protocol and define your abbreviation at the bottom of the field data sheet: unknown shorebird: UNSB; unknown brown bird (UNBR); unknown warbler (UNWA); unknown waterfowl (UNWF). For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parentheses; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded: UNBB / FO (25). You may also note on the data sheet if that particular individual is using a constructed nest box.

# 2. Bird Density

In the office, sum the Bird Survey – Field Data Sheet data by species and by behavior. Record this data in the Bird Summary Table.

### 3. Bird Behavior

Bird behavior must be identified by what is known. When a species is simply observed, the behavior that it is immediately exhibiting is what is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair individual (BP); foraging (F); flyover (FO); loafing (L; e.g. sleeping, roosting, floating with head tucked under wing are loafing behaviors); and, nesting (N). If more behaviors are observed that do have a specific descriptive word, use them and we will add it to the protocol; descriptive words or phrases such as "migrating" or "living on site" are unknown behaviors.

## 4. Bird Species Habitat Use

We are interested in what bird species are using which particular habitat within the mitigation wetlands. This data is easily collected by simply recording what habitat the species was initially observed. Use the following broad category habitat classifications: aquatic bed (AB - rooted floating, floating-leaved, or submergent vegetation); forested (FO); marsh (MA - cattail, bulrush, emergent vegetation, etc. with surface water); open water (OW - primarily unvegetated); scrubshrub (SS); and upland buffer (UP); wet meadow (WM - sedges, rushes, grasses with little to no surface water). If other categories are observed onsite that are not suggested here, we will make a new category next year.



D-2

# AQUATIC INVERTEBRATE SAMPLING PROTOCOL

## **Equipment List**

- D-frame sampling net with 1 mm mesh. Wildco is a good source of these.
- Spare net.
- 1-liter plastic sample jars, wide-mouth. VWR has these: catalog #36319-707.
- 95% ethanol: Northwest Scientific in Billings carries this.

All these other things are generally available at hardware or sporting goods stores. Make the labels on an ink jet printer preferably.

- hip waders.
- pre-printed sample labels (printed on Rite-in-the-Rain or other coated paper, two labels per sample).
- pencil.
- plastic pail (3 or 5 gallon).
- large tea strainer or framed screen.
- towel.
- tape for affixing label to jar.
- cooler with ice for sample storage.

#### Site Selection

Select the sampling site with these considerations in mind:

- Select a site accessible with hip waders. If substrates are too soft, lay a wide board down to walk on.
- Determine a location that is representative of the overall condition of the wetland.

# Sampling

Wetland invertebrates inhabit the substrate, the water column, the stems and leaves of aquatic vegetation, and the water surface. Your goal is to sweep the collecting net through each of these habitat types, and then to combine the resulting samples into the 1-liter sample jar.

Dip out about a gallon of water into the pail. Pour about a cup of ethanol into the sample jar. Fill out the top half of the sample labels, using pencil, since ink will dissolve in the ethanol.

Ideally, you can sample a swath of water column from near-shore outward to a depth of approximately 3 feet with a long sweep of the net, keeping the net at about half the depth of the water throughout the sweep. Sweep the water surface as well. Pull the net through a vegetated area, beneath the water surface, for at least a meter of distance.

Sample the substrate by pulling the net along the bottom, bumping it against the substrate several times as you pull.



This step is optional, but it gives you a chance to <u>see</u> that you've collected some invertebrates. Rinse the net out into the bucket, and look for insects, crustaceans, etc. If necessary, repeat the sampling process in a nearby location, and add the net contents to the bucket. Remember to sample all four environments.

Sieve the contents of the bucket through the straining device and pour or carefully scrape the contents of the strainer into the sample jar.

If you skip the bucket-and-sieve steps, simply lift handfuls of material out of the sampling net into the jars. In either case, please include some muck or mud and some vegetation in the jar. Often, you will have collected a large amount of vegetable material. If this is the case, lift out handfuls of material from the sieve into the jar, until the jar is about half full. Please limit material you include in the sample, so that there is only a single jar for each sample.

Top off the sample jar with enough ethanol to cover all the material in the jar. Leave as little headroom as possible.

It is not necessary to sample habitats in any specified order. Keep in mind that disturbing the habitats prior to sampling will chase off the animals you are trying to capture.

Complete the sample labels. Place one label inside the sample jar and tape the other label securely to the outside of the jar. Dry the jar before attaching the outer label if necessary. In some situations, it may be necessary to collect more than one sample at a site. If you take multiple samples from the same site, clearly indicate this by using individual sample numbers, along with the total number of samples collected at the site (e.g. Sample #3 of 5 total samples).

Photograph the sampled site.

# Sample Handling/Shipping

- In the field, keep collected samples cool by storing them in a cooler. Only a small amount of ice is necessary.
- Inventory all samples, preparing a list of all sites and enumerating all samples, before shipping or delivering to the laboratory.
- Deliver samples to Rhithron.



# **GPS Mapping and Aerial Photo Referencing Procedure**

The wetland boundaries, photograph location points and sampling locations were field located with mapping grade Trimble Geo III GPS units. The data was collected with a minimum of three positions per feature using Course/Acquisition code. The collected data was then transferred to a PC and differentially corrected to the nearest operating Community Base Station. The corrected data was then exported to ACAD drawings in Montana State Plain Coordinates NAD 83 international feet.

The GPS positions collected and processed had a 68% accuracy of 7 feet except in isolated areas of Tasks .008 and .011, where it went to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

Aerial reference points were used to position the aerial photographs. This positioning did not remove the distortion inherent in all photos; this imagery is to be used as a visual aide only. The located wetland boundaries were given a final review by the wetland biologist and adjustments were made if necessary.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

