
MONTANA DEPARTMENT OF TRANSPORTATION WETLAND MITIGATION MONITORING REPORT: YEAR 2005

*Camp Creek
Sula, Montana*



Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Ave
Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING
~ A DIVISION OF **PBS&J**
P.O. Box 239
Helena, MT 59624

December 2005

Project No: B43054.00 - 0106



MONTANA DEPARTMENT OF TRANSPORTATION

WETLAND MITIGATION MONITORING REPORT:

YEAR 2005

*Camp Creek
Sula, Montana*

Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Ave
Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING
A DIVISION OF PBS&J
P.O. Box 239
Helena, MT 59624

December 2005

Project No: B43054.00 - 0106



TABLE OF CONTENTS

1.0 INTRODUCTION.....	1
2.0 METHODS	3
2.1 Monitoring Dates and Activities	3
2.2 Hydrology	3
2.3 Vegetation	3
2.4 Soils.....	4
2.5 Wetland Delineation	4
2.6 Mammals, Reptiles, and Amphibians	4
2.7 Birds	5
2.8 Macroinvertebrates	5
2.9 Functional Assessment.....	5
2.10 Photographs.....	5
2.11 GPS Data.....	5
2.12 Maintenance Needs	5
3.0 RESULTS	6
3.1 Hydrology	6
3.2 Vegetation.....	7
3.3 Soils.....	12
3.4 Wetland Delineation	12
3.5 Wildlife	14
3.6 Macroinvertebrates	16
3.7 Functional Assessment.....	17
3.8 Photographs.....	18
3.9 Revegetation	18
3.10 Maintenance Needs/Recommendations	20
3.11 Current Credit Summary	20
4.0 REFERENCES.....	21

TABLES

Table 1	<i>2002 - 2005 Camp Creek vegetation species list.</i>
Table 2	<i>Transect 1 data summary.</i>
Table 3	<i>Wetland conditions within Camp Creek Wetland Mitigation Site.</i>
Table 4	<i>Fish and wildlife species observed at the Camp Creek Mitigation Site during 2002 – 2005 monitoring.</i>
Table 5	<i>Summary of 2001 (baseline) and 2004 wetland function/value and functional points at Camp Creek.</i>
Table 6	<i>Potential functional unit-based credit for Camp Creek Mitigation Project.</i>

FIGURES

Figure 1	<i>Project Site Location Map</i>
Figure 2	<i>Monitoring Activity Locations 2005</i>
Figure 3	<i>Mapped Site Features 2005</i>
Figure 4	<i>Pre-Developed Wetland 2001</i>
Figure 5	<i>Channel Cross Sections</i>
Figure 6	<i>Flood Channel Plan and Typical</i>

CHARTS

Chart 1	<i>Mean monthly flows for May – August from 2002-2005 as compared to the long-term mean monthly flows (1937-2004) on the Bitterroot River near Darby, MT.</i>
Chart 2	<i>Transect maps showing vegetation type from the start of transect (0 feet) to the end of transect (471 feet) for each year monitored.</i>
Chart 3	<i>Length of vegetation communities along Transect 1 for each year monitored.</i>
Chart 4	<i>Westslope cutthroat trout survey for Camp Creek (MFWP 2004).</i>
Chart 5	<i>Bioassessment scores for Camp Creek.</i>

APPENDICES

- Appendix A *Figures 2, 3, 4, and 6*
- Appendix B *2005 Wetland Mitigation Site Monitoring Form*
2005 Bird Survey Form
2005 Wetland Delineation Forms
2005 Functional Assessment Forms
- Appendix C *Representative Photographs*
- Appendix D *Original Site Plan*
- Appendix E *Bird Survey Protocol*
GPS Protocol
- Appendix F *2005 Macroinvertebrate Sampling Protocol and Data*
- Appendix G *Figure 5 Channel Cross Sections*
Planting Specifications

1.0 INTRODUCTION

This report documents the fourth year (2005) of monitoring at the Camp Creek mitigation site. The Camp Creek project was developed to mitigate wetland impacts associated with the Montana Department of Transportation (MDT) proposed Sula-North and South project, and to possibly function as a mitigation reserve to be applied against future MDT projects in the Bitterroot Valley. Camp Creek is located in Ravalli County, MDT Watershed # 3, in the Lower Clark Fork region. The mitigation site is located approximately three miles south of Sula, Montana (**Figure 1**). Elevations of the site range from 4,600 ft at the north boundary to 4,730 ft at the south boundary.

The approximate site boundary is illustrated on **Figure 2 (Appendix A)**, and the original site plans are included in **Appendix D**. The project is located within the Sula Basin and along the historic Camp Creek floodplain. Camp Creek flows across the valley bottom, until eventually draining into East Fork of the Bitterroot River. Seasonal flooding and perennial creek flow provide the primary hydrology source within the new channel/floodplain margins. Local groundwater systems serve as a secondary hydrology source, flowing through the deep alluvial substrate contained within the Sula Basin. Two smaller creeks drain into Camp Creek within the project limits: Andrews and Praine creeks.

Construction at the Camp Creek mitigation site was completed during the spring of 2002. The overall goals of this project were restoration of Camp Creek channel bottom, associated wetland functional restoration/enhancement and creation, and enhancement of heavily grazed and cleared riparian vegetation. MDT is currently developing a credit allocation scheme for this site in cooperation with the Corps of Engineers (see Section 3.11). Construction diagrams are presented in **Appendix D**. Project details for each of the three main goals are included in the following list:

Functional Restoration

- Return Camp Creek to its historic channel and establish new channel.
- Restore hydrology and vegetation, recreating high value wetland habitat along Camp Creek riparian corridor.
- Fill existing ditches.

Enhancement



- Riparian shrub and tree plantings throughout the created floodplain margins.
- Drier upland species planting in areas of created upland slopes.

Creation

- Creation of emergent/scrub shrub wetlands along the floodplain margins of the new channel.

The site was designed to mitigate for specific wetland functions impacted by MDT roadway projects, including: storm water retention, roadway runoff filtration, sediment and nutrient retention, water quality, groundwater recharge, and wildlife habitat. The Camp Creek site is typically monitored once per year to document wetland and other biological attributes. The monitoring area is illustrated in **Figure 2 (Appendix A)**.

FIGURE 1. PROJECT LOCATION
Camp Creek
Mitigation Site



PROJECT #: 130091.038
DATE: DEC 2002
LOCATION:
PROJECT MANAGER: B. DUTTON
DRAWN BY: B. NOECKER

LAND & WATER CONSULTING, INC.

1120 CEDAR PO BOX 8254 MISSOULA, MT 59807

2.0 METHODS

2.1 Monitoring Dates and Activities

The site was visited on August 1st (mid-season). Monitoring activities were conducted on the MDT-owned portion of the site, as well as within the fenced portion of the adjacent Grasser property. The mid-season visit was conducted to document vegetation, soil, and hydrologic conditions used to map jurisdictional wetlands. All information contained on the Wetland Mitigation Site Monitoring Form (**Appendix B**) was collected at this time. Activities and information conducted/collected included: wetland delineation; wetland/open water aquatic habitat boundary mapping; vegetation community mapping; vegetation transect; soils data; hydrology data; bird and general wildlife use; photograph points; macroinvertebrate sampling; GPS data points; functional assessment; (non-engineering) examination of topographic features; and stream cross section data at two established transects.

2.2 Hydrology

Wetland hydrology indicators were recorded during the mid-season visit using procedures outlined in the COE 1987 Wetland Delineation Manual (Environmental Laboratory 1987). Hydrology data were recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). Additional hydrologic data were recorded on the mitigation site monitoring form (**Appendix B**). No groundwater monitoring wells were installed at the site.

Two cross section locations were established and surveyed across Camp Creek on the MDT-owned parcel: one upstream and one downstream of the Praine Creek confluence with Camp Creek. These are designated “XS 3-A” and “XS 4A” on **Figure 2** in **Appendix A**. The cross sections will be used to monitor potential lateral and vertical channel migration over time.

2.3 Vegetation

General dominant species-based vegetation community types (e.g., *Carex/Phalaris*) were delineated on an aerial photograph during the mid-season visit. Standardized community mapping was not employed as many of these systems are geared towards climax vegetation and do not reflect yearly changes. Estimated percent cover of the dominant species in each community type was listed on the site monitoring form (**Appendix B**).

A 10-foot wide belt transect was sampled during the mid-season monitoring event to represent the range of current vegetation conditions. Percent cover was estimated for each vegetative species within each successive vegetative community encountered within the “belt” using the following values: T (few plants); P (1-5%), 1 (5-15%); 2 (15-25%); 3 (25-35%); 4 (35-45%); 5 (45-55%) and so on to 9 (85-95%). The transect location is illustrated on **Figure 2** in **Appendix A**. The transect will be used to evaluate changes over time, especially the establishment and increase of hydrophytic vegetation. The transect location was marked on the air photo and all data were recorded on the mitigation site monitoring form. Transect endpoint locations were recorded with the GPS unit in 2002. A photo was taken from both ends of the transect looking along the transect path.

A comprehensive plant species list for the site was compiled and will be updated as new species are encountered. Ultimately, observations from past years will be compared with new data to document vegetation changes over time. Revegetation enhancements were implemented in the spring of 2002. Survival rates for planted species were recorded during the mid-season monitoring visit.

2.4 Soils

Soils were evaluated during the mid-season site visit using the hydric soils determination procedures outlined in the COE 1987 Wetland Delineation Manual. Soil data were recorded for each wetland determination point on the COE Routine Wetland Delineation Data Forms (**Appendix B**). The most current terminology used by NRCS was used to describe hydric soils (USDA 1998).

2.5 Wetland Delineation

Wetland delineation was conducted during the mid-season visit according to the 1987 COE Wetland Delineation Manual. Wetland and upland areas within the monitoring area were investigated for the presence of wetland hydrology, hydrophytic vegetation and hydric soils. The information was recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). The wetland/upland boundary was originally delineated on the air photo and recorded with a resource grade GPS unit using the procedures outlined in **Appendix E**. Modifications to these boundaries in 2005 were accomplished by hand-mapping onto the 2002 aerial photograph. The wetland/upland boundary in combination with the wetland/open water boundary was used to calculate the final wetland acreage.

The Corps of Engineers concurred with a revised baseline delineation of 43.36 acres of wetland /open water channel signature on the MDT parcel and 5.37 acres of wetland / open water channel within the monitoring limits on the Grasser parcel for a total of 48.73 acres (Steinle pers. comm.). Pre-project wetlands are shown on **Figure 4, Appendix A**.

2.6 Mammals, Reptiles, and Amphibians

Mammal, reptile, and amphibian species observations and other positive indicators of use, such as vocalizations, were recorded on the wetland monitoring form during the mid-season visit. Indirect use indicators, including tracks; scat; burrows; eggshells; skins; bones; etc., were also recorded. These observations were recorded as the observer traversed the site while conducting other required activities. Direct sampling methods, such as snap traps, live traps, and pitfall traps, were not implemented. A comprehensive species list for the entire site was compiled. Observations from past years will ultimately be compared with new data.

2.7 Birds

Bird observations were also recorded during the mid-season visit. No formal census plots, spot mapping, point counts, or strip transects were conducted. Observations were recorded incidental to other monitoring activities and were categorized by species, activity code, and general habitat association.

2.8 Macroinvertebrates

Macroinvertebrate samples were collected during the mid-season site visit at one location along Camp Creek (**Figure 2**). Macroinvertebrate sampling procedures are provided in **Appendix F**. Samples were preserved as outlined in the sampling procedure and sent to Rhithron Associates for analysis.

2.9 Functional Assessment

A functional assessment form was completed using the 1999 MDT Montana Wetland Assessment Method (**Appendix B**). Field data necessary for this assessment were collected during the mid-season visit. Turnstone Biological completed a baseline functional assessment in 2001.

2.10 Photographs

Photographs were taken illustrating current land uses surrounding the site, the upland buffer, the monitored area and the vegetation transects. Each photograph point location was recorded with a resource grade GPS in 2002. The location of photo points is shown on **Figure 2, Appendix A**. All photographs were taken using a digital camera.

2.11 GPS Data

During the 2002 monitoring season, point data were collected with a resource grade GPS unit at the vegetation transect beginning and ending locations and at all photograph locations. Wetland boundaries were also recorded with a resource grade GPS unit in 2002, but were modified via hand mapping onto aerial photographs in 2005. The method used to collect these points is described in the GPS protocol in **Appendix E**.

2.12 Maintenance Needs

Observations were made of existing structures and of erosion/sediment problems to identify maintenance needs. This did not constitute an engineering-level structural inspection, but rather a cursory examination. Current or future potential problems were documented on the monitoring form.

3.0 RESULTS

3.1 Hydrology

The main source of hydrology for this site is Camp Creek, a perennial stream draining out of the south end of the Bitterroot Range. Seasonal flooding of Camp Creek occurs during spring runoff. Secondary sources of hydrology include runoff from ephemeral drainages east of the site and the persistent movement of groundwater through coarse alluvium materials located throughout the valley bottom. The mitigation site is located within the historic Camp Creek floodplain. The site consists of a constructed main channel, streambanks and floodplain terraces. Depressional wetlands are present, supported by seasonal overland flooding of Camp Creek and groundwater flows. Where it enters Grasser's parcel south of the MDT-owned parcel, the creek once was diverted into a channel running along the edge of Hwy 93. Several ditches designed to drain the wetland meadow complex were filled and closed during construction activities. Removal of drain ditches allows for groundwater systems to recharge and provide possible higher storage functions. Average high water levels were recorded at 222 cfs (Turnstone Biological, 2001). Lower water flows are on average 10 cfs.

Precipitation was close to "normal" in the general project area from January through June 2005, based on data from the Sula 3 ENE weather station (2005 data from July on were not yet available). Precipitation during this period totaled over 8.4 inches, which is 95% of the 8.85-inch mean for the January-July period between 1955 and 2005. However, stream flow was sub-normal in 2005. Based on USGS data from the stream gauge on the Bitterroot River near Darby, flows in May, June, and July of 2005 were only 65%, 50%, and 61% of the long-term mean monthly flows for these months. Flow increased in August to 112% of the mean, subsequent to the majority of the growing season. **Chart 1** provides a comparison of the long-term mean May, June, July, and August flows to mean flows for those months in 2002-2005 at the Bitterroot River near Darby..


Rock channel bottom occurred across approximately 2.15 acres or 5% of the current 46-acre mitigation site (**Figure 3**). Depths of the creek varied, ranging from 0.5 ft in the straight segments to 2 - 3 ft deep around the bends and meanders.

Cross section results are presented in **Figure 5 (Appendix G)**. These cross sections represent, in essence, post-project "baseline" (2002), as well as 2003, 2004 and 2005 channel conditions. Cross section results measured during the 2005 monitoring show that some adjustments have taken place.

Cross Section 3-A is located below the Praine Creek confluence. During 2005 runoff, this cross-section changed shape somewhat. The left bank and that side of the channel remained in the same location as 2004. Changes were observed along the channel bottom, and the gradual floodplain transition that developed along the right bank during 2004 was cut down to nearly the original channel depths observed during 2002.

Cross Section 4-A is located above the Praine Creek confluence. This cross section also adjusted slightly during 2005 runoff. The left and right banks remained in the similar location as in 2004, while the channel bottom slightly decreased in depth from the 2004 locations. Cross section monitoring will continue to ascertain stability and facilitate development of corrective measures, if necessary.

Chart 1: Mean monthly flows for May – August from 2002-2005 as compared to the long-term mean monthly flows (1937-2004) on the Bitterroot River near Darby, MT.



3.2 Vegetation

Eighty-seven plant species were identified at the site and are listed in **Table 1**. The majority of these species are herbaceous, found in wetland meadow complexes with minor tree or shrub coverage. Several remnant shrub patches exist along dry oxbows of historic Camp Creek. With the reintroduction of hydrology into the old channels, these shrub patches are now receiving water again and should flourish over time. Several mature black cottonwood (*Populus trichocarpa*) stands are also located amongst shrub patches. Large areas of wet meadows exist within the areas of lower topography. These wet meadows are seasonally inundated and groundwater-fed.

Three wetland types and three upland community types were identified and mapped at the mitigation site (**Figure 3, Appendix A**). The three wetland community types include Type 2: *Carex/Phalaris*, Type 3: *Agrostis/Deschampsia* and Type 6: *Populus/Salix*. The three upland community types include Type 1: *Agropyron/Trifolium*, Type 5: *Agropyron/Centaurea* and Type 7: *Phalaris /Centaurea*. Plant species observed within each of these communities are listed on the attached data form (**Appendix B**).

Wetland types 2 & 6 were present before construction of the main channel. Pre-construction wetland delineation mapped the majority of the site as emergent wetlands. Type 2 is a remnant wetland with heavy past alterations due to livestock grazing and historic clearing of riparian vegetation. Type 2 is the wettest community and occurs as emergent wetlands in saturated to shallow water conditions. Type 6 consists of several shrubs such as willow (*Salix*), alder (*Alnus*) and birch (*Betula*), found along the old dry oxbows and depressions. Higher on the banks, just above the streambed, mature cottonwoods are present along the old terraces.

The remaining wetland type was created during the channel reconstruction, and includes the geotextile fabric wrapped streambanks and floodplain areas. Community Type 4: *Salix/Agropyron*, mapped during the 2002 monitoring, was included within the Type 3: *Agrostis/Deschampsia* community during 2003-2005 monitoring. Community type classification for Type 4 was based on the dominant grass species and willow sprigging used during construction efforts. During the 2003 monitoring the Type 4 grasses had changed from wheatgrass (*Agropyron*) to reedtop (*Agrostis alba*) and tufted hairgrass (*Deschampsia cespitosa*). During the 2004-2005 monitoring these wetter type species have continued to increase in cover and now dominate the floodplain areas.

Revegetation efforts were implemented along the streambanks and floodplain margins during 2002 construction. These included planting of 10-cubic gallon shrubs, trees and sprigging of willows. Species planted for riparian enhancement included cottonwood, willows, dogwood (*Cornus stolonifera*) and aspen (*Populus tremuloides*). Survival data are presented in **Appendix B** and describe specific details on each species.

Adjacent upland vegetation communities are mainly dominated by rangeland and/or aggressive invasive species. Type 1 consists of several spoil piles created for upland vegetation enhancement. These areas were planted with a mix of 5-cubic gallon plantings and weed matting. Upland plantings included Douglas fir (*Pseudotsuga menziesii*), lodgepole pine (*Pinus contorta*) ponderosa pine (*Pinus ponderosa*), serviceberry (*Amelanchier alnifolia*), shrubby potentilla (*Potentilla fruticosa*), snowberry (*Symphoricarpos albus*) and woods rose (*Rosa woodsii*). Dominant species included pasture grasses and mostly weedy disturbance species such as quackgrass (*Agropyron repens*), pennycress (*Thlaspi arvensis*), dandelion (*Taraxacum officinale*), and tumble mustard (*Sisymbrium altissimum*). During monitoring, plantings did not contribute enough coverage to be considered significant in determining them as dominant in the community type.

Type 5 consists of upland areas historically grazed, dominated with pasture grasses such as quackgrass, meadow foxtail (*Alopecurus pratensis*) and smooth brome (*Bromus inermis*). Type 5 also has a high distribution of spotted knapweed (*Centaurea maculosa*), located in the

transition zone between wetland bottoms and open forest slopes. Several noxious weeds were observed throughout the Camp Creek Mitigation Site: spotted knapweed, Canada thistle (*Cirsium arvense*), Oxeye daisy (*Chrysanthemum leucanthemum*), and hound's-tongue (*Cynoglossum officinale*). Other weedy or non-native species included bull thistle (*Cirsium vulgare*), common dandelion, lambsquarters (*Chenopodium album*), clasping pepper-grass (*Lepidium perfoliatum*), pennycress, tumbleweed and quackgrass.

Vegetation transect results are detailed in the attached data forms (**Appendix B**) and are summarized below in **Table 2** and **Charts 2** and **3**. The previous years transect data is included to compare changes between monitoring years.

Table 1: 2002 - 2005 Camp Creek vegetation species list.

Scientific Name ¹	Common Name	Region 9 (Northwest) Wetland Indicator
<i>Achillea millefolium</i>	Common Yarrow	FACU
<i>Agropyron repens</i>	Quackgrass	FACU
<i>Agrostis alba</i>	Redtop	FAC+
<i>Alnus incana</i>	Thin leaved alder	FACW
<i>Alopecurus pratensis</i>	Meadow foxtail	FACW
<i>Amelanchier alnifolia</i>	Service-berry	FACU
<i>Aster integrifolius</i>	thickstem aster	--
<i>Betula occidentalis</i>	Water birch	FACW
<i>Bromus inermis</i>	Smooth brome	--
<i>Bromus tectorum</i>	Cheatgrass	--
<i>Calamagrostis canadensis</i>	Bluejoint reedgrass	FACW+
<i>Carex aquatilis</i>	Water sedge	OBL
<i>Carex bebbii</i>	Bebb's sedge	OBL
<i>Carex nebrascensis</i>	Nebraska sedge	OBL
<i>Carex crawfordii</i>	Crawford's sedge	FAC
<i>Carex lanuginosa</i>	Wooly sedge	OBL
<i>Carex praegracilis</i>	Clustered field sedge	FACW
<i>Carex utriculata</i>	Beaked sedge	OBL
<i>Centaurea maculosa</i>	Spotted Knapweed	--
<i>Cercocarpus ledifolius</i>	Mountain-mahogany	--
<i>Chenopodium album</i>	White Goosefoot	FAC
<i>Chrysanthemum leucanthemum</i>	Oxeye daisy	--
<i>Cirsium arvense</i>	Canada Thistle	FACU+
<i>Cirsium vulgare</i>	Bull thistle	FACU
<i>Cornus stolonifera</i>	Red-osier dogwood	FACW
<i>Crataegus douglasii</i>	Douglas Hawthorn	FAC
<i>Crepis tectorum</i>	Annual hawksbeard	--
<i>Cynoglossum officinale</i>	Hound's tongue	FACU
<i>Danthonia spp.</i>	Oatgrass	--
<i>Deschampsia cespitosa</i>	Tufted hairgrass	FACW
<i>Epilobium ciliatum</i>	Hairy willow-herb	FACW+
<i>Epilobium paniculatum</i>	Willow-herb	--
<i>Equisetum arvense</i>	Field horsetail	FAC
<i>Equisetum laevigatum</i>	Smooth scouring-rush	FACW
<i>Festuca pratensis</i>	Meadow fescue	FACU+

Table 1 (continued): 2002 - 2005 Camp Creek vegetation species list.

Scientific Name ¹	Common Name	Region 9 (Northwest) Wetland Indicator
<i>Geum macrophyllum</i>	Big leafed avens	OBL
<i>Glyceria elata</i>	Tall mannagrass	FACW+
<i>Glyceria grandis</i>	American mannagrass	OBL
<i>Gnaphalium palustre</i>	Cudweed	FAC+
<i>Juncus balticus</i>	Baltic rush	FACW
<i>Juncus bufonius</i>	Toad rush	FACW
<i>Juncus confusus</i>	Colorado rush	FAC
<i>Juncus ensifolius</i>	Three-stamen Rush	FACW
<i>Lactuca serriola</i>	Prickly lettuce	FAC-
<i>Lepidium perfoliatum</i>	Clasping pepper-grass	FACU+
<i>Linaria vulgaris</i>	Butter and eggs	--
<i>Lonicera involucrate</i>	Honeysuckle	FAC+
<i>Lupinus wyethii</i>	Wyeth's lupine	NI
<i>Lychnis alba</i>	White campion	--
<i>Matricaria matricarioides</i>	Pineapple-weed	FACU
<i>Melilotus officinalis</i>	Yellow Sweet clover	FACU
<i>Mentha arvensis</i>	Field mint	FAC
Mimulus guttatus	monkey-flower	OBL
<i>Phalaris arundinacea</i>	Canary Reed Grass	FACW
<i>Phleum pratense</i>	Timothy	FACU
<i>Pinus ponderosa</i>	Ponderosa pine	--
<i>Plantago major</i>	Plantain	FACU+
<i>Poa pratensis</i>	Kentucky Bluegrass	FACU+
<i>Polygonum amphibium</i>	Water smartweed	OBL
<i>Populus tremuloides</i>	Quaking aspen	FAC+
<i>Populus trichocarpa</i>	Cottonwood	FAC
<i>Potentilla fruticosa</i>	Shrubby cinquefoil	FAC-
<i>Potentilla gracilis</i>	Northwest cinquefoil	FAC
<i>Pseudotsuga menziesii</i>	Douglas fir	FACU
<i>Ranunculus aquatilis</i> var. <i>hispidulus</i>	White-water buttercup	OBL
<i>Ranunculus repens</i>	Buttercup	FACW
<i>Rosa woodsii</i>	Woods rose	FACU
<i>Rubus idaeus</i>	Wild raspberry	FACU
<i>Rumex crispus</i>	Curly Dock	FACW
<i>Salix bebbiana</i>	Bebb's willow	FACW
<i>Salix drummondiana</i>	Drummond willow	FACW
<i>Salix exigua</i>	Sandbar Willow	OBL
<i>Salix geyeriana</i>	Geyer willow	FACW+
<i>Salix lutea</i>	Yellow willow	OBL
Scirpus microcarpus	small-fruit bulrush	OBL
<i>Senecio vulgaris</i>	Common groundsel	FACU
Sium suave	water parsnip	OBL
<i>Sisymbrium altissimum</i>	Tall tumble mustard	FACU-
<i>Smilacina stellata</i>	Starry false-Solomon's-seal	FAC-
Solidago canadensis	Canada goldenrod	FACU
<i>Symphoricarpos albus</i>	Snowberry	FACU
<i>Tanacetum vulgare</i>	Common tansy	NI
<i>Taraxacum officinale</i>	Common dandelion	FACU
<i>Thlaspi arvensis</i>	Pennycress	NI
<i>Trifolium pratense</i>	Red clover	FACU
<i>Verbascum thapsus</i>	Common mullein	--
<i>Veronica Americana</i>	American speedwell	OBL

¹ **Bolded** species indicate those documented in the analysis area for the first time in 2005.

Table 2: Transect 1 data summary.

Monitoring Year	2002	2003	2004	2005
Transect Length (feet)	471	471	471	471
# Vegetation Community Transitions along Transect	4	4	4	4
# Vegetation Communities along Transect	3	3	3	3
# Hydrophytic Vegetation Communities along Transect	2	2	2	2
Total Vegetative Species	28	27	30	31
Total Hydrophytic Species	15	16	17	17
Total Upland Species	13	11	13	14
Estimated % Total Vegetative Cover	85	95	86	84
% Transect Length Comprised of Hydrophytic Vegetation Communities	59	59	59	60
% Transect Length Comprised of Upland Vegetation Communities	37	37	37	36
% Transect Length Comprised of Unvegetated Open Water	4	4	4	4
% Transect Length Comprised of Bare Substrate	0	0	0	0



Chart 2: Transect maps showing vegetation type from the start of transect (0 feet) to the end of transect (471 feet) for each year monitored.

Chart 3: Length of vegetation communities within Transect 1 for each year monitored.

3.3 Soils

The soils located at the Camp Creek site are mapped as Gallatin-shallow muck complex, gently sloping. Soil characteristics at each wetland determination point were compared with those of the Gallatin-shallow muck complex and generally matched this classification. Wetland soils observed during monitoring and documented on the Routine Wetland Determination form were mostly peat, loams, sandy loams, or sands with very low chromas (1 or 2). Mottles or oxidized rhizospheres (redoximorphic features) were not present any of the profiles. Soil profiles in the wetlands meadow mostly consisted of deep A horizons of peat or loamy materials with a sandy/gravelly layer underneath, saturated at approximately 8 inch depths. Several profiles had large cobbles, gravels and stones below a 6-8 inch A horizon with matrix colors of 10YR 2/1. Created upland slopes were constructed with fill materials removed from channel excavation. Upland soil pits consisted of a mixture of large cobbles and loamy soil, with matrix colors of 10YR 2/2.

3.4 Wetland Delineation

Delineated wetland boundaries are illustrated on **Figure 3** in **Appendix A**. Completed wetland delineation forms are included in **Appendix B**. Soils, vegetation, and hydrology are discussed in preceding sections. Approximately 47.23 wetland acres and 1.5 open water channel acres occurred within the current monitoring area prior to project implementation. Pre-project wetland locations are shown on **Figure 4** in **Appendix A**. Monitoring in 2005 identified the conditions listed in **Table 3**.

Table 3: Wetland conditions within Camp Creek Wetland Mitigation Site.

Condition	MDT Property Monitoring Area 2005 (acre)	Grasser Property Monitoring Area 2005 (acre)	MDT Property Monitoring Area 2000 Baseline (acre)	Grasser Property Monitoring Area 2000 Baseline (acre)
Wetland Area	35.13	6.93	42.61	4.62
Open Water Area	0.95	1.20	0.75	0.75
Subtotals	36.08	8.13	43.36	5.37
Total Aquatic Habitat	44.21		48.73	

Overall, the project has gained 0.65 stream acre and “lost” an estimated 5.17 wetland acres in comparison to baseline conditions. Cumulatively, approximately 42.06 wetland acres and 2.15 open water acres now occur within the monitoring area (**Figure 3 in Appendix A**), for a total of 44.21 acres of aquatic habitat. Prior to construction, the site contained approximately 47.23 acres of wetlands and 1.5 open water channel acres within the current monitoring limits. Open water channels were located in the extreme south end of the Grasser property and the in the northwest corner of the MDT property. A decrease of 2.09 acres in net wetland area was observed between 2004 and 2005. The overall cumulative change in aquatic habitat at the site since construction has been approximately $44.21 - 48.73 = (-4.52)$ acres.

During the initial 2002 monitoring, an immediate net decrease in wetland acres was observed at this mitigation site. This could have been attributable to the dry year, changes in irrigation practices, construction-related disturbance (haul routes, drive-through areas, staging areas, etc.), slight differences in pre- and post-construction delineation approaches, or a combination of all factors.

Historic irrigation practices, although unquantified, provided substantial wetland hydrology to the current MDT property. The diversion from Camp Creek onto the property was virtually uncontrolled, and water flood irrigated the site whenever the water was high enough to do so; particularly in the spring/early summer. Similarly, substantial flood irrigation was conducted on the Grasser property prior to mitigation implementation; however, most of this irrigation has now ceased due to landowner management priorities, etc. These changes in onsite and adjacent irrigation practices have had a substantive and unplanned impact on site hydrology.

Final plan designs were based on a preliminary 2000 wetland delineation. The preliminary 2000 baseline wetland delineation was smaller in acres than the baseline delineation ultimately agreed upon (post-construction) by the Corps. Consequently, some areas ultimately depicted as wetlands in the final delineation were heavily disturbed during construction efforts and were also designated as areas to deposit fill materials. Additionally, some upland areas were not created as specified in the construction plans, but were larger or in different locations. Several areas mapped during the pre-project delineation as uplands became spoil piles two to three times larger than the original size of the mapped upland.

During the 2005 monitoring, wetland acreages decreased from those observed in 2004. Wetland boundaries exhibited some change on both the MDT and Grasser owned parcels. Wetland decreases were observed along the outer reaches of some the constructed floodplain margins. The decrease of wetlands was due to the change in vegetation from mostly wetland species to

high abundance of upland species. The outer reaches of the floodplain located furthest away from the stream channel did not receive enough hydrology in 2005 to sustain wetland species. Channel down-cutting has been observed in scattered short project reaches (<100 feet long), which may be playing a role in this observed decrease. However, stream flow was sub-normal during the majority of the 2005 growing season and may have resulted in the wetland loss. Based on USGS data from the stream gauge on the Bitterroot River near Darby, flows in May, June, and July of 2005 were only 65%, 50%, and 61% of the long-term mean monthly flows for these months (**Chart 1**). Flows during 2004 were similarly low (**Chart 1**).

During 2003 and 2004, a dramatic resurgence of spotted knapweed and other upland species lead to the change in community type descriptions. Areas of heavy spotted knapweed coverage are located adjacent to and throughout the site. Disturbance from construction activities to the pre-existing seed bank, likely spreading of seed by heavy equipment, and lack of pre-project weed control could have contributed to the overall increase. It is likely that other factors such as lack of hydrology along the floodplains may be leading to the ultimate conversion of floodplains to a drier vegetation type. Thus, a combination of numerous land use (irrigation practices), construction, environmental, and baseline mapping factors likely resulted in the wetland “loss” observed at the site.

3.5 Wildlife

Wildlife species or evidence of wildlife, observed on the site during 2002, 2003, 2004 and 2005 monitoring efforts are listed in **Table 4**. Specific evidence observed, as well as activity codes pertaining to birds, is provided on the completed monitoring form in **Appendix B**.

This site provides habitat for a variety of wildlife species, although this was not necessarily reflected in the 2005 monitoring data. One mammal, two fish, one amphibian and three bird species were noted at the mitigation site during the 2005 site visits.

The newly constructed channel offers habitat for several fish species, including westslope cutthroat and brook trout. Pre-project and post-project surveys along Camp Creek were conducted by the Montana Fish Wildlife and Parks during 1999, 2003 and 2004 (**Chart 4**). The 2004 surveys documented 163 westslope cutthroat trout ranging in size from 3 to 8 inches (MFWP 2004). The 2003 surveys documented 300 westslope cutthroat trout ranging in size from 3 to 12 inches and also several small sized brook trout (MFWP 2004). The majority of fish observed were in the 3 to 6 inch size class, which is expected for new habitat because smaller fish usually colonize these areas first.


Table 4: Fish and wildlife species observed at the Camp Creek Mitigation Site during 2002-2005 monitoring.

FISH
Brook Trout (<i>Salvelinus fontinalis</i>)¹ Westslope Cutthroat Trout (<i>Oncorhynchus clarki lewisi</i>)¹
AMPHIBIANS
Spotted Frog (<i>Rana luteiventris</i>)
REPTILES
None
BIRDS
American Crow (<i>Corvus brachyrhynchos</i>) American Dipper (<i>Cinclus mexicanus</i>) American Goldfinch (<i>Carduelis tristis</i>) American Robin (<i>Turdus migratorius</i>) Black-billed Magpie (<i>Pica pica</i>) Blue Grouse (<i>Dendragapus obscurus</i>)² Brewer's Blackbird (<i>Euphagus cyanocephalus</i>) Brown-headed Cow bird (<i>Molothrus ater</i>) Canada Goose (<i>Branta canadensis</i>) Cedar Waxwing (<i>Bombycilla cedrorum</i>) Common Raven (<i>Corvus corax</i>) European Starling (<i>Sturnus vulgaris</i>) Grasshopper Sparrow (<i>Ammodramus savannarum</i>) Killdeer (<i>Charadrius vociferus</i>) Mallard (<i>Anas platyrhynchos</i>) Northern Harrier (<i>Circus cyaneus</i>)² Red-tail Hawk (<i>Buteo jamaicensis</i>)
MAMMALS
Bobcat (<i>Felis rufus</i>) Coyote (<i>Canis latrans</i>) Deer (<i>Odocoileus spp.</i>) Deer Mouse (<i>Peromyscus maniculatus</i>)² Elk (<i>Cervus elaphus</i>) Meadow Vole (<i>Microtus pennsylvanicus</i>)² Moose (<i>Alces alces</i>) Red Fox (<i>Vulpes vulpes</i>)²

¹ Survey conducted by Montana Fish, Wildlife & Parks.

² Observed by MDT


Bolded species were observed during 2005 monitoring. All other species were observed during one or more of the previous monitoring years, but not during 2005.

Chart 4: Westslope cutthroat trout survey for Camp Creek (MFWP 2004).

3.6 Macroinvertebrates

Complete results from the macroinvertebrate sampling location (**Figure 2**) are presented in **Appendix F**. Sampling points were located along one area of the creek. The following analysis was provided by Rhithron Associates (Bollman 2005). Macroinvertebrate sampling results were summarized by Rhithron Associates in the italicized section below (Bollman 2005) and in **Chart 5**.

An invertebrate assemblage characteristic of a trout stream was collected at the Camp Creek site. Two cold stenotherm taxa were present in the sample, and the mayfly taxa richness was high. The biotic index value, however, was elevated compared to expectations for a cold-water stream. This suggests that water quality may have been impaired at this site. "Clinger" richness was high, and caddisfly taxa richness was within expectations. These findings imply that sediment deposition did not appreciably limit access to stony substrate habitats. Stonefly taxa richness was blunted; this could be related to reach-scale habitat disruption. Gatherers dominated the functional mix, a finding consistent with mild impairment of water quality. All expected feeding groups, however, were collected at the site. The sampled site at Camp Creek cannot be considered a wetland, since rheophilic taxa characteristic of rapid flow conditions and cold water are supported at the site. Scores indicated in the chart were derived by means of the Montana DEQ index for Valley and Foothill Prairie streams (Bollman 1998). The score calculated for 2005 indicates slight impairment, and stable instream and reach-scale habitat conditions between 2002 and 2005.

Chart 5: Bioassessment scores for Camp Creek.

3.7 Functional Assessment

Completed 2005 functional assessment forms are included in **Appendix B**. Per Corps of Engineers direction (Steinle pers. comm.), separate functional assessments were completed for the Grasser and MDT parcels. Because of this “artificial”, ownership boundary-based separation of assessment areas (AAs), and based on development of the site and discussions with the Corps, the MDT parcel, formerly split into two AAs, was assessed in its entirety as one contiguous AA. Results are presented in **Table 5**.

The AA on the MDT parcel rated as Category II (high value). This overall rating was primarily due to high ratings for MNHP species habitat (documented primary habitat for westslope cutthroat trout based on 2003 and 2004 fish surveys conducted by Montana Department of Fish, Wildlife and Parks), surface water storage, production export / food chain support, groundwater discharge/recharge, and recreation/education ratings (public ownership with excellent access). Remaining parameters generally rated as moderate.

In 2004-2005, the MDT site received a moderate sediment/shoreline stabilization rating due to the increase in species with deep binding roots along the streambank. Shoreline species during evaluation consisted of grasses and willow sprigs; an increase in willow cover between 2003 and 2005 monitoring increased the functional rating for sediment/shoreline stabilization category. Over time, willow sprigs will develop into larger, even more robust shrubs with extensive deep binding roots systems. Enhancement of both wetland and upland vegetation should increase wildlife usage throughout the site.

The AA on the Grasser parcel is subject to a higher degree of disturbance (it is not within a conservation easement), and rated as Category III (moderate value). This AA received high ratings for MNHP species habitat (again due to west-slope cutthroat trout), production export / food chain support, and groundwater discharge/recharge. All other parameters rated low to moderate.

Pre-project (2001) and post-project (2005) wetland assessment scores are presented in **Table 5**. Turnstone Biological conducted the initial functional assessments for the Camp Creek Mitigation Site, and separated the site into three assessment areas: emergent (Type I), scrub-shrub emergent (Type II), and rock bottom with narrow mixed wetland fringe (Type III) wetland classifications.

Overall, ratings have increased substantially on the MDT parcel for MNHP species habitat (west-slope cutthroat trout), general wildlife habitat, general fish habitat, surface water storage, sediment/shoreline stabilization, production export/food chain support, uniqueness, and recreation/education potential. Rating have increased in the Grasser parcel for MNHP species habitat (west-slope cutthroat trout), general fish habitat, surface water storage, and sediment/shoreline stabilization

Approximately 141.68 functional units (score x wetland acreage) have been gained thus far at the Camp Creek mitigation site, despite the decrease in wetland acres between pre-project and post-project assessments on the MDT parcel. Approximately 113.24 functional units have been gained at the MDT parcel, and nearly 28.44 have been gained on the Grasser parcel. Refer to **Table 6** for details.

3.8 Photographs

Representative photographs taken from photo-points and transect ends are presented in **Appendix C**.

3.9 Revegetation

Upon completion of the new channel and floodplain construction, revegetation efforts were conducted in 2002 to enhance riparian and upland habitat. The streambanks were seeded with a grass mix designed by an MDT botanist and 20,480 willow cuttings were sprigged through the fabric work. Floodplain areas were planted with a mixture of native shrubs & trees associated with local riparian corridors. These included aspen, alder, black cottonwood, dogwood and willows. Upland slopes were planted with Douglas fir, lodgepole pine, ponderosa pine, serviceberry, shrubby potentilla, snowberry, and woods rose.

Species survival data is presented in **Appendix B**. The belt transect used for vegetation monitoring was also used as the survival transect. A second survival transect was added to the south of the vegetation transect across the created and planted upland berms. A third survival transect was added in 2003 to assess the channel and floodplain vegetation enhancements.

Survival rates within the upland areas were similar to those observed during the 2004 monitoring. In 2003, a majority of the survival rates ranged from 70% to 100%. Survival data recorded in 2004 showed seven out of 13 species had a survival rate below 50 %. These included such species as woods rose, ponderosa pine, snowberry, shrubby potentilla and red-osier dogwood. Almost all the Douglas-fir observed had died after initial planting.

Table 5: Summary of 2001 (baseline) and 2005 wetland function/value ratings and functional points ¹ at Camp Creek.

Function and Value Parameters From the 1999 MDT Montana Wetland Assessment Method	2001 Type I, MDT Property	2001 Type III, MDT Property	2001 Type I, Grasser Property	2001 Type II, Grasser Property	2001 Type III, Grasser Property	2005 Grasser Property	2005 MDT Property
Listed/Proposed T&E Species Habitat	Mod (0.8)	Mod (0.8)	Mod (0.8)	Mod (0.8)	Mod (0.8)	Mod (0.8)	Mod (0.8)
MNHP Species Habitat	Low (0.1)	Low (0.1)	Low (0.1)	Low (0.1)	Low (0.1)	High (0.8)	High (0.8)
General Wildlife Habitat	Low (0.3)	Mod (0.5)	Low (0.3)	Mod (0.5)	Mod (0.5)	Mod (0.5)	Mod (0.7)
General Fish/Aquatic Habitat	Low (0.1)	Mod (0.5)	Low (0.1)	Low (0.1)	Mod (0.5)	Mod (0.7)	Mod (0.7)
Flood Attenuation	Mod (0.6)	Mod (0.4)	Mod (0.6)	Mod (0.5)	Mod (0.4)	Mod (0.4)	Mod (0.6)
Short and Long Term Surface Water Storage	Low (0.3)	High (0.8)	Low (0.3)	Low (0.3)	High (0.8)	Mod (0.6)	High (1.0)
Sediment, Nutrient, Toxicant Removal	Mod (0.7)	Mod (0.6)	Mod (0.7)	Mod (0.7)	Mod (0.6)	Mod (0.6)	Mod (0.7)
Sediment/Shoreline Stabilization	Low (0.2)	Low (0.3)	Low (0.2)	Mod (0.6)	Low (0.3)	Low (0.3)	Mod (0.7)
Production Export/Food Chain Support	Mod (0.7)	High (0.9)	Mod (0.7)	Mod (0.7)	High (0.9)	High (0.9)	High (0.9)
Groundwater Discharge/Recharge	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)
Uniqueness	Low (0.1)	Low (0.2)	Low (0.1)	Low (0.3)	Low (0.2)	Low (0.2)	Mod (0.4)
Recreation/Education Potential	Low (0.2)	Low (0.1)	Low (0.2)	Low (0.3)	Low (0.1)	Low (0.3)	High (1.00)
Actual Points/Possible Points	5.1 / 12	6.2 / 12	5.1 / 12	5.9 / 12	6.2 / 12	7.1 / 12	9.3 / 12
% of Possible Score Achieved	42%	52%	42%	49%	52%	59%	78%
Overall Category	III	III	III	III	III	III	II
Total Acreage of Assessed Wetlands and Open Water within Easement	42.3	1.06²	3.51²	0.50²	1.36²	8.13	36.08
Functional Units (fu) (acreage x actual points)	215.73	6.57	17.90	2.95	8.43	57.72	335.54
Functional Unit Gain to Date by Ownership (fu)	NA	NA	NA	NA	NA	28.44³	113.24³
Total Functional Unit Gain to Date (fu)	NA	NA	NA	NA	NA	141.68	

¹ see completed functional assessment forms in **Appendix B** for further detail.² Baseline acreages adjusted per subsequent study; see Section 2.5.³ Baseline Functional Units used to determine the 2005 Functional Unit Gain included the combined totals for the 2001 MDT (222.30 fu) and Grasser (29.28 fu) properties.

The wetter species planted along the streambank and floodplain margins had a much higher survival rate ranging from 60% to 90%. These included alder, aspen, cottonwood and willows. The willow sprigs are spreading out along the banks, increasing in sizes and density each growing season. Several other planted shrubs had increased in overall stature and exhibited vigorous growth.

These survival rates are based on a low number of total observations and might misrepresent the true survival rate. The current survival rates are based on the occurrences recorded during the 2002 monitoring. The 2002 planting specifications are presented in **Appendix G**.

3.10 Maintenance Needs/Recommendations

Per Corps recommendations, the potential for enhancing the surface connection between Camp Creek and the large emergent complex on the MDT parcel was investigated. Based on field survey investigations, a shallow flood channel was excavated during fall 2005 between the creek and existing swales to enhance the connectivity of these two systems during high water events.

Figure 6 (Appendix B) illustrates the location and approximate cross-sectional view of the channel.

Several Category 1 Noxious weeds are present on both MDT and Grasser parcels including Canada thistle, hound's-tongue, oxeye daisy and spotted knapweed. Weed control and re-vegetation of disturbed sites is needed to prevent further weed spread, reduce the risk of new weeds invading, reduce wind and water erosion and reduce sediment input to surface waters. Survival of plantings will continue to be monitored, and supplemental planting may need to be implemented if success of current plantings is low.

The MDT parcel has the least amount of invasive species and distribution is limited to upland areas not affected during construction efforts. Control measures for these areas should be implemented to avoid potential spread of invasive species into the wetland areas. Planted upland areas within the MDT parcel which were observed to have a low survival rates should be replanted with appropriate native plant stock, and irrigated.

The Grasser parcel supports the majority of the noxious weed species with extensive distribution along the floodplain corridor. A weed management plan for this site should be developed and implemented to control the spread of noxious weeds. Areas of invading spotted knapweed located along floodplain margins should be controlled and reseeded or planted with appropriate wetland species to help control further spread of invasive species.

3.11 Current Credit Summary

As of 2005, the project has gained 0.65 stream acre and “lost” an estimated 5.17 wetland acres in comparison to baseline conditions. Cumulatively, approximately 42.06 wetland acres and 2.15 open water acres now occur within the monitoring area (**Figure 3, Appendix A**), for a total of 44.21 acres of aquatic habitat. Prior to construction, the site contained approximately 47.23 acres of wetlands and 1.5 open water channel acres within the current monitoring limits. Open water channels were located in the extreme south end of the Grasser property and in the

northwest corner of the MDT property. A 2.09-acre decrease in the net wetland area was observed between 2004 and 2005. The overall cumulative change in aquatic habitat at the site since construction has been approximately $44.21 - 48.73 = (-4.52)$ acres.

This “decrease” in wetland acreage could be attributable to several factors. However, a primary cause is thought to be the virtual termination of flood irrigation on both the MDT and adjacent Grasser parcels, and low stream flows during 2004 and 2005, which appear to have had a substantive impact on site hydrology. Other possible causes include long-term drought, scattered channel adjustments (down-cutting), fire, construction-related disturbance, slight differences in pre- and post-construction delineation approaches, or a combination of all factors.

Despite the apparent decrease in wetland acreage, approximately 141.68 functional units (score x wetland acreage) have been gained thus far at the Camp Creek mitigation site. Approximately 113.24 functional units have been gained at the MDT parcel, and 28.44 have been gained on the Grasser parcel.

A final method of credit allocation for this site is being worked out between MDT and COE, and will be based upon this monitoring data and other information. As such, the current amount of credit applicable to this site is undetermined. However, one approach under consideration pertains to the use of functional units, whereby wetland acreage for each AA is multiplied by the total score for that AA to arrive at an overall functional unit score. This is done both pre-project and post-project. The difference between these two numbers (the functional unit “gain”) is then divided by the post-project score to arrive at an approximate credit acreage for that AA. Credit acreages from each AA are summed to arrive at a total for the site. This approach is illustrated below in **Table 6**. Using this approach, a current maximum of approximately 16.17 credit acres could be assigned to the Camp Creek site as of 2005.

Table 6: Potential functional unit-based credit for Camp Creek Mitigation Project.

Property	2005 Wetland & Channel Acreage	2005 Score	2005 Functional Units	Baseline Functional Units	Functional Unit “Gain”	“Gain” Divided by Current Score (potential credit acres)
MDT	36.08	9.3	335.54	222.30	113.24	12.17
Grasser	8.13	7.1	57.72	29.28	28.44	4.00
Total	44.21	--	393.26	251.58	141.68	16.17

4.0 REFERENCES

- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. U.S. Army Corps of Engineers, Washington, DC.
- Ralph, C.J., Geupel, G.R., Pyle, P., Martin, T.E., and D.F. DeSante. 1993. *Handbook of field methods for monitoring landbirds*. Gen. Tech. Rep. PSW-GTR-144. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Dept. of Agriculture. 41 p.
- Montana Department of Fish Wildlife and Parks (MFWP). 2004. Camp Creek Fish Survey. Personal Communication: Chris Clancy (MFWP)
- Steinle, A. 2004. Montana Program Manager, U.S. Army Corps of Engineers, Helena Regulatory Office, Helena, Montana. May 6th meeting.
- USDA Soil Conservation Service. 1951. *Soil Survey of Bitterroot Valley Area, Montana*.
- USDA Natural Resources Conservation Service. 1998. *Field Indicators of Hydric Soils in the United States*, Version 4. G. Hurt, P. Whited and R. Pringle (eds.). USDA, NRCS Fort Worth, TX.

Appendix A

FIGURES 2, 3, 4, AND 6

*MDT Wetland Mitigation Monitoring
Camp Creek
Sula, Montana*

Figure 2 - Monitoring Activity Locations






Figure 3 - Mapped Site Features 2005



PROJECT NAME		MDT Camp Creek Wetland Mitigation	
DRAWING TITLE		Mapped Site Features 2005	
PROJ NO:	330054.106	DRAWN:	R.A.I.L.
FILE NAME:	CAMPOR2005	CHECKED:	B.D.
SCALE:	1"=300ft	APPVD:	G.H.
LOCATION:	Sula, Mt.	PROJECT MGR:	B.Dutton
LAND & WATER CONSULTING, INC.		FIGURE	
P.O. BOX 8254		F3	
Missoula, MT 59807		REV -	
		DATE: 11-11-05	

Figure 4 - Pre-Developed Wetland





LEGEND

- Monitoring Area Limits
- Pre Developed Wetland Area Limits Within Monitoring Limits
- Base Photograph Date: July 23, 2002

Pre Developed Wetland Area : (within monitoring area limits)
 Gross Area (2001) MDT 45.59 + Grasser 5.37 = 50.96 Acres
 Open Water Grasser 0.75 Acres+ MDT 0.75 Acres= 1.50 Acres
 Upland Islands MDT-2.23 Acres Grasser -0.00 = 2.23 Acres
 Net Wetland Area(2001) 47.23 Acres

PROJECT NAME		MDT Camp Creek Wetland Mitigation	
DRAWING TITLE		Pre-Developed Wetland	
PROJ NO:	B43054.00 0106	DRAWN:	R.A.
FILE NAME:	TASK106BASE	CHECKED:	J.B.
SCALE:	1"=300ft	APPVD:	J.B.
LOCATION:	Sula, Mt.	PROJ MGR:	J.B.
LAND & WATER CONSULTING, INC.		FIGURE	
P.O. BOX 8254		F4	
Missoula, MT 59807		REV -	
		DATE: 6-13-05	



Typical Section of Excavated Flood Channel

PROJECT NAME		MDT Camp Creek Wetland Mitigation	
DRAWING TITLE		Flood Channel Plan and Typical	
PROJ NO: 130091	DRAWN: R.A.	FILE NAME: TASK38BASE2003	CHECKED: B.D.
SCALE: 1"=150ft	APPVD: G.H.	LOCATION: Sula, Mt.	PROJ MGR: B. Dutton
LAND & WATER CONSULTING, INC.		FIGURE	
P.O. BOX 8254		F6	
Missoula, MT 59807		REV	
		DATE: 10-28-03	

Appendix B

2005 WETLAND MITIGATION SITE MONITORING FORM

2005 BIRD SURVEY FORM

2005 WETLAND DELINEATION FORMS

2005 FUNCTIONAL ASSESSMENT FORM

MDT Wetland Mitigation Monitoring

Camp Creek

Sula, Montana

LWC / MDT WETLAND MITIGATION SITE MONITORING FORM

Project Name: Camp Creek Project Number: B43054.00 - 0106 Assessment Date: 08/01/05
Location: Sula Valley MDT District: Lower Clark Fork Milepost:
Legal description: T 1 N R 19 W Section 27 & 34 Time of Day: Morning to early afternoon
Weather Conditions: Cloudy & overcast Person(s) conducting the assessment: Greg Howard
Initial Evaluation Date: 09/05/02 Visit #: 3 Monitoring Year: 2005
Size of evaluation area: 200 acres Land use surrounding wetland: Agriculture; livestock grazing & pasture

HYDROLOGY

Surface Water Source: Camp Creek

Inundation: Present Absent X Average depths: - ft Range of depths: - ft

Assessment area under inundation: %

Depth at emergent vegetation-open water boundary: ft

If assessment area is not inundated are the soils saturated w/in 12" of surface: Yes X No

Other evidence of hydrology on site (drift lines, erosion, stained vegetation etc.): Hydrology on this site is mostly comes from Camp Creek. Surface and groundwater observed in areas of lower topography, undisturbed wetland meadows.

Groundwater

Monitoring wells: Present Absent: X

Record depth of water below ground surface

Well #	Depth	Well #	Depth	Well #	Depth

Additional Activities Checklist:

X Map emergent vegetation-open water boundary on air photo

X Observe extent of surface water during each site visit and look for evidence of past surface water elevations (drift lines, erosion, vegetation staining etc.)

 GPS survey groundwater monitoring wells locations if present

COMMENTS/PROBLEMS: Vegetation cover along floodplains and creek margins dominated by wetland species. Certain areas of the floodplain saturated throughout the season. The planted shrubs and trees showing new growth for this season. Willows sprigged along the streambanks spreading by rhizomes. Vegetation community types and mapping remained similar to 2004 monitoring. Grassier parcel remained similar to 2004 with higher coverage values of spotted knapweed and other undesirable species within floodplain margins.

VEGETATION COMMUNITIES

Community No.: 1 Community Title (main species): Agropyron / Trifolium (Created upland)

Dominant Species	% Cover	Dominant Species	% Cover
<i>Agropyron repens</i>	50	Planted Species	10
<i>Thlaspi arvensis</i>	P	<i>Trifolium pratense</i>	P
<i>Sisymbrium altissimum</i>	20	<i>Centaurea maculosa</i>	20
<i>Lychnis alba</i>	P	<i>Alopecurus pratensis</i>	P
<i>Agrostis alba</i>	P		

COMMENTS/PROBLEMS: Created uplands, planted with several drier species: *Pinus ponderosa*, *Pseudotsuga menziesii*, *Symphoricarpos albus*, *Rosa woodsii*, *Potentilla fruticosa*, and *Amelanchier alnifolia*. Community No. 1 with similar condition as found in 2004 monitoring, except for an increase in *Sisymbrium altissimum* and *Centaurea maculosa*.

Community No.: 2 Community Title (main species): Carex / Phalaris (Undisturbed wetland)

Dominant Species	% Cover	Dominant Species	% Cover
<i>Carex aquatilis</i>	P	<i>Alopecurus pratensis</i>	P
<i>Phalaris arundinacea</i>	20	<i>Phleum pratense</i>	P
<i>Carex utriculata</i>	20	<i>Agrostis alba</i>	P
<i>Carex nebrascensis</i>	50	<i>Sisymbrium altissimum</i>	10
<i>Geum macrophyllum</i>	P		

COMMENTS/PROBLEMS: Open wetland meadow with extensive sedges, intermixed with a few drier grass species.

Community No.: 3 Community Title (main species): Agrostis / Deschampsia (Floodplain / Streambank)

Dominant Species	% Cover	Dominant Species	% Cover
<i>Alopecurus pratensis</i>	P	<i>Carex nebrascensis</i>	P
<i>Populus trichocarpa</i> - Planted	10	<i>Phalaris arundinacea</i>	10
<i>Populus tremuloides</i> - Planted	10	<i>Phleum pratense</i>	T
<i>Mimulus guttatus</i>	T	<i>Salix</i> – sprigged	20
<i>Agrostis alba</i>	30	<i>Alnus incana</i>	10
<i>Deschampsia cespitosa</i>	20	<i>Carex crawfordii</i>	10
<i>Glyceria grandis</i>	10	<i>Carex stipata</i>	P

COMMENTS/PROBLEMS: Vegetation communities along streambanks and floodplain are further transitioning from upland to wetland species. Coverage of wetland species continues to increase with each season. Saturated soils observed along much of the floodplain margin. Shrubs and trees installed during first year have increased in cover and show more new growth. Heights of several planted shrubs and trees ranging from 3-4 ft. tall. Some areas along the floodplain previously mapped as wetland have been changed to upland, mostly areas along the outer fringes of floodplain.

Additional Activities Checklist:

X Record and map vegetative communities on air photo

VEGETATION COMMUNITIES (continued)

Community No.: 4 Community Title (main species): Surface flow within channel

Dominant Species	% Cover	Dominant Species	% Cover
<i>Ranunculus aquatilis</i> var. <i>hispidulus</i>	10		

COMMENTS/PROBLEMS: Vegetation Community No. 4 combined with No. 3. Aquatic vegetation invading areas of slower water along banks. Volunteer *Populus trichocarpa* seedlings along cobble banks.

Community No.: 5 Community Title (main species): Agropyron / Centaurea

Dominant Species	% Cover	Dominant Species	% Cover
<i>Centaurea maculosa</i>	60	<i>Agropyron repens</i>	20
<i>Sisymbrium altissimum</i>	P	<i>Linaria vulgaris</i>	P
<i>Bromus inermis</i>	10		
<i>Bromus tectorum</i>	10		
<i>Alopecurus pratensis</i>	P		

COMMENTS/PROBLEMS: Upland slopes observed on both the east and west sides of site. Area dominated by spotted knapweed and several other pasture grasses such as smooth brome and quackgrass.

Community No.: 6 Community Title (main species): Populus / Salix

Dominant Species	% Cover	Dominant Species	% Cover
<i>Populus trichocarpa</i>	30	<i>Rosa woodsii</i>	10
<i>Salix bebbiana</i>	P	<i>Symphoricarpos albus</i>	P
<i>Alnus incana</i>	P	<i>Salix drummondiana</i>	P
<i>Salix geyeriana</i>	10	<i>Salix exigua</i>	P
<i>Cornus stolonifera</i>	T		

COMMENTS/PROBLEMS: Mature cottonwood and shrub communities found along the old channel.

Community No.: 7 Community Title (main species): Centaurea / Phalaris

Dominant Species	% Cover	Dominant Species	% Cover
<i>Phalaris arundinacea</i>	20	<i>Taraxacum officinale</i>	T
<i>Centaurea maculosa</i>	50	<i>Chrysanthemum leucanthemum</i>	P
<i>Verbascum thapsus</i>	T	<i>Trifolium pratense</i>	P
<i>Bromus inermis</i>	P	<i>Rumex crispus</i>	T
<i>Agropyron repens</i>	20	Plantings	10

COMMENTS/PROBLEMS: Vegetation type found along the upland fringes of constructed floodplain on Grasser-owned parcels. Community No. 7 dominated by spotted knapweed. Aggressive reed canarygrass also invading in many areas with spotted knapweed.

COMPREHENSIVE VEGETATION LIST

Species	Vegetation Community Number(s)	Species	Vegetation Community Number(s)
<i>Achillea millefolium</i>	1,5	<i>Lepidium perfoliatum</i>	1
<i>Agropyron repens</i>	1,3,5,7	<i>Linaria vulgaris</i>	1,7
<i>Agrostis alba</i>	2,3	<i>Lonicera involucrata</i>	6
<i>Alnus incana</i>	6	<i>Lupinus wyethii</i>	1
<i>Alopecurus pratensis</i>	2,3,5	<i>Lychnis alba</i>	1
<i>Amelanchier alnifolia</i>	1	<i>Matricaria matricarioides</i>	1
<i>Aster integrifolius</i>	1	<i>Melilotus officinalis</i>	1,5
<i>Betula occidentalis</i>	3	<i>Mentha arvensis</i>	2,3
<i>Bromus inermis</i>	5,7	<i>Mimulus guttatus</i>	3
<i>Bromus tectorum</i>	1,5	<i>Phalaris arundinacea</i>	2,3,7
<i>Calamagrostis canadensis</i>	2	<i>Phleum pratense</i>	2,3
<i>Carex aquatilis</i>	2	<i>Pinus ponderosa</i>	1
<i>Carex bebbii</i>	2	<i>Plantago major</i>	1,3
<i>Carex crawfordii</i>	3	<i>Poa pratensis</i>	1,5
<i>Carex lanuginosa</i>	2,3	<i>Polygonum amphibium</i>	2
<i>Carex nebrascensis</i>	2,3	<i>Populus tremuloides</i>	3,4
<i>Carex praegracilis</i>	2	<i>Populus trichocarpa</i>	3,6
<i>Carex utriculata</i>	2	<i>Potentilla fruticosa</i>	1
<i>Centaurea maculosa</i>	1,5,7	<i>Potentilla gracilis</i>	1
<i>Cercocarpus ledifolius</i>	1	<i>Pseudotsuga menziesii</i>	1
<i>Chenopodium album</i>	1,3	<i>Ranunculus aquatilis</i> var. <i>hispidulus</i>	4
<i>Chrysanthemum leucanthemum</i>	1,5,7	<i>Ranunculus repens</i>	2
<i>Cirsium arvense</i>	1	<i>Rosa woodsii</i>	1,6
<i>Cirsium vulgare</i>	1,2	<i>Rubus idaeus</i>	6
<i>Cornus stolonifera</i>	3,6	<i>Rumex crispus</i>	1,2,3,7
<i>Crataegus douglasii</i>	1	<i>Salix bebbiana</i>	6
<i>Crepis tectorum</i>	1	<i>Salix drummondiana</i>	4
<i>Cynoglossum officinale</i>	1	<i>Salix exigua</i>	2,3,4
<i>Danthonia</i> spp.	1	<i>Salix geyeriana</i>	4,6
<i>Deschampsia cespitosa</i>	2,3	<i>Salix lutea</i>	3
<i>Epilobium ciliatum</i>	2,3	<i>Scirpus microcarpus</i>	3,4
<i>Epilobium paniculatum</i>	2,3	<i>Senecio vulgaris</i>	1
<i>Equisetum arvense</i>	2,3	<i>Sium suave</i>	3
<i>Equisetum laevigatum</i>	2,3	<i>Sisymbrium altissimum</i>	1,5
<i>Festuca pratensis</i>	1	<i>Smilacina stellata</i>	2
<i>Geum macrophyllum</i>	2,3	<i>Solidago canadensis</i>	2,3
<i>Glyceria elata</i>	2	<i>Symphoricarpos albus</i>	1,5
<i>Glyceria grandis</i>	3	<i>Tanacetum vulgare</i>	2,3
<i>Gnaphalium palustre</i>	1	<i>Taraxacum officinale</i>	1,2,3,4,5,7
<i>Juncus balticus</i>	2	<i>Thlaspi arvensis</i>	1,3,5
<i>Juncus bufonius</i>	2,3	<i>Trifolium pratense</i>	1,7
<i>Juncus confusus</i>	3	<i>Verbascum thapsus</i>	1,3,5,7
<i>Juncus ensifolius</i>	2,3	<i>Veronica americana</i>	2
<i>Lactuca serriola</i>	1		

COMMENTS/PROBLEMS: Six new species added to the list for 2005. These included common monkey-flower (*Mimulus guttatus*), small-fruit bulrush (*Scirpus microcarpus*), wooly sedge (*Carex lanuginosa*), water parsnip (*Sium suave*), thickstem aster (*Aster integrifolius*) and Canada goldenrod (*Solidago canadensis*).

PLANTED WOODY VEGETATION SURVIVAL

[illegible]

COMMENTS/PROBLEMS: Three transects were used to assess overall survival. Transect 1 was located along the same line as the vegetation monitoring transect, using the same belt width. The transect no. 2, starts at the beginning of transect no. 1, running towards the east (45°), approximately 165 ft long. Transect no. 2 bisects an area of created uplands and associated drier species plantings. Transect 3 was located along floodplain margins near vegetation transect.

BIRDS

MAMMALS AND HERPTILES

Additional Activities Checklist:

COMMENTS/PROBLEMS: Macroinvertebrate samples taken at one location along the Camp Creek.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

PHOTOGRAPHS

Using a camera with a 50 mm lenses and color film take photographs of the following permanent reference points listed in the checklist below. Record the direction of the photograph using a compass. (The first time at each site establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3' above ground, survey the location with a resource grade GPS and mark the location on the air photo.)

Checklist:

- ☒ One photo for each of the 4 cardinal directions surrounding wetland
- ☒ At least one photo showing upland use surrounding wetland – if more than one upland use exists, take additional photos
- ☒ At least one photo showing buffer surrounding wetland
- ☒ One photo from each end of vegetation transect showing transect

Location	Frame #	Photograph Description	Compass Reading
1	R1 16	Looking north at transect end.	0°
1	R1 17	Looking south, uplands w/plantings.	180°
1	R1 18	Looking west, Hwy 93 and created uplands.	270°
1	R1 19	Looking northwest, upland and floodplain.	315°
2	R1 20	Looking southwest at start of vegetation transect.	225°
3	R1 21-22	Looking north along transect line.	0°
4	R1 23	Looking northwest, downstream along channel.	315°
4	R1 24	Looking south, upstream along channel.	180°
4	R1 25	Looking north, curve in creek, fabric failure.	0°
5	R1 26-31	Looking south to north, panoramic of channel & floodplain.	180° – 0°
6	R1 32	Looking east along survival transect.	45°
7	R1 34-35	Looking south, lower section, creek leaving MDT parcel.	180°
8	R2 1-5	Looking east, panoramic from west side.	180° – 0°
9	R2 6-8	Looking north, main channel entering culvert.	270° – 0°
9	R2 9-12	Looking south, main channel entering culvert.	135° – 225°
10	R2 13-14	Looking south, channel and floodplain.	180° – 225°
10	R2 15	Looking north, channel and floodplain.	0°
11	R2 16-19	Looking north, channel and floodplain, upper culvert.	0° – 315°
12	R2 20	Looking south, channel and floodplain, Grasser parcel.	180° – 225°
13	R2 21	Looking south, channel & floodplain.	180°
14	R2 22	Looking north, creek entering Grasser parcel.	225°

COMMENTS/PROBLEMS:

GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points with the GPS unit set at 5 second recording rate. Record file numbers for site in designated GPS field notebook

Checklist:

- ☐ Jurisdictional wetland boundary
- ☐ 4-6 landmarks recognizable on the air photo
- ☐ Start and end points of vegetation transect(s)
- ☐ Photo reference points
- ☐ Groundwater monitoring well locations

COMMENTS/PROBLEMS: GPS surveying completed during first year monitoring.

WETLAND DELINEATION

(Attach Corps of Engineers delineation forms)

At each site conduct the items on the checklist below:

- ☒ Delineate wetlands according to the 1987 Army Corps manual.
- ☒ Delineate wetland-upland boundary on the air photo
- ☐ Survey wetland-upland boundary with a resource grade GPS survey

COMMENTS/PROBLEMS: _____

FUNCTIONAL ASSESSMENT

(Complete and attach full MDT Montana Wetland Assessment Method field forms; also attach abbreviated field forms, if used)

COMMENTS/PROBLEMS: Functional assessments similar to 2004 monitoring. No dramatic changes or difference between monitoring periods, similar conditions exist.

MAINTENANCE

Were man-made nesting structures installed at this site? YES___ NO X

If yes, do they need to be repaired? YES___ NO___

If yes, describe problems below and indicate if any actions were taken to remedy the problems.

Were man-made structures built or installed to impound water or control water flow into or out of the wetland?
YES X NO___

If yes, are the structures working properly and in good working order? YES X NO ___

If no, describe the problems below.

COMMENTS/PROBLEMS: _____

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: Camp Creek Date: 08/01/05 Examiner: Greg Howard Transect # 1

Approx. transect length: 471 ft Compass Direction from Start (Upland): 225°

Vegetation type 1:	Agropyron / Trifolium (Community No. 1)	
Length of transect in this type:	111	feet
Species:	Cover:	
Agropyron repens	50	
Thlaspi arvensis	P	
Potentilla fruticosa	P	
Agrostis alba	10	
Cirsium vulgare	T	
Alopecurus pratensis	10	
Matricaria matricarioides	T	
Sisymbrium altissimum	T	
Phalaris arundinacea	T	
Centaurea maculosa	P	
Lychnis alba	T	
Total Vegetative Cover:		80%

Vegetation type 2:	Carex / Phalaris (Community No. 2)	
Length of transect in this type:	102	feet
Species:	Cover:	
Carex nebrascensis	70	
Carex utriculata	T	
Phalaris arundinacea	10	
Geum macrophyllum	T	
Cirsium arvense	P	
Lychnis alba	P	
Agrostis alba	P	
Salix exigua	P	
Sisymbrium altissimum	P	
Cirsium vulgare	P	
Trifolium pratense	T	
Total Vegetative Cover:		90%

Vegetation type 3:	Agropyron / Trifolium (Community No. 1)	
Length of transect in this type:	63	feet
Species:	Cover:	
Carex nebrascensis	P	
Chrysanthemum leucanthemum	T	
Epilobium ciliatum	P	
Agropyron repens	T	
Festuca pratensis	30	
Phalaris arundinacea	P	
Sisymbrium altissimum	T	
Cirsium arvense	T	
Centaurea maculosa	T	
Verbascum thapsus	T	
Deschampsia cespitosa	20	
Poa pratensis	10	
Total Vegetative Cover:		70%

Vegetation type 4:	Agrostis / Deschampsia (Community No. 3)	
Length of transect in this type:	16	feet
Species:	Cover:	
Carex utriculata	T	
Epilobium ciliatum	P	
Agrostis alba	20	
Mentha arvensis	T	
Alopecurus pratensis	P	
Juncus ensifolius	P	
Trifolium pratense	20	
Carex nebrascensis	T	
Deschampsia cespitosa	20	
Plantings (Populus tremuloides & Populus trichocarpa)	P	
Willow Sprigs	20	
Phalaris arundinacea	P	
Total Vegetative Cover:		90%

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: Camp Creek Date: 08/17/04 Examiner: Greg Howard Transect # 1

Approx. transect length: 471 ft Compass Direction from Start (Upland): 225°

[illegible][illegible]

Vegetation type 6:	Agrostis / Deschampsia (Community No. 3)	
Length of transect in this type:	164	feet
Species:	Cover:	
Carex utriculata	T	
Epilobium ciliatum	T	
Agrostis alba	20	
Alopecurus pratensis	P	
Juncus ensifolius	P	
Juncus confusus	P	
Trifolium pratense	20	
Glyceria grandis	P	
Carex nebrascensis	T	
Deschampsia cespitosa	20	
Plantings (Populus tremuloides & Populus trichocarpa)	P	
Willow Sprigs	P	
Phalaris arundinacea	10	
Carex crawfordii	10	
Total Vegetative Cover:		90%

[illegible]

MDT WETLAND MONITORING – VEGETATION TRANSECT (back of form)

Cover Estimate

+	=	<1%	3	=	11-20%
1	=	1-5%	4	=	21-50%
2	=	6-10%	5	=	>50%

Indicator Class:

+ = Obligate
 - = Facultative/Wet
 0 = Facultative

Source:

P = Planted
 V = Volunteer

Percent of perimeter _____ % developing wetland vegetation – excluding dam/berm structures.

Establish transects perpendicular to the shoreline (or saturated perimeter). The transect should begin in the upland area. Permanently mark this location with a standard metal fencepost. Extend the imaginary transect line towards the center of the wetland, ending at the 3 foot depth (in open water), or at a point where water depths or saturation are maximized. Mark this location with another metal fencepost.

Estimate cover within a 10 ft wide “belt” along the transect length. At a minimum, establish a transect at the windward and leeward sides of the wetland. Remember that the purpose of this sampling is to monitor, not inventory, representative portions of the wetland site.

Notes:

BIRD SURVEY – FIELD DATA SHEET

Page__1_of_1__

Date: 8/01/05

Survey Time: 10:00-4:00

SITE: Camp Creek

[illegible]

Notes:

Behavior: BP – one of a breeding pair; BD – breeding display; F – foraging; FO – flyover; L – loafing; N – nesting

Habitat: AB – aquatic bed; FO – forested; I – island; MA – marsh; MF – mud flat; OW – open water; SS – scrub/shrub; UP – upland buffer; WM – wet meadow, US – unconsolidated shoreline

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Camp Creek</u> Applicant/Owner: <u>MDT</u> Investigator: <u>Greg Howard</u>	Date: <u>08/01/05</u> County: <u>Ravalli</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <u> x </u> Yes <u> </u> No Is the site significantly disturbed (Atypical Situation)? <u> </u> Yes <u> </u> No Is the area a potential Problem Area? <u> </u> Yes <u> </u> No (If needed, explain on reverse.)	Community ID: <u>Upland</u> Transect ID: <u>1</u> Plot ID: <u>1</u>

VEGETATION

	Dominant Plant Species	Stratum	Indicator		Dominant Plant Species	Stratum	Indicator
1	<i>Agropyron repens</i>	H	FAC-				
2	<i>Thlaspi arvensis</i>	H	--				
3	<i>Agrostis alba</i>	H	FAC+				
4	<i>Sisymbrium altissimum</i>	H	FACU-				
5	<i>Centaurea maculosa</i>	H	--				
6	<i>Potentilla fruticosa</i>	S	FAC-				

Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). 1/6 = 17%

Area dominated by upland vegetation.

HYDROLOGY

<u> </u> Recorded Data (Describe in Remarks): <u> </u> Stream, Lake, or Tide Gauge <u> </u> Aerial Photographs <u> </u> Other <u> X </u> No Recorded Data Available	Wetland Hydrology Indicators: Primary Indicators: <u> </u> Inundated <u> </u> Saturated in Upper 12 Inches <u> </u> Water Marks <u> </u> Drift Lines <u> </u> Sediment Deposits <u> </u> Drainage Patterns in Wetlands Secondary Indicators (2 or more required): <u> </u> Oxidized Root Channels in Upper 12 Inches <u> </u> Water-Stained Leaves <u> </u> Local Soil Survey Data <u> </u> FAC-Neutral Test <u> </u> Other (Explain in Remarks)
Field Observations: Depth of Surface Water: <u> </u> - <u> </u> (in.) Depth to Free Water in Pit: <u> </u> - <u> </u> (in.) Depth to Saturated Soil: <u> </u> - <u> </u> (in.)	
Remarks: No hydrology indicators present.	

SOILS

Map Unit Name (Series and Phase):		Gallatin-Shallow Muck Complex Gallatin		Drainage Class:	Imperfectly and Poorly-drained
Taxonomy (Subgroup):				Field Observations	
				Confirm Mapped Type?	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

Profile Description:					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 – 6+	A	10 YR 2/1	--	--	Loam with large cobbles

Hydric Soil Indicators:	
<input type="checkbox"/> Histosol	<input type="checkbox"/> Concretions
<input type="checkbox"/> Histic Epipedon	<input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils
<input type="checkbox"/> Sulfidic Odor	<input type="checkbox"/> Organic Streaking in Sandy Soils
<input type="checkbox"/> Aquic Moisture Regime	<input type="checkbox"/> Listed on Local Hydric Soils List
<input type="checkbox"/> Reducing Conditions	<input type="checkbox"/> Listed on National Hydric Soils List
<input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors	<input type="checkbox"/> Other (Explain in Remarks)

Soil pit located in area of created upland habitat, soils consisting of fill material excavated from channel reconstruction and removed from historic wetland.
--

WETLAND DETERMINATION

Hydrophytic Vegetation Present? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No Wetland Hydrology Present? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No Hydric Soils Present? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Is this Sampling Point Within a Wetland? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No
Remarks: Sampling point considered within an upland area.	

Approved by HQUSACE 2/92

DATA FORM

ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Camp Creek</u>				Date: <u>08/01/05</u>
Applicant/Owner: <u>MDT</u>				County: <u>Ravalli</u>
Investigator: <u>Greg Howard</u>				State: <u>MT</u>
Do Normal Circumstances exist on the site:	<u>X</u>	Yes	<u> </u>	No
Is the site significantly disturbed (Atypical Situation)?	<u> </u>	Yes	<u> </u>	No
Is the area a potential Problem Area?:	<u> </u>	Yes	<u> </u>	No
(If needed, explain on reverse.)				
Community ID:				<u>Emergent</u>
Transect ID:				<u>1</u>
Plot ID:				<u>2</u>

VEGETATION

VEGETATION			
	Dominant Plant Species	Stratum	Indicator
1	<i>Carex nebrascensis</i>	H	OBL
2	<i>Phalaris arundinacea</i>	H	FACW
3	<i>Geum macrophyllum</i>	H	OBL
4	<i>Agrostis alba</i>	H	FAC+
5	<i>Cirsium arvense</i>	H	FACU+
6	<i>Carex utriculata</i>	H	OBL
7	<i>Salix exigua</i>	S	OBL
Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). <u>6/7 = 85%</u>			
Area dominated by hydrophytic vegetation.			

HYDROLOGY

<p>Recorded Data (Describe in Remarks):</p> <p>_____ Stream, Lake, or Tide Gauge</p> <p>_____ Aerial Photographs</p> <p>_____ Other</p> <p><u> X </u> No Recorded Data Available</p>	<p>Wetland Hydrology Indicators:</p> <p>Primary Indicators:</p> <p>_____ Inundated</p> <p><u> X </u> Saturated in Upper 12 Inches</p> <p>_____ Water Marks</p> <p>_____ Drift Lines</p> <p>_____ Sediment Deposits</p> <p><u> X </u> Drainage Patterns in Wetlands</p> <p>Secondary Indicators (2 or more required):</p> <p>_____ Oxidized Root Channels in Upper 12 Inches</p> <p>_____ Water-Stained Leaves</p> <p>_____ Local Soil Survey Data</p> <p>_____ FAC-Neutral Test</p> <p>_____ Other (Explain in Remarks)</p>
<p>Field Observations:</p> <p>Depth of Surface Water: _____ (in.)</p> <p>Depth to Free Water in Pit: _____ (in.)</p> <p>Depth to Saturated Soil: <u> 8 </u> (in.)</p> <p>Remarks:</p> <p>Hydrology indicators present with saturated soils. Drainage patterns evident, area of lower topography. Historic side channel of Camp Creeks wet meadow complex.</p>	

SOILS

Map Unit Name		Gallatin-Shallow Muck Complex		Drainage Class: <u>Imperfectly and Poorly-drained</u>	
(Series and Phase):		<u>Gallatin</u>		Field Observations	
Taxonomy (Subgroup):				Confirm Mapped Type? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	

Profile Description:					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 – 3	O	10 YR 2/2	--	--	Roots & organics
3 – 6	A1	10 YR 2/1	--	--	Sandy loam & roots
6 – 8	A2	10 YR 2/1	--	--	Peat & sandy loam

Hydric Soil Indicators:	
<input type="checkbox"/> Histosol <input type="checkbox"/> Histic Epipedon <input type="checkbox"/> Sulfidic Odor <input type="checkbox"/> Aquic Moisture Regime <input type="checkbox"/> Reducing Conditions <input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors	<input type="checkbox"/> Concretions <input checked="" type="checkbox"/> High Organic Content in surface Layer in Sandy Soils <input type="checkbox"/> Organic Streaking in Sandy Soils <input type="checkbox"/> Listed on Local Hydric Soils List <input type="checkbox"/> Listed on National Hydric Soils List <input type="checkbox"/> Other (Explain in Remarks)

Hydric soils indicator present with low-chroma colors and high organic content (peat).

WETLAND DETERMINATION

Hydrophytic Vegetation Present?	<input checked="" type="checkbox"/>	Yes	<input type="checkbox"/>	No	Is this Sampling Point Within a Wetland? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
Wetland Hydrology Present?	<input checked="" type="checkbox"/>	Yes	<input type="checkbox"/>	No	
Hydric Soils Present?	<input checked="" type="checkbox"/>	Yes	<input type="checkbox"/>	No	

Remarks:
 Sampling point is considered within a wetland. Area of lower topography dominated by emergent type vegetation. Undisturbed wetlands mapped during initial delineation.

Approved by HQUSACE 2/92

ROUTINE WETLAND DETERMINATION (1987 COE Wetlands Delineation Manual)

Project/Site: <u>Camp Creek</u>				Date: <u>08/01/05</u>		
Applicant/Owner: <u>MDT</u>				County: <u>Ravalli</u>		
Investigator: <u>Greg Howard</u>				State: <u>MT</u>		
Do Normal Circumstances exist on the site:				X	Yes	No
Is the site significantly disturbed (Atypical Situation)?				<u> </u>	Yes	<u> </u> No
Is the area a potential Problem Area?:				<u> </u>	Yes	<u> </u> No
(If needed, explain on reverse.)						
Community ID:				<u>Emergent / Rock Bottom</u>		
Transect ID:				<u>1</u>		
Plot ID:				<u>3</u>		

VEGETATION

	Dominant Plant Species	Stratum	Indicator		Dominant Plant Species	Stratum	Indicator
1	<i>Carex utriculata</i>	H	OBL	7	<i>Agrostis alba</i>	H	FAC+
2	<i>Alopecurus pratensis</i>	H	FACW	8	<i>Centaurea maculosa</i>	H	--
3	<i>Carex nebrascensis</i>	H	OBL	9	<i>Veronica americana</i>	H	OBL
4	<i>Phalaris arundinacea</i>	H	FACW	10	<i>Juncus ensifolius</i>	H	FACW
5	<i>Epilobium ciliatum</i>	H	FACW	11	<i>Trifolium pratense</i>	H	FACU
6	<i>Deschampsia cespitosa</i>	H	FACW				
Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-).				9/11 = 81%			
Area consisting of streambank, creek and floodplain margins, dominated by hydrophytic vegetation.							

HYDROLOGY

_____ Recorded Data (Describe in Remarks): _____ Stream, Lake, or Tide Gauge _____ Aerial Photographs _____ Other <u> X </u> No Recorded Data Available		Wetland Hydrology Indicators: Primary Indicators: _____ Inundated <u> X </u> Saturated in Upper 12 Inches _____ Water Marks _____ Drift Lines _____ Sediment Deposits <u> X </u> Drainage Patterns in Wetlands Secondary Indicators (2 or more required): _____ Oxidized Root Channels in Upper 12 Inches _____ Water-Stained Leaves _____ Local Soil Survey Data _____ FAC-Neutral Test _____ Other (Explain in Remarks)
Field Observations: Depth of Surface Water: _____ (in.) Depth to Free Water in Pit: _____ (in.) Depth to Saturated Soil: <u> 0 </u> (in.)		
Remarks: Hydrology indicator present with soils saturated to ground surface and drainage patterns.		

SOILS

Map Unit Name		Gallatin-Shallow Muck Complex		Drainage Class: <u>Imperfectly and Poorly-drained</u>	
(Series and Phase):		<u>Gallatin</u>		Field Observations	
Taxonomy (Subgroup):				Confirm Mapped Type? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	

Profile Description:					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 – 8+	B	10 YR 2/1	--	--	Loam with large cobbles

Hydric Soil Indicators:	
<input type="checkbox"/> Histosol <input type="checkbox"/> Histic Epipedon <input type="checkbox"/> Sulfidic Odor <input type="checkbox"/> Aquic Moisture Regime <input type="checkbox"/> Reducing Conditions <input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors	<input type="checkbox"/> Concretions <input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils <input type="checkbox"/> Organic Streaking in Sandy Soils <input type="checkbox"/> Listed on Local Hydric Soils List <input type="checkbox"/> Listed on National Hydric Soils List <input type="checkbox"/> Other (Explain in Remarks)

Hydric soil indicator present with low-chroma colors.

WETLAND DETERMINATION

Hydrophytic Vegetation Present?	<input checked="" type="checkbox"/>	Yes	<input type="checkbox"/>	No	Is this Sampling Point Within a Wetland? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
Wetland Hydrology Present?	<input checked="" type="checkbox"/>	Yes	<input type="checkbox"/>	No	
Hydric Soils Present?	<input checked="" type="checkbox"/>	Yes	<input type="checkbox"/>	No	

Remarks:
 Sampling point considered within a wetland and Waters of the U.S. Floodplain along Camp Creek developing into emergent and scrub-shrub wetland vegetation types.

Approved by HQUSACE 2/92

ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Camp Creek</u> Applicant/Owner: <u>MDT</u> Investigator: <u>Greg Howard</u>	Date: <u>08/01/05</u> County: <u>Ravalli</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <u>X</u> Yes <u> </u> No Is the site significantly disturbed (Atypical Situation)? <u> </u> Yes <u> </u> No Is the area a potential Problem Area?: <u> </u> Yes <u> </u> No (If needed, explain on reverse.)	Community ID: <u>Emergent</u> Transect ID: <u>1</u> Plot ID: <u>4</u>

VEGETATION

	Dominant Plant Species	Stratum	Indicator		Dominant Plant Species	Stratum	Indicator
1	<i>Phalaris arundinacea</i>	H	FACW				
2	<i>Agrostis alba</i>	H	FAC+				
3	<i>Carex lanuginosa</i>	H	OBL				
4	<i>Carex nebrascensis</i>	H	OBL				
5	<i>Deschampsia cespitosa</i>	H	FACW				
Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). <u>5/5 = 100%</u>							
Area dominated by hydrophytic vegetation.							

HYDROLOGY

<p>_____ Recorded Data (Describe in Remarks):</p> <p style="margin-left: 40px;">_____ Stream, Lake, or Tide Gauge</p> <p style="margin-left: 40px;">_____ Aerial Photographs</p> <p style="margin-left: 40px;">_____ Other</p> <p><u>X</u> No Recorded Data Available</p> <hr/> <p>Field Observations:</p> <p style="margin-left: 40px;">Depth of Surface Water: _____ (in.)</p> <p style="margin-left: 40px;">Depth to Free Water in Pit: _____ (in.)</p> <p style="margin-left: 40px;">Depth to Saturated Soil: <u>8</u> (in.)</p>	<p>Wetland Hydrology Indicators:</p> <p>Primary Indicators:</p> <p style="margin-left: 20px;"><u>X</u> Inundated</p> <p style="margin-left: 20px;"><u> </u> Saturated in Upper 12 Inches</p> <p style="margin-left: 20px;"><u> </u> Water Marks</p> <p style="margin-left: 20px;"><u> </u> Drift Lines</p> <p style="margin-left: 20px;"><u> </u> Sediment Deposits</p> <p style="margin-left: 20px;"><u>X</u> Drainage Patterns in Wetlands</p> <p>Secondary Indicators (2 or more required):</p> <p style="margin-left: 20px;"><u> </u> Oxidized Root Channels in Upper 12 Inches</p> <p style="margin-left: 20px;"><u> </u> Water-Stained Leaves</p> <p style="margin-left: 20px;"><u> </u> Local Soil Survey Data</p> <p style="margin-left: 20px;"><u> </u> FAC-Neutral Test</p> <p style="margin-left: 20px;"><u> </u> Other (Explain in Remarks)</p>
Remarks: Hydrology indicators present with saturated soils and drainage patterns.	

SOILS

Map Unit Name (Series and Phase):		Gallatin-Shallow Muck Complex Gallatin		Drainage Class: Imperfectly and Poorly-drained	
Taxonomy (Subgroup):				Field Observations Confirm Mapped Type? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	

Profile Description:					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0 – 3	O	10 YR 2/1	--	--	Roots & organics
3 – 5	A	10 YR 2/1	--	--	Sandy loam & roots
5 – 7	B	--	--	--	Sand with fine gravels
7 – 10+	A	10 YR 2/1	--	--	Sandy loam with fine to medium gravels

Hydric Soil Indicators:	
<input type="checkbox"/> Histosol <input type="checkbox"/> Histic Epipedon <input type="checkbox"/> Sulfidic Odor <input type="checkbox"/> Aquic Moisture Regime <input type="checkbox"/> Reducing Conditions <input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors	<input type="checkbox"/> Concretions <input checked="" type="checkbox"/> High Organic Content in surface Layer in Sandy Soils <input type="checkbox"/> Organic Streaking in Sandy Soils <input type="checkbox"/> Listed on Local Hydric Soils List <input type="checkbox"/> Listed on National Hydric Soils List <input type="checkbox"/> Other (Explain in Remarks)

Hydric soil indicators present with low-chroma colors and high organic content in sandy soils.
--

WETLAND DETERMINATION

Hydrophytic Vegetation Present? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No Wetland Hydrology Present? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No Hydric Soils Present? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Is this Sampling Point Within a Wetland? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
--	--

Remarks: Sampling point is considered within an emergent wetland type. Located on upper terrace adjacent to created floodplain. Remnant wetlands not disturbed during construction efforts.
--

Approved by HQUSACE 2/92

1. Project Name: Camp Creek

2. Project #: B43054.00-0106 Control #:

3. Evaluation Date: 8/1/2005

4. Evaluator(s): Greg Howard

5. Wetland / Site #(s): AA-1, MDT Property

6. Wetland Location(s) i. T: 1 N R: 19 W S: 27 & 34 T: N R: E S:

ii. Approx. Stationing / Mileposts:

iii. Watershed: 3 - Lower Clark Fork GPS Reference No. (if applies):

Other Location Information: Located in Sula Basin; Reconstructed Camp Creek channel, floodplain, and adjacent emergent wetlands.

7. A. Evaluating Agency MDT

8. Wetland Size (total acres): (visually estimated)
36.08 (measured, e.g. GPS)

B. Purpose of Evaluation:

☐ Wetlands potentially affected by MDT project

☐ Mitigation wetlands; pre-construction

☒ Mitigation wetlands; post-construction

☐ Other

9. Assessment Area (total acres): (visually estimated)
36.08 (measured, e.g. GPS)

Comments: AA includes 0.95 acres of stream channel.

HGM CLASS ¹	SYSTEM ²	SUBSYSTEM ²	CLASS ²	WATER REGIME ²	MODIFIER ²	% OF AA
Riverine	Riverine	Upper Perennial	Rock Bottom	Permanently Flooded	---	3
Riverine	Palustrine	---	Emergent Wetland	Seasonally Flooded	---	82
Riverine	Palustrine	---	Scrub-Shrub Wetland	Seasonally Flooded	---	15
---	---	---	---	---	---	

¹ = Smith et al. 1995. ² = Cowardin et al. 1979.

Comments: _____

11. ESTIMATED RELATIVE ABUNDANCE (of similarly classified sites within the same Major Montana Watershed Basin)

Common Comments:

12. GENERAL CONDITION OF AA

i. Regarding Disturbance: (Use matrix below to select appropriate response.)

Conditions Within AA	Predominant Conditions Adjacent (within 500 Feet) To AA		
	Land managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings.	Land not cultivated, but moderately grazed or hayed or selectively logged or has been subject to minor clearing; contains few roads or buildings.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density.
AA occurs and is managed in predominantly a natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings.	---	low disturbance	---
AA not cultivated, but moderately grazed or hayed or selectively logged or has been subject to relatively minor clearing, or fill placement, or hydrological alteration; contains few roads or buildings.	---	---	---
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density.	---	---	---

Comments: (types of disturbance, intensity, season, etc.) Past disturbances include grazing, clearing, and hydrologic alterations; currently low disturbance.

ii. **Prominent weedy, alien, & introduced species:** Spotted knapweed, bull thistle, Canada thistle, hound' tongue, pennycress, common dandelion, tumble mustard.

iii. **Briefly describe AA and surrounding land use / habitat:** AA located in Sula Basin, Camp Creek, and adjacent wetlands. USFS land & private ownership. Surround land use includes pasture, livestock grazing, and logging.

13. STRUCTURAL DIVERSITY (Based on 'Class' column of #10 above.)

Number of 'Cowardin' Vegetated Classes Present in AA	≥3 Vegetated Classes or ≥ 2 if one class is forested	2 Vegetated Classes or 1 if forested	≤ 1 Vegetated Class
Select Rating	---	Moderate	---

Comments:

14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS

i. AA is Documented (D) or Suspected (S) to contain (check box):

Primary or Critical habitat (list species) ☐ D ☐ S _____
 Secondary habitat (list species) ☒ D ☐ S bull trout, bald eagle
 Incidental habitat (list species) ☐ D ☐ S _____
 No usable habitat ☐ D ☐ S _____

ii. Rating (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point & Rating	---	---	.8 (M)	---	---	---	---

If documented, list the source (e.g., observations, records, etc.): FWP records and observations.

14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.

Do not include species listed in 14A(i).

i. AA is Documented (D) or Suspected (S) to contain (check box):

Primary or Critical habitat (list species) ☐ D ☒ S west-slope cutthroat trout
 Secondary habitat (list species) ☐ D ☐ S _____
 Incidental habitat (list species) ☐ D ☐ S _____
 No usable habitat ☐ D ☐ S _____

ii. Rating: Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point & Rating	---	.8 (H)	---	---	---	---	---

If documented, list the source (e.g., observations, records, etc.): FWP records.

14C. GENERAL WILDLIFE HABITAT RATING

i. Evidence of overall wildlife use in the AA: Check either substantial, moderate, or low.

☐ Substantial (based on any of the following)

- ☐ observations of abundant wildlife #s or high species diversity (during any period)
- ☐ abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ presence of extremely limiting habitat features not available in the surrounding area
- ☐ interviews with local biologists with knowledge of the AA

☐ Low (based on any of the following)

- ☐ few or no wildlife observations during peak use periods
- ☐ little to no wildlife sign
- ☐ sparse adjacent upland food sources
- ☐ interviews with local biologists with knowledge of AA

☒ Moderate (based on any of the following)

- ☒ observations of scattered wildlife groups or individuals or relatively few species during peak periods
- ☒ common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☒ adequate adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

ii. Wildlife Habitat Features: Working from top to bottom, select the AA attribute to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from 13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see 10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A = absent.

Structural Diversity (from 13)	<input type="checkbox"/> High								<input checked="" type="checkbox"/> Moderate								<input type="checkbox"/> Low			
	<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even				<input checked="" type="checkbox"/> Uneven				<input type="checkbox"/> Even			
Class Cover Distribution (all vegetated classes)	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Duration of Surface Water in ≥ 10% of AA																				
Low disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	H	--	--	--	--	--	--
Moderate disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
High disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

iii. Rating: Use 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.

Evidence of Wildlife Use from 14C(i)	Wildlife Habitat Features Rating from 14C(ii)			
	<input type="checkbox"/> Exceptional	<input checked="" type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
Substantial	--	--	--	--
Moderate	--	.7 (M)	--	--
Low	--	--	--	--

Comments: _____

14D. GENERAL FISH / AQUATIC HABITAT RATING ☐ NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat or excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

i. Habitat Quality: Pick the appropriate AA attributes in matrix to determine the quality rating of exceptional (E), high (H), moderate (M), or low (L).

Duration of Surface Water in AA	<input checked="" type="checkbox"/> Permanent/Perennial			<input type="checkbox"/> Seasonal / Intermittent			<input type="checkbox"/> Temporary / Ephemeral		
	>25%	10-25%	<10%	>25%	10-25%	<10%	>25%	10-25%	<10%
Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation)									
Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities	--	--	--	--	--	--	--	--	--
Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--
Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	M	--	--	--	--	--	--	--

ii. Modified Habitat Quality: Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support?

☐ Y ☒ N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating: ☐ E ☐ H ☐ M ☐ L

iii. Rating: Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).

Types of Fish Known or Suspected within AA	Modified Habitat Quality from 14D(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> Low
Native game fish	--	--	.7 (M)	--
Introduced game fish	--	--	--	--
Non-game fish	--	--	--	--
No fish	--	--	--	--

Comments: Reconstructed channel supports native fish populations. Enhancement of habitat: pools, riffles, and overhanging banks. Ratings will improve with establishment of woody vegetation.

14E. FLOOD ATTENUATION ☐ NA (proceed to 14G)

Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA do not flood from in-channel or overbank flow, then check NA.

i. Rating: Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Estimated wetland area in AA subject to periodic flooding	<input checked="" type="checkbox"/> ≥ 10 acres			<input type="checkbox"/> <10, >2 acres			<input type="checkbox"/> ≤2 acres		
	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
% of flooded wetland classified as forested, scrub/shrub, or both									
AA contains no outlet or restricted outlet	--	--	.6 (M)	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	--	--	--	--

ii. Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA? (check)

☒ Y ☐ N **Comments:** USFS offices downstream, adjacent parcel with MDT boundary.

14F. SHORT AND LONG TERM SURFACE WATER STORAGE ☐ NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow.

If no wetlands in the AA are subject to flooding or ponding, then check NA above.

i. Rating: Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.
P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding.	<input checked="" type="checkbox"/> >5 acre feet			<input type="checkbox"/> <5, >1 acre feet			<input type="checkbox"/> ≤1 acre foot		
	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Duration of surface water at wetlands within the AA									
Wetlands in AA flood or pond ≥ 5 out of 10 years	1 (H)	--	--	--	--	--	--	--	--
Wetlands in AA flood or pond < 5 out of 10 years	--	--	--	--	--	--	--	--	--

Comments: _____

14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL ☐ NA (proceed to 14H)

Applies to wetlands with the potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input.

If no wetlands in the AA are subject to such input, check NA above.

i. Rating: Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Sediment, Nutrient, and Toxicant Input Levels Within AA	AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
	<input type="checkbox"/> ≥ 70%		<input checked="" type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
% cover of wetland vegetation in AA	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Evidence of flooding or ponding in AA								
AA contains no or restricted outlet	--	--	.7 (M)	--	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	--	--	--

Comments: Minor sedimentation due to logging and recent forest fires.

14H. SEDIMENT/ShORELINE STABILIZATION☐ NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, then check NA above.

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

% Cover of wetland streambank or shoreline by species with deep, binding rootmasses.	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input checked="" type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
≥ 65 %	--	--	--
35-64 %	.7 (M)	--	--
< 35 %	--	--	--

Comments: Increased density of willows and wetland grasses / grass-like plants along streambanks.

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet. P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

A	<input checked="" type="checkbox"/> Vegetated component >5 acres						<input type="checkbox"/> Vegetated component 1-5 acres						<input type="checkbox"/> Vegetated component <1 acre					
B	<input type="checkbox"/> High		<input checked="" type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low	
C	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input checked="" type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N
P/P	--	--	.9H	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
S/I	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
T/E/A	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Comments: _____

14J. GROUNDWATER DISCHARGE / RECHARGE (DR) (Check the indicators in i & ii below that apply to the AA.)i. ☒ Discharge Indicators

- ☐ Springs are known or observed.
☐ Vegetation growing during dormant season / drought.
☒ Wetland occurs at the toe of a natural slope.
☐ Seeps are present at the wetland edge.
☐ AA permanently flooded during drought periods.
☐ Wetland contains an outlet, but no inlet.
☐ Other _____

ii. ☒ Recharge Indicators

- ☒ Permeable substrate presents without underlying impeding layer.
☐ Wetland contains inlet but not outlet.
☐ Other _____

iii. **Rating:** Use information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

Criteria	Functional Point and Rating
AA has known Discharge/Recharge area or one or more indicators of D/R present	1 (H)
No Discharge/Recharge indicators present	--
Available Discharge/Recharge information inadequate to rate AA D/R potential	--

Comments: Channel & floodplain located in Sula Basin. Steep slopes on both sides of basin. Wetland occurring along toe of slope.

14K. UNIQUENESS

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP.			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP.			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate.		
Estimated Relative Abundance from 11	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input checked="" type="checkbox"/> common	<input type="checkbox"/> abundant
Low disturbance at AA (12i)	--	--	--	--	--	--	--	.4M	--
Moderate disturbance at AA (12i)	--	--	--	--	--	--	--	--	--
High disturbance at AA (12i)	--	--	--	--	--	--	--	--	--

Comments: _____

14L. RECREATION / EDUCATION POTENTIAL

i. Is the AA a known recreational or educational site? ☐ Yes [Rate ☐ High (1.0), then proceed to 14L(ii) only] ☒ No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA: ☒ Educational / scientific study ☒ Consumptive rec. ☒ Non-consumptive rec. ☐ Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

☒ Yes [Proceed to 14L (ii) and then 14L(iv)] ☐ No [Rate as low in 14L(iv)]

iv. **Rating** Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Ownership	Disturbance at AA from 12(i)		
	<input checked="" type="checkbox"/> Low	<input type="checkbox"/> Moderate	<input type="checkbox"/> High
Public ownership	1(H)	--	--
Private ownership	--	--	--

Comments: Good potential for recreation/education as located along Highway 93.

FUNCTION, VALUE SUMMARY, AND OVERALL RATING

Function and Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	moderate	0.80	1	
B. MT Natural Heritage Program Species Habitat	high	0.80	1	
C. General Wildlife Habitat	moderate	0.70	1	
D. General Fish/Aquatic Habitat	moderate	0.70	1	
E. Flood Attenuation	moderate	0.60	1	
F. Short and Long Term Surface Water Storage	high	1.00	1	
G. Sediment/Nutrient/Toxicant Removal	moderate	0.70	1	
H. Sediment/Shoreline Stabilization	moderate	0.70	1	
I. Production Export/Food Chain Support	high	0.90	1	
J. Groundwater Discharge/Recharge	high	1.00	1	
K. Uniqueness	moderate	0.40	1	
L. Recreation/Education Potential	moderate	1.00	1	
Total:		<u>9.30</u>	<u>12.00</u>	
Percent of Total Possible Points:		<u>78%</u> (Actual / Possible) x 100 [rd to nearest whole #]		

Category I Wetland: (Must satisfy one of the following criteria. If not satisfied, proceed to Category II.) <input type="checkbox"/> Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; or <input type="checkbox"/> Score of 1 functional point for Uniqueness; or <input type="checkbox"/> Score of 1 functional point for Flood Attenuation and answer to Question 14E(ii) is "yes"; or <input type="checkbox"/> Percent of total Possible Points is > 80%.
Category II Wetland: (Criteria for Category I not satisfied and meets any one of the following Category II criteria. If not satisfied, proceed to Category IV.) <input type="checkbox"/> Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; or <input type="checkbox"/> Score of .9 or 1 functional point for General Wildlife Habitat; or <input type="checkbox"/> Score of .9 or 1 functional point for General Fish/Aquatic Habitat; or <input type="checkbox"/> "High" to "Exceptional" ratings for both General Wildlife Habitat and General Fish / Aquatic Habitat; or <input type="checkbox"/> Score of .9 functional point for Uniqueness; or <input checked="" type="checkbox"/> Percent of total possible points is > 65%.
<input type="checkbox"/> Category III Wetland: (Criteria for Categories I, II, or IV not satisfied.)
Category IV Wetland: (Criteria for Categories I or II are not satisfied and <u>all</u> of the following criteria are met; If not satisfied, return to Category III.) <input type="checkbox"/> "Low" rating for Uniqueness; and <input type="checkbox"/> "Low" rating for Production Export / Food Chain Support; and <input type="checkbox"/> Percent of total possible points is < 30%.

OVERALL ANALYSIS AREA (AA) RATING: (Check appropriate category based on the criteria outlined above.)

☐ **I**
☒ **II**
☐ **III**
☐ **IV**

1. Project Name: Camp Creek

2. Project #: B43054.00-00106

Control #: _____

3. Evaluation Date: 8/1/2005

4. Evaluator(s): Greg Howard

5. Wetland / Site #(s): AA-2; Grasser Property

6. Wetland Location(s) i. T: 1 N R: 19 W S: 22, 27, & 34 T: __ N R: __ E S: _____

ii. Approx. Stationing / Mileposts: _____

iii. Watershed: 3 - Lower Clark Fork GPS Reference No. (if applies): _____

Other Location Information: Located in Sula Basin; Reconstructed Camp Creek channel, floodplain, and adjacent emergent wetlands.

7. A. Evaluating Agency MDT

8. Wetland Size (total acres): _____ (visually estimated)
8.13 (measured, e.g. GPS)

B. Purpose of Evaluation:

☐ Wetlands potentially affected by MDT project

☐ Mitigation wetlands; pre-construction

☒ Mitigation wetlands; post-construction

☐ Other

9. Assessment Area (total acres): _____ (visually estimated)
8.13 (measured, e.g. GPS)

Comments: AA includes 1.2 acres of stream channel.

HGM CLASS ¹	SYSTEM ²	SUBSYSTEM ²	CLASS ²	WATER REGIME ²	MODIFIER ²	% OF AA
Riverine	Palustrine	None	Emergent Wetland	Seasonally Flooded	---	80
Riverine	Riverine	Upper Perennial	Rock Bottom	Permanently Flooded	---	15
Riverine	Palustrine	None	Scrub-Shrub Wetland	Seasonally Flooded	---	5
---	---	---	---	---	---	

¹ = Smith et al. 1995. ² = Cowardin et al. 1979.

Comments: _____

11. ESTIMATED RELATIVE ABUNDANCE (of similarly classified sites within the same Major Montana Watershed Basin)

Abundant **Comments:**

12. GENERAL CONDITION OF AA

i. Regarding Disturbance: (Use matrix below to select appropriate response.)

Conditions Within AA	Predominant Conditions Adjacent (within 500 Feet) To AA		
	Land managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings.	Land not cultivated, but moderately grazed or hayed or selectively logged or has been subject to minor clearing; contains few roads or buildings.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading clearing, or hydrological alteration; high road or building density.
AA occurs and is managed in predominantly a natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings.	---	---	---
AA not cultivated, but moderately grazed or hayed or selectively logged or has been subject to relatively minor clearing, or fill placement, or hydrological alteration; contains few roads or buildings.	---	moderate disturbance	---
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density.	---	---	---

Comments: (types of disturbance, intensity, season, etc.) Historically and currently grazed.

ii. **Prominent weedy, alien, & introduced species:** Spotted knapweed, Canada thistle, hound' tongue, pennycress, common dandelion, and tumble mustard.

iii. Briefly describe AA and surrounding land use / habitat: Camp Creek and adjacent wetland within the Sula Basin. Land uses include pasture, logging, private, and USFS.

13. STRUCTURAL DIVERSITY (Based on 'Class' column of #10 above.)

Number of 'Cowardin' Vegetated Classes Present in AA	≥3 Vegetated Classes or ≥ 2 if one class is forested	2 Vegetated Classes or 1 if forested	≤ 1 Vegetated Class
Select Rating	---	Moderate	---

Comments:

14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS

i. AA is Documented (D) or Suspected (S) to contain (check box):

Primary or Critical habitat (list species) ☐ D ☐ S _____
 Secondary habitat (list species) ☒ D ☐ S bull trout, bald eagle
 Incidental habitat (list species) ☐ D ☐ S _____
 No usable habitat ☐ D ☐ S _____

ii. Rating (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point & Rating	---	---	.8 (M)	---	---	---	---

If documented, list the source (e.g., observations, records, etc.): FWP records and observations on MDT site.

14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.

Do not include species listed in 14A(i).

i. AA is Documented (D) or Suspected (S) to contain (check box):

Primary or Critical habitat (list species) ☐ D ☒ S west-slope cutthroat trout
 Secondary habitat (list species) ☐ D ☐ S _____
 Incidental habitat (list species) ☐ D ☐ S _____
 No usable habitat ☐ D ☐ S _____

ii. Rating: Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point & Rating	---	.8 (H)	---	---	---	---	---

If documented, list the source (e.g., observations, records, etc.): FWP records.

14C. GENERAL WILDLIFE HABITAT RATING

i. Evidence of overall wildlife use in the AA: Check either substantial, moderate, or low.

☐ Substantial (based on any of the following)

- ☐ observations of abundant wildlife #s or high species diversity (during any period)
- ☐ abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ presence of extremely limiting habitat features not available in the surrounding area
- ☐ interviews with local biologists with knowledge of the AA

☐ Low (based on any of the following)

- ☐ few or no wildlife observations during peak use periods
- ☐ little to no wildlife sign
- ☐ sparse adjacent upland food sources
- ☐ interviews with local biologists with knowledge of AA

☒ Moderate (based on any of the following)

- ☒ observations of scattered wildlife groups or individuals or relatively few species during peak periods
- ☒ common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☒ adequate adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

ii. Wildlife Habitat Features: Working from top to bottom, select the AA attribute to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from 13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see 10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A = absent.

Structural Diversity (from 13)	<input type="checkbox"/> High								<input checked="" type="checkbox"/> Moderate								<input type="checkbox"/> Low			
Class Cover Distribution (all vegetated classes)	<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even				<input checked="" type="checkbox"/> Uneven				<input type="checkbox"/> Even			
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Low disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	M	--	--	--	--	--	--
High disturbance at AA (see 12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

iii. Rating: Use 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.

Evidence of Wildlife Use from 14C(i)	Wildlife Habitat Features Rating from 14C(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> Low
Substantial	--	--	--	--
Moderate	--	--	.5 (M)	--
Low	--	--	--	--

Comments: _____

14D. GENERAL FISH / AQUATIC HABITAT RATING ☐ NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat or excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

i. Habitat Quality: Pick the appropriate AA attributes in matrix to determine the quality rating of exceptional (E), high (H), moderate (M), or low (L).

Duration of Surface Water in AA	<input checked="" type="checkbox"/> Permanent/Perennial			<input type="checkbox"/> Seasonal / Intermittent			<input type="checkbox"/> Temporary / Ephemeral		
	>25%	10-25%	<10%	>25%	10-25%	<10%	>25%	10-25%	<10%
Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation)									
Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities	--	--	--	--	--	--	--	--	--
Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--
Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	M	--	--	--	--	--	--

ii. Modified Habitat Quality: Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support?

☐ Y ☒ N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating: ☐ E ☐ H ☐ M ☐ L

iii. Rating: Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).

Types of Fish Known or Suspected within AA	Modified Habitat Quality from 14D(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> Low
Native game fish	--	--	.7 (M)	--
Introduced game fish	--	--	--	--
Non-game fish	--	--	--	--
No fish	--	--	--	--

Comments: Reconstructed channel supports native fish populations. Enhancement of habitat: pools, riffles, and overhanging banks. Ratings will improve with establishment of woody vegetation.

14E. FLOOD ATTENUATION ☐ NA (proceed to 14G)

Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA do not flood from in-channel or overbank flow, then check NA.

i. Rating: Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Estimated wetland area in AA subject to periodic flooding	<input type="checkbox"/> ≥ 10 acres			<input checked="" type="checkbox"/> <10, >2 acres			<input type="checkbox"/> ≤2 acres		
	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
% of flooded wetland classified as forested, scrub/shrub, or both									
AA contains no outlet or restricted outlet	--	--	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	.4 (M)	--	--	--

ii. Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA? (check)

☒ Y ☐ N **Comments:** USFS offices downstream & several other homes located nearby.

14F. SHORT AND LONG TERM SURFACE WATER STORAGE ☐ NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow.

If no wetlands in the AA are subject to flooding or ponding, then check NA above.

i. Rating: Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.
P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding.	<input type="checkbox"/> >5 acre feet			<input checked="" type="checkbox"/> <5, >1 acre feet			<input type="checkbox"/> ≤1 acre foot		
	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Duration of surface water at wetlands within the AA									
Wetlands in AA flood or pond ≥ 5 out of 10 years	--	--	--	--	.6 (M)	--	--	--	--
Wetlands in AA flood or pond < 5 out of 10 years	--	--	--	--	--	--	--	--	--

Comments: _____

14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL ☐ NA (proceed to 14H)

Applies to wetlands with the potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input.

If no wetlands in the AA are subject to such input, check NA above.

i. Rating: Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Sediment, Nutrient, and Toxicant Input Levels Within AA	AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
	<input type="checkbox"/> ≥ 70%		<input checked="" type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
% cover of wetland vegetation in AA	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
Evidence of flooding or ponding in AA								
AA contains no or restricted outlet	--	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	.6 (M)	--	--	--	--	--

Comments: Minor sediment from nearby burned forest. Potential nutrient input due to heavy livestock grazing in Sula Basin.

14H. SEDIMENT/ShORELINE STABILIZATION☐ NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, then check NA above.

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

% Cover of wetland streambank or shoreline by species with deep, binding rootmasses.	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input checked="" type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
≥ 65 %	--	--	--
35-64 %	--	--	--
< 35 %	.3 (L)	--	--

Comments: _____

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet. P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

A	<input checked="" type="checkbox"/> Vegetated component >5 acres						<input type="checkbox"/> Vegetated component 1-5 acres						<input type="checkbox"/> Vegetated component <1 acre					
B	<input type="checkbox"/> High		<input checked="" type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low	
C	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input checked="" type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N
P/P	--	--	.9H	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
S/I	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
T/E/A	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Comments: _____

14J. GROUNDWATER DISCHARGE / RECHARGE (DR) (Check the indicators in i & ii below that apply to the AA.)i. ☒ Discharge Indicators

- ☐ Springs are known or observed.
☐ Vegetation growing during dormant season / drought.
☒ Wetland occurs at the toe of a natural slope.
☐ Seeps are present at the wetland edge.
☐ AA permanently flooded during drought periods.
☐ Wetland contains an outlet, but no inlet.
☐ Other _____

ii. ☒ Recharge Indicators

- ☒ Permeable substrate presents without underlying impeding layer.
☐ Wetland contains inlet but not outlet.
☐ Other _____

iii. **Rating:** Use information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

Criteria	Functional Point and Rating
AA has known Discharge/Recharge area or one or more indicators of D/R present	1 (H)
No Discharge/Recharge indicators present	--
Available Discharge/Recharge information inadequate to rate AA D/R potential	--

Comments: _____

14K. UNIQUENESS

i. **Rating:** Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP.			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP.			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate.		
Estimated Relative Abundance from 11	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input checked="" type="checkbox"/> abundant
Low disturbance at AA (12i)	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (12i)	--	--	--	--	--	--	--	--	.2L
High disturbance at AA (12i)	--	--	--	--	--	--	--	--	--

Comments: _____

14L. RECREATION / EDUCATION POTENTIAL

i. Is the AA a known recreational or educational site? ☐ Yes [Rate ☐ High (1.0), then proceed to 14L(ii) only] ☒ No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA: ☒ Educational / scientific study ☒ Consumptive rec. ☒ Non-consumptive rec. ☐ Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

☒ Yes [Proceed to 14L (ii) and then 14L(iv)] ☐ No [Rate as low in 14L(iv)]

iv. **Rating** Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Ownership	Disturbance at AA from 12(i)		
	<input type="checkbox"/> Low	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> High
Public ownership	--	--	--
Private ownership	--	.3(L)	--

Comments: Good potential for recreation/education as it is adjacent to Highway 93, though under private ownership.

FUNCTION, VALUE SUMMARY, AND OVERALL RATING

Function and Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	moderate	0.80	1	
B. MT Natural Heritage Program Species Habitat	high	0.80	1	
C. General Wildlife Habitat	moderate	0.50	1	
D. General Fish/Aquatic Habitat	moderate	0.70	1	
E. Flood Attenuation	moderate	0.40	1	
F. Short and Long Term Surface Water Storage	moderate	0.60	1	
G. Sediment/Nutrient/Toxicant Removal	moderate	0.60	1	
H. Sediment/Shoreline Stabilization	low	0.30	1	
I. Production Export/Food Chain Support	high	0.90	1	
J. Groundwater Discharge/Recharge	high	1.00	1	
K. Uniqueness	low	0.20	1	
L. Recreation/Education Potential	low	0.30	1	
Total:		<u>7.10</u>	<u>12.00</u>	
Percent of Total Possible Points:		<u>59%</u> (Actual / Possible) x 100 [rd to nearest whole #]		

Category I Wetland: (Must satisfy **one** of the following criteria. If not satisfied, proceed to Category II.)

- ☐ Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**
- ☐ Score of 1 functional point for Uniqueness; **or**
- ☐ Score of 1 functional point for Flood Attenuation **and** answer to Question 14E(ii) is "yes"; **or**
- ☐ Percent of total Possible Points is > 80%.

Category II Wetland: (Criteria for Category I not satisfied **and** meets any **one** of the following Category II criteria. If not satisfied, proceed to Category IV.)

- ☐ Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; **or**
- ☐ Score of .9 or 1 functional point for General Wildlife Habitat; **or**
- ☐ Score of .9 or 1 functional point for General Fish/Aquatic Habitat; **or**
- ☐ "High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish / Aquatic Habitat; **or**
- ☐ Score of .9 functional point for Uniqueness; **or**
- ☐ Percent of total possible points is > 65%.

☒ **Category III Wetland:** (Criteria for Categories I, II, or IV not satisfied.)

Category IV Wetland: (Criteria for Categories I or II are not satisfied **and** all of the following criteria are met; If not satisfied, return to Category III.)

- ☐ "Low" rating for Uniqueness; **and**
- ☐ "Low" rating for Production Export / Food Chain Support; **and**
- ☐ Percent of total possible points is < 30%.

OVERALL ANALYSIS AREA (AA) RATING: (Check appropriate category based on the criteria outlined above.)

☐ **I**
☐ **II**
☒ **III**
☐ **IV**

Appendix C

REPRESENTATIVE PHOTOGRAPHS

MDT Wetland Mitigation Monitoring
Camp Creek
Sula, Montana

CAMP CREEK MITIGATION SITE 2005



Photo Point No. 1: View looking northeast along vegetation transect, end point in foreground.



Photo Point No. 2: View looking southwest along vegetation transect, starting point in foreground, located in upland community type.



Photo Point No. 3: View looking northeast, constructed Camp Creek channel and floodplain margins. Area becoming dominated by wetland species.



Photo Point No. 4: View looking north, floodplain margins with emergent wetland and riparian vegetation enhancements. Large containerized cottonwood and aspen plantings.



Photo Point No. 7: View looking south; lowest section of Camp Creek channel, north boundary of MDT parcel.




Photo Point No. 8: View looking west across mitigation site, upland community type in foreground. Area dominated by mostly invasive species.

CAMP CREEK MITIGATION SITE 2005



Photo Point No. 9: View looking north, main channel just below second culvert. Example of fabric work along constructed streambanks.



Photo Point No. 10: View looking south, section of channel with shrub communities present.



Photo Point No. 12: View looking south, main channel running along Grasser structures, shrub community present.



Photo Point No. 13: View looking south, straight sections of main channel running across upper portion of Grasser parcel.

CAMP CREEK MITIGATION SITE 2005



Photo Point No. 5: Panoramic looking west across site. Representative photo of typical channel and floodplain section present at Camp Creek with MDT owned parcels. Floodplain areas dominated by mostly wetland species. Areas of floodplain with saturated soils during late summer visit. The shrub and tree plantings showing new vigorous growth.



Photo Point No. 11: View looking north along creek, below upper road crossing and culvert near Grasser complex. Mature cottonwoods and remnant shrub communities present along creek. Floodplain areas with spotted knapweed infestations.

Appendix D

ORIGINAL SITE PLAN

*MDT Wetland Mitigation Monitoring
Camp Creek
Sula, Montana*

STATE	PROJECT NUMBER	SHEET
MONTANA	NH 41 (24)	29

CAMP CREEK RESTORATION

LEGEND

(B2) BANK PROTECTION
TREATMENT TYPE 2
SEE DETAILS

PR POOL RIGHT

PL POOL LEFT

R RIFFLE - 10

EACH POOL AND RIFFLE

XXXXXXXXXX FILL IRR. DITCH

NEW FENCE

FLOOD SLA


***** FLOOD PLAIN
***** FLOOD PLAIN

SCALE 1:1000

0 50m 100m

NOTES

1. ALL DIMENSIONS ARE METERS
UNLESS OTHERWISE NOTED.



Public Transit 516-322-6700

WISMA Group, Inc.

WATER CONSULTING, INC.

AND &

43

Mr. A. C. 1900-1901

12

144C

1430

1420

NEWARK DEPARTMENT
 OF TRANSPORTATION

Walsh Group, Inc.

INTER CONSULTING, INC.

MAR 2001



CAAD


3
2

DGN

1428510

CAMP CREEK RESTORATION







NOTES

1. ALL DIMENSIONS ARE METERS
UNLESS OTHERWISE NOTED.

- LEGEND**
- (B2) BANK PROTECTION TREATMENT TYPE 2 SEE DETAILS
 - PR POOL RIGHT
 - PL POOL LEFT
 - R RIFFLE - INCLUDE 3.6m TRANSITION BETWEEN EACH POOL AND RIFFLE
 - XXXXXXXXXX F&L IRR. DITCH
 - NEW FENCE
 - FLOOD PLAIN
 - CONST. LIMITS





LEGEND


- (B2) BANK PROTECTION TREATMENT TYPE 2 SEE DETAILS
- PR POOL RIGHT
- PL POOL LEFT
- R RIFFLE - INCLUDE 3.6m TRANSITION BETWEEN EACH POOL AND RIFFLE

- XXXXXXXXXX F&L IRR. DITCH
- NEW FENCE
- FLOOD PLAN
- CONST. LIMITS

NOTES

- ALL DIMENSIONS ARE METERS UNLESS OTHERWISE NOTED.





Appendix E

BIRD SURVEY PROTOCOL GPS PROTOCOL

*MDT Wetland Mitigation Monitoring
Camp Creek
Sula, Montana*

BIRD SURVEY PROTOCOL

The following is an outline of the MDT Wetland Mitigation Site Monitoring Bird Survey Protocol. Though each site is vastly different, the bird survey data collection methods must be standardized to a certain degree to increase repeatability. An Area Search within a restricted time frame will be used to collect the following data: a bird species list, density, behavior, and habitat-type use. There will be some decisions that team members must make to fit the protocol to their particular site. Each of the following sections and the desired result describes the protocol established to reflect bird species use over time.

Species Use within the Mitigation Wetland: Survey Method

Result: To conduct a bird survey of the wetland mitigation site within a restricted period of time and the budget allotment.

Sites that can be circumambulated or walked throughout.

These types of sites will include ponds, enhanced historic river channels, wet meadows, and any area that can be surveyed from the entirety of its perimeter or walked throughout. If the wetland is not uncomfortably inundated, conduct several “meandering” transects through the site in an orderly fashion (record the number and approximate location/direction of the transects in the field notebook; they do not have to be formalized or staked). If a very small portion of the site cannot be crossed due to inundation, this method will also apply. Though the sizes of the site vary, each site will require surveying to the fullest extent possible within a set time limit. The optimum times to conduct the survey are in the morning hours. Conduct the survey from sunrise to no later than 11:00 AM. (Note: some sites may have to be surveyed in the late afternoon or evening due to time constraints or weather; if this is the case, record the time of day and include this information in your report discussion.) If the survey is completed before 11:00 AM and no additions are being made to the list, then the task is complete. The overall limiting factor regarding the number of hours that are spent conducting this survey is the number of budgeted hours; this determination must be made by site by each individual.

In many cases, binoculars will be the only instrument that is needed to identify and count the birds using the wetland. If the wetland includes deep water habitat that can not be assessed with binoculars, then a scope and tripod are necessary. If this is the case, establish as many lookout posts as necessary from key vantage points to collect the data. Depending on the size of the open water, more time may be spent viewing the mitigation area from these vantage points than is spent walking the peripheries of more shallow-water wetlands.

Sites that cannot be circumambulated.

These types of sites will include large-bodied waters, such as reservoirs, particularly those with deep water habitat (>6 ft) close to the shore and no wetland development in that area of the shoreline. If one area of the reservoir was graded in such a way to create or enhance the development of a wetland, then that will be the area in which the ambulatory bird survey is conducted. The team member must then determine the length of the shoreline that will be surveyed during each visit.

As stated above in the ambulatory site section, these large sites most likely will have to be surveyed from established vantage points.

Species Use within the Mitigation Wetland: Data Recording

Result: A complete list of bird species using the site, an estimate of bird densities and associated behaviors, and identification of habitat use.

1. Bird Species List

Record the bird species on the Bird Survey - Field Data Sheet using the appropriate 4-letter code of the common name. The coding uses the first two letters of the first two words of the birds' common name or if one name, the first four (4) letters. For example, mourning dove is coded MODO and mallard is MALL. If an unknown individual is observed, use the following protocol and define your abbreviation at the bottom of the field data sheet: unknown shorebird: UNSB; unknown brown bird (UNBR); unknown warbler (UNWA); unknown waterfowl (UNWF). For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parentheses; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded: UNBB / FO (25). You may also note on the data sheet if that particular individual is using a constructed nest box.

2. Bird Density

In the office, sum the Bird Survey – Field Data Sheet data by species and by behavior. Record this data in the Bird Summary Table.

3. Bird Behavior

Bird behavior must be identified by what is known. When a species is simply observed, the behavior that it is immediately exhibiting is what is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair individual (BP); foraging (F); flyover (FO); loafing (L; e.g. sleeping, roosting, floating with head tucked under wing are loafing behaviors); and, nesting (N). If more behaviors are observed that do have a specific descriptive word, use them and we will add it to the protocol; descriptive words or phrases such as “migrating” or “living on site” are unknown behaviors.

4. Bird Species Habitat Use

We are interested in what bird species are using which particular habitat within the mitigation wetlands. This data is easily collected by simply recording what habitat the species was initially observed. Use the following broad category habitat classifications: aquatic bed (AB - rooted floating, floating-leaved, or submergent vegetation); forested (FO); marsh (MA – cattail, bulrush, emergent vegetation, etc. with surface water); open water (OW – primarily unvegetated); scrub-shrub (SS); and upland buffer (UP); wet meadow (WM – sedges, rushes, grasses with little to no surface water). If other categories are observed onsite that are not suggested here, we will make a new category next year.

GPS Mapping and Aerial Photo Referencing Procedure

The wetland boundaries, photograph location points and sampling locations were field located with mapping grade Trimble Geo III GPS units. The data was collected with a minimum of three positions per feature using Course/Acquisition code. The collected data was then transferred to a PC and differentially corrected to the nearest operating Community Base Station. The corrected data was then exported to ACAD drawings in Montana State Plane Coordinates NAD 83 international feet.

The GPS positions collected and processed had a 68% accuracy of 7 feet except in isolated areas of Tasks .008 and .011, where it went to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

Aerial reference points were used to position the aerial photographs. This positioning did not remove the distortion inherent in all photos; this imagery is to be used as a visual aide only. The located wetland boundaries were given a final review by the wetland biologist and adjustments were made if necessary.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

Appendix F

2005 MACROINVERTEBRATE SAMPLING PROTOCOL AND DATA

MDT Wetland Mitigation Monitoring
Camp Creek
Sula, Montana

AQUATIC INVERTEBRATE SAMPLING PROTOCOL

Equipment List

- D-frame sampling net with 1 mm mesh. Wildco is a good source of these.
- Spare net.
- 1-liter plastic sample jars, wide-mouth. VWR has these: catalog #36319-707.
- 95% ethanol: Northwest Scientific in Billings carries this.

All these other things are generally available at hardware or sporting goods stores. Make the labels on an ink jet printer preferably.

- hip waders.
- pre-printed sample labels (printed on Rite-in-the-Rain or other coated paper, two labels per sample).
- pencil.
- plastic pail (3 or 5 gallon).
- large tea strainer or framed screen.
- towel.
- tape for affixing label to jar.
- cooler with ice for sample storage.

Site Selection

Select the sampling site with these considerations in mind:

- Select a site accessible with hip waders. If substrates are too soft, lay a wide board down to walk on.
- Determine a location that is representative of the overall condition of the wetland.

Sampling

Wetland invertebrates inhabit the substrate, the water column, the stems and leaves of aquatic vegetation, and the water surface. Your goal is to sweep the collecting net through each of these habitat types, and then to combine the resulting samples into the 1-liter sample jar.

Dip out about a gallon of water into the pail. Pour about a cup of ethanol into the sample jar. Fill out the top half of the sample labels, using pencil, since ink will dissolve in the ethanol.

Ideally, you can sample a swath of water column from near-shore outward to a depth of approximately 3 feet with a long sweep of the net, keeping the net at about half the depth of the water throughout the sweep. Sweep the water surface as well. Pull the net through a vegetated area, beneath the water surface, for at least a meter of distance.

Sample the substrate by pulling the net along the bottom, bumping it against the substrate several times as you pull.

This step is optional, but it gives you a chance to see that you've collected some invertebrates. Rinse the net out into the bucket, and look for insects, crustaceans, etc. If necessary, repeat the sampling process in a nearby location, and add the net contents to the bucket. Remember to sample all four environments.

Sieve the contents of the bucket through the straining device and pour or carefully scrape the contents of the strainer into the sample jar.

If you skip the bucket-and-sieve steps, simply lift handfuls of material out of the sampling net into the jars. In either case, please include some muck or mud and some vegetation in the jar. Often, you will have collected a large amount of vegetable material. If this is the case, lift out handfuls of material from the sieve into the jar, until the jar is about half full. Please limit material you include in the sample, so that there is only a single jar for each sample.

Top off the sample jar with enough ethanol to cover all the material in the jar. Leave as little headroom as possible.

It is not necessary to sample habitats in any specified order. Keep in mind that disturbing the habitats prior to sampling will chase off the animals you are trying to capture.

Complete the sample labels. Place one label inside the sample jar and tape the other label securely to the outside of the jar. Dry the jar before attaching the outer label if necessary. In some situations, it may be necessary to collect more than one sample at a site. If you take multiple samples from the same site, clearly indicate this by using individual sample numbers, along with the total number of samples collected at the site (e.g. Sample #3 of 5 total samples).

Photograph the sampled site.

Sample Handling/Shipping

- In the field, keep collected samples cool by storing them in a cooler. Only a small amount of ice is necessary.
- Inventory all samples, preparing a list of all sites and enumerating all samples, before shipping or delivering to the laboratory.
- Deliver samples to Rhithron.

MDT Mitigated Wetland Monitoring Project

Aquatic Invertebrate Monitoring Summary 2001 - 2005

METHODS

Among other monitoring activities, aquatic invertebrate assemblages were collected at a number of mitigated wetlands throughout Montana. This report summarizes data generated from five years of collection. In 2001, 29 sites were sampled statewide. Nineteen of these sites were revisited in 2002, and 13 new sites were sampled. In 2003, 17 sites that had been visited in both 2001 and 2002 were re-sampled, and 11 sites sampled for the first time in 2001 were re-visited. In addition, 2 new sites were sampled. In 2004, 25 sites were re-visited, and 6 new sites were sampled. In 2005, an additional 2 sites were added. Over all years of sampling, a total of 151 sites were sampled for invertebrates. Table 2 summarizes sites and sampling years.

The method employed to assess these wetlands is based on an index incorporating a battery of 12 bioassessment metrics or attributes (Table 1) tested and recommended by Stribling et al. (1995) in a report to the Montana Department of Health and Environmental Science. In that study, it was determined that some of the metrics were of limited use in some geographic regions, and for some wetland types. Despite that finding, all 12 metrics are used in this evaluation of mitigated wetlands, since detailed geographic information and wetland classifications were unavailable.

Scoring criteria for metrics were developed by generally following the tactic used by Stribling et al. Boxplots were generated using a statistical software package (Statistica), and distributions, median values, ranges, and quartiles for each metric were examined. All sites in all years of sampling were used. Camp Creek, which was sampled in 2002, 2003, 2004, and 2005, and Kleinschmidt Creek, sampled in 2003, 2004, and 2005, were assessed using the tested metric battery developed for montane streams of Western Montana (Bollman 1998). Invertebrate assemblages at these sites were different from that of the other sites, and suggested montane or foothill stream conditions rather than wetland conditions. For the wetland sites, "optimal" scores were generally those that fell above the 75th percentile (for those metrics that decrease in value in response to stress) or below the 25th percentile (for metrics that respond to stress by an increase in value) of all scores. Additional scoring ranges were established by bisecting the range below the 75th percentile for decreasing scores (or above the 25th percentile for increasing scores) into "sub-optimal" and "poor" assessment categories. A score of 5, 3, or 1 was assigned to optimal, sub-optimal, and poor metric performance, respectively. In this way, metric values were translated into normalized metric scores, and scores for all metrics were summed to produce a total bioassessment score. Total bioassessment scores were classified according to a similar process, using the ranges and distributions of total scores for all sites studied in all years.

The purpose of constructing an index from biological attributes or metrics is to provide a means of integrating information to facilitate the determination of whether management action is needed. The nature of the action needed is not determined solely by the index score, however, but by consideration of an analysis of the component metrics, the taxonomic composition of the assemblages, and other issues. The diagnostic functions of the metrics and taxonomic data need more study; our understanding of the interrelationships of natural environmental factors and anthropogenic disturbances are tentative. Thus, the further interpretive remarks accompanying the raw taxonomic and metric data are offered cautiously.

Table 1. Montana Department of Transportation Mitigated Wetlands Monitoring Project sites, 2001 – 2005.

2001	2002	2003	2004	2005
Beaverhead 1	Beaverhead 1	Beaverhead 1	Beaverhead 1	Beaverhead 1
Beaverhead 2	Beaverhead 2			
Beaverhead 3	Beaverhead 3		Beaverhead 3	Beaverhead 3
Beaverhead 4	Beaverhead 4	Beaverhead 4		
Beaverhead 5	Beaverhead 5	Beaverhead 5	Beaverhead 5	Beaverhead 5
Beaverhead 6	Beaverhead 6	Beaverhead 6	Beaverhead 6	Beaverhead 6
Big Sandy 1				
Big Sandy 2				
Big Sandy 3				
Big Sandy 4				
Johnson-Valier				
VIDA				
Cow Coulee	Cow Coulee	Cow Coulee		
Fourchette – Puffin	Fourchette - Puffin	Fourchette - Puffin	Fourchette - Puffin	
Fourchette – Flashlight	Fourchette – Flashlight	Fourchette – Flashlight	Fourchette – Flashlight	
Fourchette – Penguin	Fourchette – Penguin	Fourchette – Penguin	Fourchette – Penguin	
Fourchette – Albatross	Fourchette – Albatross	Fourchette – Albatross	Fourchette – Albatross	
Big Spring	Big Spring	Big Spring	Big Spring	Big Spring
Vince Ames				
Ryegate				
Lavinia				
Stillwater	Stillwater	Stillwater	Stillwater	Stillwater
Roundup	Roundup	Roundup	Roundup	Roundup
Wigeon	Wigeon	Wigeon	Wigeon	Wigeon
Ridgeway	Ridgeway	Ridgeway	Ridgeway	Ridgeway
Musgrave – Rest. 1	Musgrave – Rest. 1	Musgrave – Rest. 1	Musgrave – Rest. 1	Musgrave – Rest. 1
Musgrave – Rest. 2	Musgrave – Rest. 2	Musgrave – Rest. 2	Musgrave – Rest. 2	Musgrave – Rest. 2
Musgrave – Enh. 1	Musgrave – Enh. 1	Musgrave – Enh. 1	Musgrave – Enh. 1	Musgrave – Enh. 1
Musgrave – Enh. 2				
	Hoskins Landing	Hoskins Landing	Hoskins Landing	Hoskins Landing
	Peterson - 1	Peterson – 1	Peterson – 1	Peterson – 1
	Peterson – 2		Peterson – 2	Peterson – 2
	Peterson – 4	Peterson – 4	Peterson – 4	Peterson – 4
	Peterson – 5	Peterson – 5	Peterson – 5	Peterson – 5
	Jack Johnson - main	Jack Johnson - main		
	Jack Johnson - SW	Jack Johnson - SW		
	Creston	Creston	Creston	Creston
	Lawrence Park			
	Perry Ranch			Perry Ranch
	SF Smith River	SF Smith River	SF Smith River	SF Smith River
	Camp Creek	Camp Creek	Camp Creek	Camp Creek
	Kleinschmidt	Kleinschmidt – pond	Kleinschmidt – pond	Kleinschmidt – pond
		Kleinschmidt – stream	Kleinschmidt – stream	Kleinschmidt – stream
		Ringling - Galt		
			Circle	
			Cloud Ranch Pond	Cloud Ranch Pond
			Cloud Ranch Stream	
			Colloid	Colloid
			Jack Creek	Jack Creek
			Norem	Norem
				Rock Creek Ranch
				Wagner Marsh

Sample Processing

Aquatic invertebrate samples were collected at mitigation wetland sites in the summer months of 2001, 2002, 2003, 2004, and 2005 by personnel of Land and Water Consulting, Inc. Sampling procedures utilized were based on the protocols developed by the Montana Department of Environmental Quality (MT DEQ). Sampling consisted of D-frame net sweeps through emergent vegetation (when present), the water column, over the water surface, and included disturbing and scraping substrates at each sampled sites. Samples were preserved in ethanol at each wetland site and subsequently delivered to Rhithron Associates, Inc. for processing, taxonomic determinations, and data analysis.

At Rhithron's laboratory, Caton subsamplers and stereomicroscopes with 10X magnification were used to randomly select a minimum of 100 organisms, when possible, from each sample. In some cases, the entire sample contained fewer than 100 organisms; in these cases, all organisms from the sample were taken. Taxa were identified in general accordance with the taxonomic resolution standards set out in the MT DEQ Standard Operating Procedures for Sampling and Sample Analysis (Bukantis 1998). All samples were re-identified by a second taxonomist for quality assurance purposes. The identified samples have been archived at Rhithron's laboratory. Taxonomic data and organism counts were entered into an Excel 2000 spreadsheet, and metrics were calculated and scored using spreadsheet formulae.

Bioassessment Metrics

An index based on the performance of 12 metrics was constructed, as described above. Table 2 lists those metrics, describes their calculation and the expected response of each to increased degradation or impairment of the wetland.

In addition to the summed scores of each metric and the associated impairment classification described above, each individual metric informs the bioassessment to some degree. The four richness metrics (Total taxa, POET, Chironomidae taxa, and Crustacea taxa + Mollusca taxa) can be interpreted to express habitat complexity as well as water quality. Complex, diverse habitats consist of variable substrates, emergent vegetation, variable water depths and other factors, and are potential features of long-established stable wetlands with minimal human disturbance. In the study conducted by Stribling et al. (1995), all four richness metrics were found to be significantly associated with water quality parameters including conductance, salinity, and total dissolved solids.

Four composition metrics (%Chironomidae, %Orthocladiinae of Chironomidae, %Crustacea + %Mollusca, and %Amphipoda) measure the relative contributions of certain taxonomic groups that may have significant responses to habitat and/or water quality impacts. For example, amphipods have been demonstrated to increase in abundance in alkaline conditions. Short-lived, relatively mobile taxa such as chironomids dominate ephemeral environments; many are hemoglobin-bearers capable of tolerating de-oxygenated conditions.

Two tolerance metrics (the Hilsenhoff Biotic Index and %Dominant taxon) were included in the bioassessment battery. The HBI indicates the overall invertebrate assemblage tolerance to nutrient enrichment, warm water, and/or low dissolved oxygen conditions. The percent abundance of the dominant taxon has been demonstrated to be strongly associated with pH, conductance, salinity, total organic carbon, and total dissolved solids.

Two trophic measures (%Collector-gatherers and %Filterers) may be helpful in expressing functional integrity of the invertebrate assemblage, which can be impacted by poor water quality or habitat degradation. High proportions of filtering organisms suggest nutrient and/or organic enrichment, while abundant collectors suggest more positive functional conditions and well-developed wetland morphology. These organisms graze periphyton growing on stable surfaces such as macrophytes.

Metric scoring criteria were re-examined each year as new data was added. For 2005, all 151 records were utilized. Ranges of individual metrics, as well as median metric values remained remarkably consistent over all 5 years of analysis. Since metric value distributions changed insignificantly with the addition of the 2005 data, no changes were made to scoring criteria this year. Summary metric values and scores for the 2005 samples are given in Tables 3a-3d.

Table 2. Aquatic invertebrate metrics employed in the MTDT mitigation wetland monitoring study, 2001-2005.

Metric	Metric calculation	Expected response to degradation or impairment
Total taxa	Count of unique taxa identified to lowest recommended taxonomic level	Decrease
POET	Count of unique Plecoptera, Trichoptera, Ephemeroptera, and Odonata taxa identified to lowest recommended taxonomic level	Decrease
Chironomidae taxa	Count of unique midge taxa identified to lowest recommended taxonomic level	Decrease
Crustacea taxa + Mollusca taxa	Count of unique Crustacea taxa and Mollusca taxa identified to lowest recommended taxonomic level	Decrease
% Chironomidae	Percent abundance of midges in the subsample	Increase
Orthoclaadiinae/Chironomidae	Number of individual midges in the sub-family Orthoclaadiinae / total number of midges in the subsample.	Decrease
% Amphipoda	Percent abundance of amphipods in the subsample	Increase
% Crustacea + % Mollusca	Percent abundance of crustaceans in the subsample plus percent abundance of molluscs in the subsample	Increase
HBI	Relative abundance of each taxon multiplied times that taxon's modified Hilsenhoff Biotic Index value. These numbers are summed over all taxa in the subsample.	Increase
% Dominant taxon	Percent abundance of the most abundant taxon in the subsample	Increase
% Collector-Gatherers	Percent abundance of organisms in the collector-gatherer functional group	Decrease
% Filterers	Percent abundance of organisms in the filterer functional group	Increase

RESULTS

(Note: Individual site discussions were removed from this report by Land & Water Consulting / PBS&J and are included in the Macro-Invertebrate sections of individual reports. Summary tables are provided on the following pages.)

Table 3a. Metric values and scores for Montana Department of Transportation mitigated wetland sites in 2005.

	BEAVERHEAD #1	BEAVERHEAD #3	BEAVERHEAD #5	BEAVERHEAD #6	BIG SPRING CREEK	STILLWATER	ROUNDUP	WIDGEON
Total taxa	22	9	14	18	28	17	7	19
POET	2	0	0	2	4	4	0	0
Chironomidae taxa	7	4	4	4	9	5	3	11
Crustacea + Mollusca	4	3	1	4	7	5	2	4
% Chironomidae	59.80%	7.55%	50.00%	16.67%	33.65%	9.43%	22.22%	76.47%
Orthoclaadiinae/Chir	0.197	0.625	0.059	0.067	0.457	0.500	0.000	0.205
% Amphipoda	1.96%	0.94%	0.00%	1.11%	18.27%	7.55%	0.00%	10.78%
% Crustacea + % Mollusca	10.78%	90.57%	2.94%	55.56%	33.65%	53.77%	72.65%	15.69%
HBI	7.71	7.88	7.88	7.98	7.55	7.28	8.33	8.25
% Dominant taxon	34.31%	76.42%	35.29%	25.56%	18.27%	33.02%	71.79%	44.12%
% Collector-Gatherers	56.86%	93.40%	47.06%	21.11%	70.19%	64.15%	82.05%	26.47%
% Filterers	0.00%	0.00%	0.00%	0.00%	0.96%	3.77%	0.00%	6.86%
Total taxa	5	1	1	3	5	3	1	3
POET	1	1	1	1	5	5	1	1
Chironomidae taxa	5	3	3	3	5	3	3	5
Crustacea + Mollusca	3	1	1	3	5	3	1	3
% Chironomidae	1	5	1	5	3	5	3	1
Orthoclaadiinae/Chir	3	5	1	1	5	5	1	3
% Amphipoda	5	5	5	5	3	3	5	3
% Crustacea + % Mollusca	5	1	5	3	3	3	1	5
HBI	1	1	1	1	3	3	1	1
% Dominant taxon	3	1	3	5	5	5	1	3
% Collector-Gatherers	3	5	3	1	3	3	5	1
% Filterers	3	3	3	3	3	3	3	1
Total score	38	32	28	34	48	44	26	30
Percent of maximum score	0.633333	0.533333	0.466667	0.566667	0.8	0.733333	0.433333	0.5
Impairment classification	sub-optimal	poor	poor	sub-optimal	optimal	optimal	poor	poor

Table 3b. Metric values and scores for Montana Department of Transportation mitigated wetland sites in 2005.

	RIDGEWAY	MUSGRAVE REST. 1	MUSGRAVE REST. 2	MUSGRAVE ENH. 1	HOSKINS LANDING	PETERSON RANCH 1	PETERSON RANCH 2	PETERSON RANCH 4	PETERSON RANCH 5
Total taxa	19	19	23	19	27	29	16	25	16
POET	3	1	3	1	5	4	2	4	4
Chironomidae taxa	6	6	8	3	6	11	6	8	7
Crustacea + Mollusca	5	5	3	7	6	6	5	6	2
% Chironomidae	9.26%	14.55%	22.00%	2.80%	17.58%	17.48%	13.91%	24.55%	16.96%
Orthoclaadiinae/Chir	0.600	0.750	0.136	0.667	0.188	0.556	0.563	0.630	0.632
% Amphipoda	6.48%	3.64%	0.00%	0.93%	0.00%	0.97%	7.83%	1.82%	8.04%
% Crustacea + % Mollusca	22.22%	30.91%	38.00%	58.88%	27.47%	31.07%	72.17%	20.00%	8.93%
HBI	7.71	7.22	7.77	7.16	6.81	7.16	7.43	7.65	8.08
% Dominant taxon	53.70%	21.82%	35.00%	28.04%	14.29%	26.21%	33.04%	18.18%	31.25%
% Collector-Gatherers	68.52%	40.00%	15.00%	11.21%	31.87%	59.22%	28.70%	43.64%	68.75%
% Filterers	0.00%	0.00%	0.00%	2.80%	0.00%	4.85%	33.91%	5.45%	1.79%
Total taxa	3	3	5	3	5	5	3	5	3
POET	3	1	3	1	5	5	1	5	5
Chironomidae taxa	3	3	5	3	3	5	3	5	5
Crustacea + Mollusca	3	3	1	5	5	5	3	5	1
% Chironomidae	5	5	3	5	5	5	5	3	5
Orthoclaadiinae/Chir	5	5	1	5	3	5	5	5	5
% Amphipoda	3	5	5	5	5	5	3	5	3
% Crustacea + % Mollusca	5	5	3	3	5	5	1	5	5
HBI	1	3	1	3	5	3	3	1	1
% Dominant taxon	1	5	3	5	5	5	5	5	5
% Collector-Gatherers	3	1	1	1	1	3	1	1	3
% Filterers	3	3	3	3	3	3	1	3	3
Total score	38	42	34	42	50	54	34	48	44
Percent of maximum score	0.633333	0.7	0.566667	0.7	0.833333	0.9	0.566667	0.8	0.733333
Impairment classification	sub-optimal	optimal	sub-optimal	optimal	optimal	optimal	sub-optimal	optimal	optimal

Table 3c. Metric values and scores for Montana Department of Transportation mitigated wetland sites in 2005.

	CRESTON	PERRY RANCH	SOUTH FORK SMITH RIVER	CAMP CREEK	KLEINSCH MIDT POND	KLEINSCH MIDT STREAM	CLOUD RANCH POND	COLLOID	JACK CREEK
Total taxa	16	18	19	36	27	23	22	9	16
POET	0	0	4	14	6	5	2	1	1
Chironomidae taxa	4	8	6	13	6	9	11	4	9
Crustacea + Mollusca	6	4	5	0	2	3	3	1	4
% Chironomidae	27.62%	43.69%	21.67%	45.54%	8.85%	45.08%	37.50%	25.83%	29.41%
Orthoclaadiinae/Chir	0.931	0.622	0.192	0.804	0.200	0.473	0.256	0.000	0.467
% Amphipoda	0.00%	0.00%	29.17%	0.00%	5.31%	0.82%	0.00%	0.00%	0.98%
% Crustacea + % Mollusca	52.38%	38.83%	62.50%	0.00%	7.96%	3.28%	7.69%	67.50%	41.18%
HBI	7.52	7.31	7.54	5.06	7.40	5.83	6.96	8.53	7.39
% Dominant taxon	25.71%	25.24%	29.17%	18.81%	30.09%	32.79%	41.35%	67.50%	35.29%
% Collector-Gatherers	64.76%	47.57%	65.00%	47.52%	37.17%	50.82%	75.96%	88.33%	91.18%
% Filterers	6.67%	27.18%	8.33%	5.94%	0.88%	2.46%	2.88%	0.00%	2.94%
Total taxa	3	3	3	5	5	5	5	1	3
POET	1	1	5	5	5	5	1	1	1
Chironomidae taxa	3	5	3	5	3	5	5	3	5
Crustacea + Mollusca	5	3	3	1	1	1	1	1	3
% Chironomidae	3	1	3	1	5	1	3	3	3
Orthoclaadiinae/Chir	5	5	3	5	3	5	3	1	1
% Amphipoda	5	5	1	5	3	5	5	5	5
% Crustacea + % Mollusca	3	3	3	5	5	5	5	1	3
HBI	3	3	3	5	3	5	3	1	3
% Dominant taxon	5	5	5	5	5	5	3	1	3
% Collector-Gatherers	3	3	3	3	1	3	3	5	5
% Filterers	1	1	1	3	3	3	3	3	3
Total score	40	38	36	48	42	48	40	26	38
Percent of maximum score	0.666667	0.633333	0.6	0.8	0.7	0.8	0.666667	0.433333	0.633333
Impairment classification	sub-optimal	sub-optimal	sub-optimal	optimal	optimal	optimal	sub-optimal	poor	sub-optimal

Table 3d. Metric values and scores for Montana Department of Transportation mitigated wetland sites in 2005.

	NOREM	ROCK CREEK RANCH	WAGNER MARSH
Total taxa	4	24	23
POET	0	2	5
Chironomidae taxa	2	8	8
Crustacea + Mollusca	2	4	5
% Chironomidae	37.50%	22.00%	24.00%
Orthocladiinae/Chir	0.000	0.318	0.167
% Amphipoda	0.00%	3.00%	7.00%
% Crustacea + % Mollusca	62.50%	40.00%	19.00%
HBI	7.50	7.61	8.58
% Dominant taxon	56.25%	18.00%	38.00%
% Collector-Gatherers	6.25%	57.00%	40.00%
% Filterers	0.00%	0.00%	3.00%
Total taxa	1	5	5
POET	1	1	5
Chironomidae taxa	1	5	5
Crustacea + Mollusca	1	3	3
% Chironomidae	3	3	3
Orthocladiinae/Chir	1	3	1
% Amphipoda	5	5	3
% Crustacea + % Mollusca	3	3	5
HBI	3	1	1
% Dominant taxon	1	5	3
% Collector-Gatherers	1	3	1
% Filterers	3	3	3
Total score	24	40	38
Percent of maximum score	0.4	0.666667	0.633333
Impairment classification	poor	sub-optimal	sub-optimal

Literature Cited

Bollman, W. 1998. Montana Valleys and Foothill Prairies Ecoregion. Master's Thesis. (M.S.) University of Montana. Missoula, Montana.

Bukantis, R. 1998. Rapid bioassessment macroinvertebrate protocols: Sampling and sample analysis SOP's. Working draft. Montana Department of Environmental Quality. Planning Prevention and Assistance Division. Helena, Montana.

McCune, B. and J.B. Grace. 2002. Analysis of Ecological Communities. MjM Software Design, Gleneden Beach, Oregon, USA.

McCune, B. and M.J. Mefford. 2002. PC-ORD. Multivariate Analysis of Ecological Data, Version 4. MjM Software Design, Gleneden Beach, Oregon, USA.

Stribling, J.B., J. Lathrop-Davis, M.T. Barbour, J.S. White, and E.W. Leppo. 1995. Evaluation of environmental indicators for the wetlands of Montana: the multimetric approach using benthic macroinvertebrates. Report to the Montana Department of Health and Environmental Science. Helena, Montana.

Taxa Listing

Project ID: MDT05LW
RAI No.: MDT05LW016

RAI No.: MDT05LW016

Sta. Name: CAMP CREEK

Client ID:

Date Coll.: 8/1/2005

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
Non-Insect							
Acari	2	2.04%	Yes	Unknown		5	PR
Branchiobdellidae							
Branchiobdellidae	1	1.02%	Yes	Unknown		6	PA
Enchytraeidae							
Enchytraeidae	1	1.02%	Yes	Unknown		4	CG
Naididae							
Naididae	3	3.06%	Yes	Unknown		8	CG
Ephemeroptera							
Ameletidae							
<i>Ameletus</i> sp.	1	1.02%	Yes	Larva		0	CG
Baetidae							
<i>Baetis tricaudatus</i>	1	1.02%	Yes	Larva		4	CG
Ephemerellidae							
<i>Drunella grandis</i>	1	1.02%	Yes	Larva		2	SC
<i>Serratella tibialis</i>	1	1.02%	Yes	Larva		2	CG
<i>Timpanoga hecuba</i>	1	1.02%	Yes	Larva		2	CG
Heptageniidae							
<i>Nixe</i> sp.	1	1.02%	Yes	Larva		4	SC
Plecoptera							
Pteronarcyidae							
<i>Pteronarcella</i> sp.	1	1.02%	Yes	Larva	Early Instar	4	SH
<i>Pteronarcys</i> sp.	3	3.06%	Yes	Larva	Early Instar	2	SH
Trichoptera							
Brachycentridae							
Brachycentridae	2	2.04%	Yes	Larva	Early Instar	1	CG
<i>Micrasema</i> sp.	2	2.04%	Yes	Larva		1	SH
Hydroptilidae							
<i>Hydroptila</i> sp.	1	1.02%	Yes	Larva		6	PH
<i>Ochrotrichia</i> sp.	2	2.04%	Yes	Larva		4	PH
Lepidostomatidae							
<i>Lepidostoma</i> sp.	1	1.02%	Yes	Larva		1	SH
Limnephilidae							
<i>Psychoglypha</i> sp.	2	2.04%	Yes	Larva		0	CG
Coleoptera							
Dytiscidae							
<i>Oreodytes</i> sp.	1	1.02%	Yes	Adult		5	PR
Elmidae							
<i>Optioservus</i> sp.	7	7.14%	Yes	Adult		5	SC
<i>Optioservus</i> sp.	12	12.24%	No	Larva	Larva	5	SC
Diptera							
Athericidae							
<i>Atherix</i> sp.	2	2.04%	Yes	Larva		5	PR
Simuliidae							
Simuliidae	1	1.02%	No	Pupa	Pupa	6	CF
<i>Simulium</i> sp.	5	5.10%	Yes	Larva		6	CF

Monday, November 07, 2005

Taxa Listing

Project ID: MDT05LW
RAI No.: MDT05LW016

RAI No.: MDT05LW016

Sta. Name: CAMP CREEK

Client ID:

Date Coll.: 8/1/2005

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
Chironomidae							
Chironomidae							
Chironomidae	3	3.06%	No	Pupa	Pupa	10	CG
<i>Cricotopus (Cricotopus) sp.</i>	2	2.04%	Yes	Larva		7	SH
<i>Cricotopus (Nostococladius) sp.</i>	1	1.02%	Yes	Larva		6	SH
<i>Cricotopus bicinctus</i>	6	6.12%	Yes	Larva		7	SH
<i>Eukiefferiella sp.</i>	2	2.04%	No	Larva	Early Instar	8	CG
<i>Eukiefferiella Brehmi Gr.</i>	3	3.06%	Yes	Larva		8	CG
<i>Eukiefferiella Devonica Gr.</i>	1	1.02%	Yes	Larva		8	CG
<i>Micropsectra sp.</i>	5	5.10%	Yes	Larva		4	CG
<i>Orthocladius sp.</i>	15	15.31%	Yes	Larva		6	CG
<i>Parametriocnemus sp.</i>	1	1.02%	Yes	Larva		5	CG
<i>Stempellinella sp.</i>	1	1.02%	Yes	Larva		4	CG
<i>Tvetenia Bavarica Gr.</i>	2	2.04%	Yes	Larva		5	CG
<i>Tvetenia vitracies</i>	1	1.02%	Yes	Larva		5	CG
Sample Count	98						

Metrics Report


Project ID: MDT05LW
RAI No.: MDT05LW016
Sta. Name: CAMP CREEK
Client ID:
STORET ID
Coll. Date: 8/1/2005

Abundance Measures

Sample Count: 98
Sample Abundance: 420.00 23.33% of sample used
Total Abundance: 564.90
Coll. Procedure:
Sample Notes:

Taxonomic Composition

Category	R	A	PRA
Non-Insect	4	7	7.14%
Odonata			
Ephemeroptera	6	6	6.12%
Plecoptera	2	4	4.08%
Heteroptera			
Megaloptera			
Trichoptera	6	10	10.20%
Lepidoptera			
Coleoptera	2	20	20.41%
Diptera	2	8	8.16%
Chironomidae	11	43	43.88%




Dominant Taxa

Category	A	PRA
Optioservus	19	19.39%
Orthocladus	15	15.31%
Cricotopus binctus	6	6.12%
Simulium	5	5.10%
Micropsectra	5	5.10%
Pteronarcys	3	3.06%
Naididae	3	3.06%
Eukiefferiella Brehmi Gr.	3	3.06%
Chironomidae	3	3.06%
Tvetenia Bavarica Gr.	2	2.04%
Psychodgylpha	2	2.04%
Ochrotrichia	2	2.04%
Brachycentridae	2	2.04%
Atherix	2	2.04%
Acari	2	2.04%

Functional Composition

Category	R	A	PRA
Predator	3	5	5.10%
Parasite	1	1	1.02%
Collector Gatherer	16	46	46.94%
Collector Filterer	1	6	6.12%
Macrophyte Herbivore			
Piercer Herbivore	2	3	3.06%
Xylophage			
Scraper	3	21	21.43%
Shredder	7	16	16.33%
Omnivore			
Unknown			




Metric Values and Scores

Metric	Value	BIBI	MTP	MTV	MTM
<i>Composition</i>					
Taxa Richness	33	3	3		3
Non-Insect Percent	7.14%				
E Richness	6	3		3	
P Richness	2	1		2	
T Richness	6	3		3	
EPT Richness	14		3		0
EPT Percent	20.41%		1		0
Oligochaeta+Hirudinea Percent	5.10%				
Baetidae/Ephemeroptera	0.167				
Hydropsychidae/Trichoptera	0.000				
<i>Dominance</i>					
Dominant Taxon Percent	19.39%		3		3
Dominant Taxa (2) Percent	34.69%				
Dominant Taxa (3) Percent	40.82%	5			
Dominant Taxa (10) Percent	65.31%				
<i>Diversity</i>					
Shannon H (loge)	3.106				
Shannon H (log2)	4.481		3		
Margalef D	7.303				
Simpson D	0.056				
Evenness	0.045				
<i>Function</i>					
Predator Richness	3		1		
Predator Percent	5.10%	1			
Filterer Richness	1				
Filterer Percent	6.12%			2	
Collector Percent	53.06%		3		3
Scraper+Shredder Percent	37.76%		3		1
Scraper/Filterer	3.500				
Scraper/Scraper+Filterer	0.778				
<i>Habit</i>					
Burrower Richness	0				
Burrower Percent	0.00%				
Swimmer Richness	3				
Swimmer Percent	3.06%				
Clinger Richness	15	3			
Clinger Percent	50.00%				
<i>Characteristics</i>					
Cold Stenotherm Richness	2				
Cold Stenotherm Percent	3.06%				
Hemoglobin Bearer Richness					
Hemoglobin Bearer Percent					
Air Breather Richness	1				
Air Breather Percent	1.02%				
<i>Volturnism</i>					
Univoltine Richness	13				
Semivoltine Richness	4	3			
Multivoltine Percent	50.00%		2		
<i>Tolerance</i>					
Sediment Tolerant Richness	0				
Sediment Tolerant Percent	0.00%				
Sediment Sensitive Richness	1				
Sediment Sensitive Percent	1.02%				
Metals Tolerance Index	4.158				
Pollution Sensitive Richness	2			2	
Pollution Tolerant Percent	24.49%	3		1	
Hilsenhoff Biotic Index	5.173		2		0
Intolerant Percent	14.29%				
Supertolerant Percent	12.24%				
CTQa	73.083				

Bioassessment Indices

BioIndex	Description	Score	Pct	Rating
BIBI	B-IBI (Karr et al.)	26	52.00%	
MTP	Montana DEQ Plains (Bukantis 1998)	24	80.00%	Slight
MTV	Montana Revised Valleys/Foothills (Bollman 1998)	13	72.22%	Slight
MTM	Montana DEQ Mountains (Bukantis 1998)	10	47.62%	Moderate




Appendix G


FIGURE 5 - CAMP CREEK CHANNEL CROSS SECTIONS PLANTING SPECIFICATIONS

*MDT Wetland Mitigation Monitoring
Camp Creek
Sula, Montana*

Cross Section 3-A



Cross Section 4-A



LEGEND

- Existing Surface 12-3-02
- Existing Surface 9-10-03
- Existing Surface 9-24-04
- Existing Surface 8-01-05

Nature's Enhancement, Inc.

2980 Eastside Highway
 Stevensville, Montana 59870
 Phone: (406) 777-3560
 FAX: (406) 777-3560

SOLD TO:

Department of Transportation

Project No:

NH7-1(58)9 F Sula-North & South

NH41(24) Camp Creek Restoration

SHIPPED TO:

Sula North & South/ Camp Creek Restoration
 Project Site
 Sula, Montana

MONITORING

INVOICE NUMBR
 PURCHASE ORDER #
 ORDER DATE
 SHIP DATE (EST.)
 TERMS
 DUE DATE
 SALES REP
 SHIP VIA

Greg
 NE

GC5: REVEGETATION

446	Alnus Incana	1 Gallon	1-2'
315	Alnus Incana	5 Gallon	3-4'
752	Amelanchier alnifolia	1 Gallon	1-2'
374	Betula occidentalis	5 Gallon	3-4'
687	Cornus stolonifera	1 Gallon	2-3'
369	Cornus stolonifera	5 Gallon	4-5'
213	Pinus contorta	1 Gallon	1-2'
89	Pinus contorta	5 Gallon	2-3'
213	Pinus ponderosa	1 Gallon	1-2'
89	Pinus ponderosa	5 Gallon	2-3'
303	Populus tremuloides	1 Gallon	18-24"SS
15	Populus tremuloides	5 Gallon	4-5"SS
791	Populus tremuloides	1 Gallon	18-24"MS
311	Populus tremuloides	5 Gallon	4-5"MS
800	Populus trichocarpa	1 Gallon	2-3'
518	Populus trichocarpa	5 Gallon	5-8'
2025	Potentilla fruticosa	1 Gallon	12-18"
213	Pseudotsuga menziesii	1 Gallon	12-18"
89	Pseudotsuga menziesii	5 Gallon	24-30"
1178	Rosa woodsii	1 Gallon	2-3'
1902	Willow (Salix spp.)	1 Gallon	2-3"MS



429	Willow (Salix spp.)	5 Gallon	4-5 MS
1178	Symphoricarpos albus	1 Gallon	18-24
10681	Installation of above 1 Gallon Plants		
2598	Installation of above 5 Gallon Plants		
20,480	Willow Cuttings 12" long with a minimum base of .75 inches (800/Hectare) Includes collection, installation	12" x .75 Base	
57	<u>WILLOW SALVAGE</u> Tree Spade dig at a minimum diameter of 24", burlap, basket, crimp, tie Storage of the above on site in .75m fine soil, to be provided by prime contractor Replant willow clumps		
Shipping Charges:			
Common Carrier (CMN):		billed COD from the trucking company.	
Nature's Enhancement (NE):		billed from NE on the Final Invoice.	
Nursery Pick Up (NPU):		no charge.	
		SUBTOTAL	
		QTY. DISCOUNT	Included
		SHIPPING (ESTIMATE)	Included
		BOXING & HANDLING	Included
		TOTAL	0.00

Questions concerning this order?

Call: PHONE: (406) 777-3560

FAX: (406) 777-3500

MAKE ALL CHECKS PAYABLE TO:

Nature's Enhancement, Inc.
2980 Eastside Highway
Stevensville, Montana 59870

\$0.00
PAY THIS
AMOUNT

THANK YOU FOR YOUR ORDER! WE LOOK FORWARD TO SERVING YOU AGAIN.

U.S. 6702

SEED BLENDING REPORT
Dept. of Transportation, Great Falls, MT

1-Materials Bureau, (Pat Hoy)
1-District Lab Gt. Falls
1-E.P.M. *T. DENIEDIK.*

PROJECT NO.: NH 41(24)

TERMINI: CAMP CREEK RESTORATION

MISSOULA

BLENDING WITNESSED BY: JAMES O. BLOSSOM

DATE: 04/22/2002

LOCATION: Fairfield, Montana

SEED SUPPLIER: Treasure State Seed Inc.

SEED SUPPLIER: Treasure State Seed Inc.		ACRES Hectare Area 1/.25		(e) Total Bulk Seed Blended For Area 1	MSU Seed Laboratory test results			Hectare Area 2		(e) Total Bulk Seed Blended for Area 2	Mat'l's. Bureau Pretest Lab. No	MSU Test Date Expires
Type Of Seed	Lot No.	LOS kg Pls per AC.	(d) Total kg LOS Pls		(a) % Purity	(b) % Germ	(c) % Pls	kg Pls Per ha	(d) Total kg Pls			
MEADOW BARLEY •	NBS-1-05381	0.5	0.6	0.7	93.59	97	90.78					
BLUEJOINT REEDGRASS •	CACA24204	0.3	0.4	0.6	85.85	77	66.13					
FOWL BLUEGRASS •	00-043	2.0	2.5	3.2	86.91	89	77.35					
TUFFED HAIRGRASS •	99-1438-75	2.0	2.5	2.7	94.2	99	93.25					
BLUE WILCRYE •	6BS-0-300	7.0	8.8	9.7	99.05	92	91.15					
BROMAR MOUNTAIN BROME	006-026-124	6.0	7.5	7.8	98.85	97	95.88					
TOTAL				24.7 LOS								

BULK SEEDING RATE AREA 1 19.76 KILOGRAMS (kg) PER HECTARE (ha). BULK AREA 2 KILOGRAMS (kg) PER HECTARE (ha)

$$\% \text{ PURITY (a)} \times \% \text{ GERMINATION (b)} = \% \text{ PURE LIVE SEED @ } \times 100.$$

TOTAL KILOGRAMS (kg) PURE LIVE SEED (d) = % PURE LIVE SEED © X 100 = BULK SEED NEEDED (e)

REMARKS: