
MONTANA DEPARTMENT OF TRANSPORTATION WETLAND MITIGATION MONITORING REPORT: YEAR 2008

*Wagner Marsh
Billings, Montana*



Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Avenue
Helena, MT 59620-1001

Prepared by:

POST, BUCKLEY, SCHUH, AND JERNIGAN
801 North Last Chance Gulch, Suite 101
Helena, MT 59601-3360

December 2008

PBS&J Project No: 0B4308801.06.07



MONTANA DEPARTMENT OF TRANSPORTATION

WETLAND MITIGATION MONITORING REPORT:

YEAR 2008

*Wagner Marsh
Billings, Montana*

Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Ave
Helena, MT 59620-1001

Prepared by:

POST, BUCKLEY, SCHUH, AND JERNIGAN
801 North Last Chance Gulch, Suite 101
Helena, MT 59601-3360

December 2008

PBS&J Project No: 0B4308801.06.07

“MDT attempts to provide accommodations for any known disability that may interfere with a person participating in any service, program, or activity of the Department of Transportation. Alternative accessible formats of this information will be provided upon request. For further information, call 406-444-7228 or TTY (800-335-7592) or by calling Montana Relay at 711.”



TABLE OF CONTENTS

1.0 INTRODUCTION.....	1
2.0 METHODS.....	3
2.1 Monitoring Dates and Activities.....	3
2.2 Hydrology	3
2.3 Vegetation.....	3
2.4 Soils.....	4
2.5 Wetland Delineation	4
2.6 Mammals, Reptiles and Amphibians	4
2.7 Birds.....	5
2.8 Macroinvertebrates	5
2.9 Functional Assessment.....	5
2.10 Photographs.....	6
2.11 GPS Data.....	6
2.12 Maintenance Needs.....	6
3.0 RESULTS	6
3.1 Hydrology	6
3.2 Vegetation.....	8
3.3 Soils.....	12
3.4 Wetland Delineation	12
3.5 Wildlife and Fish.....	13
3.6 Macroinvertebrates	14
3.7 Functional Assessment.....	15
3.8 Photographs.....	15
3.9 Maintenance Needs/Recommendations	16
3.10 Current Credit Summary.....	16
4.0 REFERENCES.....	18

TABLES

Table 1	<i>2005 - 2008 vegetation species list for the Wagner Marsh Wetland Mitigation Site.</i>
Table 2	<i>2005 – 2008 vegetation transect data summary.</i>
Table 3	<i>2008 observed mortality of planted woody species for the Wagner Marsh Wetland Mitigation Site.</i>
Table 4	<i>Fish and wildlife species observed at the Wagner Marsh Mitigation Site during 2005 to 2008.</i>
Table 5	<i>Summary of 2001 and 2005 through 2008 wetland function/value ratings and functional points at the Wagner Marsh Wetland Mitigation Site.</i>
Table 6	<i>Summary of open water and wetland acreages at the Wagner Marsh Wetland Mitigation Site for 2001, 2005, 2006, 2007, and 2008.</i>
Table 7	<i>2008 mitigation credit summary for the Wagner Marsh Wetland Mitigation Site.</i>

FIGURES

Figure 1	<i>Project Site Location Map</i>
Figure 2	<i>Monitoring Activity Locations 2008</i>
Figure 3	<i>Mapped Site Features 2008</i>

CHARTS

Chart 1	<i>An example of the variation in groundwater levels at the Wagner Marsh Wetland Mitigation Site.</i>
Chart 2	<i>Transect maps showing vegetation types from the start of transect (0 feet) to the end of transect (530 feet) for each year monitored.</i>
Chart 3	<i>Length of vegetation communities within Transect 1 for each year monitored.</i>
Chart 4	<i>Macroinvertebrate bioassessment scores for the Wagner Marsh Wetland Mitigation Site from 2005 to 2008.</i>

APPENDICES

Appendix A	<i>Figures 2 & 3</i>
Appendix B	<i>2008 Wetland Mitigation Site Monitoring Form</i>
	<i>2008 Bird Survey Forms</i>
	<i>2008 COE Wetland Delineation Forms</i>
	<i>2008 Functional Assessment Forms</i>

APPENDICES (continued)

Appendix C *2008 Representative Photographs*

Appendix D *Conceptual Site Layout*

Appendix E *Bird Survey Protocol*

GPS Protocol

Appendix F *Macroinvertebrate Sampling Protocol and Data*

1.0 INTRODUCTION

This report presents the results of the fourth year (2008) of wetland monitoring at the Wagner Marsh wetland mitigation project. This mitigation site was constructed during the spring of 2005 in the eastern portion of the Upper Yellowstone River watershed (Watershed #13). It is anticipated that this site will compensate for wetland impacts resulting from Montana Department of Transportation (MDT) highway and bridge reconstruction projects in the watershed. Wagner Marsh was constructed on MDT property originally purchased in 1954 and used as a borrow area (gravel mining) for construction of the Interstate 90 (I-90) corridor. For this reason the Wagner Marsh is also known as the 'Wagner Pit'. The goal of the project is to create wetland hydrology at the site, and thereby ultimately provide approximately 21.59 acres of palustrine emergent and scrub-shrub wetland within the confines of the 39 acre site. Prior to construction, approximately 2.12 acres of palustrine emergent and scrub-shrub wetland and 1.75 acres of open water had been incidentally created by MDT via pit excavation.




The site occurs at an elevation of approximately 3,240 feet above mean sea level and is located on the west edge of Billings, MT just north and east of the intersection of Danford Road and 56th Street in the SW ¼ of Section 28, Township 1 South, Range 25 East, Yellowstone County (**Figure 1**). Approximate universal transverse mercator (UTM) coordinates for the central portion of the site are (Zone 12N) 5,065,220 Northing, 682,385 Easting.

The approximate site boundary is illustrated in **Figure 2 (Appendix A)**, and the original conceptual layout is provided in **Appendix D**. The project incorporates the two incidentally created wetland/open water areas totaling 3.87 acres and seven wetland creation areas (i.e., wetland cells) totaling approximately 17.72 acres for a total projected aquatic habitat size of 21.59 acres. Wetland hydrology is supplied primarily through interception of the groundwater table, with some minimal contributions from precipitation. No surface outlet exists at the site. To ensure sufficient water for the wetland creation areas into the future, MDT previously secured groundwater rights. The establishment of an upland buffer is also a part of this project and is tied into the crediting for the project. Monitoring occurs on the site in mid-summer when wetland data is collected, and in the fall when bird and other wildlife use is documented.

Wetland credits for the site are determined by the following ratios:

- 1:1 for wetland establishment/reestablishment for in-kind mitigation conducted prior to wetland impacts
- 1.5:1 for out-of-kind wetland mitigation, or if wetland impacts occurred prior to the reserve's establishment
- Credit for open water is limited to no more than 20% of the amount of actual wetland acreage that develops onsite.
- Upland buffers are limited to a maximum width of 50 feet and are credited at a ratio of 4:1.

FIGURE 1. PROJECT LOCATION
Wagner Marsh
Mitigation Site



0 800 1,600
FEET
1:24,000

PROJECT #: 0B4308801
DATE: NOVEMBER 2008
LOCATION: WAGNER MARSH
PROJECT MGR: J. BERGLUND
DRAWN BY: JJC



801 N. LAST CHANCE GULCH
SUITE 101
HELENA, MT 59601-3360

2.0 METHODS

2.1 Monitoring Dates and Activities

The site was visited on August 8, 2008 (mid-season visit) and again on October 7, 2008 (fall visit). The mid-season visit was conducted to document vegetation, soil, and hydrologic conditions used to map jurisdictional wetlands. The majority of the information contained on the Wetland Mitigation Site Monitoring Form (**Appendix B**) was collected at this time. Activities and information conducted/collected included: wetland delineation; wetland/open water boundary mapping; vegetation community mapping; vegetation transects; soils data; hydrology data; bird and general wildlife use; photograph points; macroinvertebrate sampling; functional assessment; and survival of planted woody vegetation.

The primary purpose of the fall visit was to conduct bird/general wildlife reconnaissance of the site. The fall visit was timed to coincide with fall bird migrations. Based on past experience with the hydrology of the site, vegetation community mapping was finalized during the fall visit.

2.2 Hydrology

Hydrologic indicators were primarily evaluated at the site during the mid-season visit, but additional notes were also taken during the fall visit. Wetland hydrology indicators were recorded using procedures outlined in the Corps of Engineers (COE) Wetlands Delineation Manual (Environmental Laboratory 1987) and hydrology data were recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). If located within 18 inches of the ground surface (soil pit depth for purposes of delineation), groundwater depths were documented on the routine wetland delineation data form at each data point.

All additional hydrologic data were recorded on the mitigation site monitoring form (**Appendix B**). The boundary between wetlands and open water (no rooted vegetation) aquatic habitats was mapped on the aerial photograph and an estimate of the average water depth at this boundary was recorded.

2.3 Vegetation

General dominant species-based vegetation community types (e.g., *Typha latifolia*/*Scirpus acutus*) were delineated on an aerial photograph during the fall visit. Standardized community mapping was not employed as many of these systems are geared towards climax vegetation and may not reflect yearly changes. Estimated percent cover of the dominant species in each community type was listed on the site monitoring form (**Appendix B**).

A 10-foot wide belt transect was established in 2005 (**Figure 2 in Appendix A**). Within the transect belt, percent cover was estimated for each vegetative species for each vegetation community encountered within the “belt” using the following values: + (<1%); 1 (1-5%); 2 (6-10%); 3 (11-20%); 4 (21-50%); and 5 (>50%).

The purpose of the transect is to evaluate changes over time, especially the establishment and increase of hydrophytic vegetation. The transect location was marked on the aerial photo and all data recorded on the mitigation site monitoring form. Transect endpoint locations were recorded with a global positioning system (GPS) unit. Metal fence posts were installed to physically mark the transect ends. Photos of the transect were taken from both ends during the mid-season visit. A comprehensive plant species list for the site was compiled.

Seven woody species were planted at this mitigation site. Planting locations were documented as point data with a GPS unit. Observers recorded the number of dead individuals for each species observed and compared them to known planting numbers.

2.4 Soils

Soils were evaluated during the mid-season visit according to hydric soils determination procedures outlined in the COE 1987 Wetland Delineation Manual. Soil data were recorded for each wetland determination point on the COE Routine Wetland Delineation Data Form (**Appendix B**). The most current terminology used by NRCS was used to describe hydric soils (USDA-NRCS 2006).

2.5 Wetland Delineation

Wetland delineation was conducted during the mid-season visit in accordance with the 1987 COE Wetland Delineation Manual. In July 2008, consultation with the COE (Steinle pers. comm.) confirmed that, where the 1987 manual was used to establish baseline wetland conditions at MDT wetland mitigation sites, it should continue to be applied at such sites for the duration of the monitoring period. Consequently, application of the new *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Great Plains Region* (COE 2008) was not required or undertaken at this site in 2008. Wetland and upland areas within the monitoring area were investigated for the presence of wetland hydrology, hydrophytic vegetation and hydric soils. The indicator status of vegetation was derived from the National List of Plant Species that occur in Wetlands: Northwest (Region 9) (Reed 1988).

The information was recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). The wetland/upland boundary was delineated using a resource grade GPS unit during the fall visit. The wetland/upland boundary in combination with the wetland/open water habitat boundary was used to calculate the wetland area that has developed within the monitoring area.

2.6 Mammals, Reptiles, and Amphibians

Mammal, reptile, and amphibian species observations and other positive indicators of use, such as vocalizations, were recorded on the wetland monitoring form during each visit. Indirect use indicators, including tracks; scat; burrows; eggshells; skins; bones; etc., were also recorded. Observations were recorded as the observer traversed the site while conducting other required activities. Direct sampling methods, such as snap traps, live traps, and pitfall traps, were not implemented. A comprehensive list of observed species was compiled. Observations from past monitoring is compared to this data.

2.7 Birds

Bird observations were recorded during each visit. No formal census plots, spot mapping, point counts, or strip transects were conducted. During the mid-season visit, bird observations were recorded incidental to other monitoring activities. During the fall visit, observations were recorded in compliance with the Bird Survey Protocol in **Appendix E**. During both visits, observations were categorized by species, activity code, and general habitat association (**Bird Survey Field Data Sheets in Appendix B**).

2.8 Macroinvertebrates

One macroinvertebrate sample was collected during the mid-season site visit and data recorded on the wetland mitigation monitoring form. Macroinvertebrate sampling procedures and analysis are included in **Appendix F**. The approximate location of this sample point, within emergent marsh habitat in the northeast portion of the site, is shown on **Figure 2 in Appendix A**. The sample was preserved as outlined in the sampling procedure and sent to a laboratory for analysis. The sample point in 2008 and 2007 differs from the sample points in 2005 and 2006. The 2005 sample macroinvertebrate sample point was taken in one of the ponds that had been established for several years. This information helps evaluators to understand the site's potential. The sample point taken in 2006 was in one of the new shallow pond/emergent marsh areas and represents the early stages of ecosystem evolution at the Wagner Marsh. The 2006 sample point was dry during the 2007 mid-season visit, therefore a new site was selected that has had water during all three years of monitoring, and therefore, presumably, will be able to be sampled in subsequent years. The 2008/2007 sampling site is similar to the 2006 sample site in that the site was also newly constructed in 2005.

2.9 Functional Assessment

Since 2001, a functional assessment for each delineated wetland was conducted using the 1999 MDT Montana Wetland Assessment Method (Berglund 1999). In 2008 the 2008 MDT Montana Wetland Assessment Method (Berglund and McEldowney 2008) was applied. Field data necessary for this assessment were generally collected during the mid-season site visit. The remainder of the functional assessment was completed in the office. For each wetland or group of wetlands (that share similar functions and values) a Functional Assessment form was completed (**Appendix B**)

2.10 Photographs

Photographs were taken during the mid-season visit showing the current land use surrounding the site, the upland buffer, the monitored area, macroinvertebrate sampling location, and the vegetation transect (**Appendix C**). Each photograph point location was recorded with a GPS in 2005. The approximate location of photo points is shown on **Figure 2** in **Appendix A**. All photographs were taken using a digital camera, with no optical zoom used. A description and compass direction for each photograph was recorded on the wetland monitoring form.

2.11 GPS Data

During the 2005 monitoring season, data were collected with a Garmin 12CT GPS unit at the vegetation transect beginning and ending locations, at all photograph locations, wetland sample points, and at aerial photograph reference points. These data were not re-collected in 2008. A resource-grade Magellan MobileMapper GPS unit was used to map wetland boundaries in 2007. Procedures for GPS mapping and aerial photograph referencing are in **Appendix E**.

2.12 Maintenance Needs

Where encountered, current or potential future problems were documented and conveyed to MDT.

3.0 RESULTS

3.1 Hydrology

Groundwater has been the primary hydrologic component of Wagner Marsh, with precipitation playing a minor role in the overall water budget. Recently, excavation of the gravel pit on the west side of S. 56th Street has altered groundwater routing, which without corrective measures would likely have caused the dewatering of the Wagner Marsh mitigation site. However, MDT has developed an agreement with the gravel mining company and they now pump water from their gravel pits into the mitigation site. This has resulted in an overall increase in water within the mitigation area.


The closest weather station to the wetland monitoring area is Laurel, MT station #244894, but it was closed in 1994. According to the Western Regional Climate Center (WRCC) (2008), mean annual precipitation at this station is approximately 14.61 inches; with the majority of precipitation occurring in April, May, June, and September. The closest *active* weather station is Billings WSO (Sta. #240807). Excluding the month of June, the precipitation total through October 2008 at the Billings weather station was 12.36 inches (WRCC 2008). No data were collected during the month of June 2008. If the average value for June is used, the total amount of precipitation is 14.44 inches, which is 1.47 inches more than the average of approximately 12.97 inches for this time of year. Annual evaporation pan rates are estimated to be approximately 41.27 inches at the Huntley Experiment Station (WRCC 2008), almost three times the yearly precipitation rate.

Inundation was present at all wetland cells within the monitoring area during the mid-season visit. Open water areas are shown on **Figure 3 (Appendix A)**.

MDT has contracted with the U.S. Geologic Survey (USGS) to monitor groundwater wells at the Wagner Marsh since 1998. **Chart 1** depicts groundwater fluctuations for one well and provides an example of groundwater fluctuations in the area. Based on the dates of recorded high and low water levels, it is clear that groundwater levels are typically highest in August and September and often the lowest in the spring. Water levels are presumably linked to agricultural use and irrigation periods. This hydroperiod is the opposite of most wetlands in Montana and may hinder the establishment of hydrophytic plant species that have evolved under a more natural hydrologic regime (i.e., wettest in spring, driest in late summer/early fall). The graph also shows that groundwater levels dropped in 2005 when the mitigation site was constructed. It is unclear if the drop in groundwater levels is due to the construction of the mitigation site, groundwater dewatering at nearby gravel pit operations, an increase in evaporation, a change in irrigation practices, drought, or a combination of these factors. From 2005 through 2008 there is a noticeable downward trend for groundwater elevations at this well. This trend is assumed to be linked to the gravel pit located on the west side of 56th St. Supplemental water from that gravel pit is being pumped into the Wagner Mitigation site to supplement water availability at the site. Though a different source of water than what existed prior to construction of the new gravel pit in 2007, this supplemental water appears to be working well because surface water levels were high in 2008 and all wetland cells contained water.

Of the 39 acres in the monitoring area, approximately 42 percent was inundated (**Figure 3 in Appendix A**), with an average depth of 1 foot and a range of depths from zero to an estimated five feet. As in previous years, the pond located immediately south of the crescent shaped pond on the west side of the site appeared to have the greatest maximum depths; approximately 5 feet deep.

Chart 1: An example of the variation in groundwater levels at the Wagner Marsh Wetland Mitigation Site (USGS Well #5).



NOTE: The line connecting points is for display purposes only and are included to show general trends in groundwater levels. It should be understood that groundwater levels can vary substantially between monitoring dates.

3.2 Vegetation

Vegetation species identified on the site are presented in **Table 1** and on the **Monitoring Form (Appendix B)**. Construction of the site was completed in June 2005. In 2008 a total of seven community types were documented at the site, of which five are vegetated wetland community types. These wetland community types were identified and mapped (**Figure 3 in Appendix A**) as: Type 2 - *Salix exigua*-*Eleagnus angustifolia*/*Carex lanuginosa* (*Salix* type), Type 3 - *Eleocharis palustris*-*Typha* sp./*Mixed graminoids* (*Eleocharis*-*Typha* type), Type 10 - *Mixed graminoids*, Type 11 - *Phalaris arundinaceae*, and Type 12 - *Scirpus acutus* (*Scirpus* type). Dominant species within each of these communities are listed on the **Monitoring Form (Appendix B)**. The mixed graminoid and *Phalaris arundinaceae* types occur as wetland fringes around previously existing ponds on the west and northwest sides of the site (**Figure 3 in Appendix A**) and evolved from the *Polypogon* and *Polygonum lapathifolium* types from previous years.

Table 1: 2005 – 2008 vegetation species list for the Wagner Marsh Wetland Mitigation Site.

Scientific Name*	1988 Region 9 (Northwest) Wetland Indicator	Scientific Name*	1988 Region 9 (Northwest) Wetland Indicator
<i>Agropyron cristatum</i>	--	<i>Medicago lupulina</i>	FAC
<i>Agropyron repens</i>	FACU	<i>Medicago sativa</i>	--
<i>Agropyron smithii</i>	FACU	<i>Melilotus officinalis</i>	FACU
<i>Agropyron</i> spp.	--	<i>Nepeta cataria</i>	FAC
<i>Agrostis alba</i>	FACW	<i>Oenothera biennis</i>	FACU
<i>Alyssum</i> spp.	--	<i>Onopordum acanthium</i>	--
<i>Asclepias</i> spp.	--	<i>Panicum capillare</i>	FAC
<i>Aster brachyactis</i>	FACW	<i>Polygonum aviculare</i>	FACW-
<i>Aster</i> spp. (white)	--	<i>Polygonum lapathifolium</i>	FACW+
<i>Beckmannia syzigachne</i>	OBL	<i>Polygonum persicaria</i>	FACW
Brassicaceae (mustard)	--	<i>Polypogon monspeliensis</i>	FACW
<i>Bromus inermis</i>	--	<i>Populus deltoides</i>	FAC
<i>Bromus japonicus</i>	FACU	<i>Potentilla anserina</i>	OBL
<i>Carex lanuginosa</i>	OBL	<i>Prunus virginiana</i> (planted)	FACU
<i>Carex nebrascensis</i>	OBL	<i>Ribes aureum</i> (planted)	FAC+
<i>Carex</i> spp.	--	<i>Rosa woodsii</i> (planted)	FACU
<i>Centaurea maculosa</i>	--	<i>Rumex crispus</i>	FACW
<i>Chenopodium album</i>	FAC	<i>Salix amygdaloides</i>	FACW
<i>Cirsium arvense</i>	FACU+	<i>Salix exigua</i>	OBL
<i>Convolvulus arvensis</i>	--	<i>Salsola iberica</i>	--
<i>Conyza canadensis</i>	FACU	<i>Scirpus acutus</i>	OBL
<i>Echinochloa muricata</i>	FACW	<i>Scirpus maritimus</i>	OBL
<i>Eleagnus angustifolia</i>	FAC	<i>Scirpus pungens</i>	OBL
<i>Eleagnus commutata</i> (planted)	NI	<i>Shepherdia argentea</i> (planted)	--
<i>Eleocharis palustris</i>	OBL	<i>Sisymbrium altissimum</i>	FACU-
<i>Epilobium ciliatum</i>	FACW-	<i>Solidago canadensis</i>	FACU
<i>Erodium cicutarium</i>	--	<i>Sonchus arvensis</i>	FACU+
<i>Festuca pratensis</i>	FACU+	<i>Tamarix ramosissima</i>	FACW
<i>Grindelia squarrosa</i>	FACU	<i>Taraxacum officinale</i>	FACU
<i>Hordeum jubatum</i>	FAC+	<i>Thlaspi arvense</i>	NI
<i>Juncus torreyi</i>	FACW	<i>Tragopogon dubius</i>	--
<i>Juniperus scopulorum</i> (planted)	--	<i>Typha angustifolia</i>	OBL
<i>Lactuca serriola</i>	FACU	<i>Typha latifolia</i>	OBL
<i>Leptochloa fusca</i>	FACW	Unidentified shrub	--
<i>Linum lewisii</i>	--	<i>Verbena bracteata</i>	FACU+
<i>Lotus unifoliolatus</i>	--		

***Bolded plant species** were observed for the first time in 2008.

The *Eleocharis-Typha* type is the most common wetland type on the site and occurs as scattered pockets throughout the mitigation area. With the supplemental water being added to the site from the gravel mine, the *Carex* type that, in 2007, had taken the place of the *Eleocharis-Typha* type in the northwest portion of the site east-adjacent to the *Salix* type reverted back to the *Eleocharis-Typha* type in 2008. The *Echinochloa* type that occurred in the northeastern portion of the site has now developed into the *Eleocharis-Typha* type.

Upland communities are primarily dominated by seeded and/or weedy herbaceous species including, smooth brome (*Bromus inermis*), crested wheatgrass (*Agropyron cristatum*), western wheatgrass (*Agropyron smithii*), meadow fescue (*Festuca pratensis*), Japanese brome (*Bromus japonicus*), quackgrass (*Agropyron repens*), field bindweed (*Convolvulus arvensis*), lambsquarters (*Chenopodium album*), and spotted knapweed (*Centaurea maculosa*). Weed control efforts primarily for knapweed and Canada thistle (*Cirsium arvense*) were implemented in upland areas in 2007 and 2008 and have been largely successful in controlling these weeds.

Vegetation community data were recorded from a transect (**Monitoring Forms in Appendix B**) and summarized in **Table 2**. The types of communities and their relative extent did not change substantially from 2006 to 2007 (**Charts 2 and 3**). In 2007 the number of hydrophytic and upland plant species was consistent with 2006 results (**Table 2**). The overall percent cover in 2008 was about 30%, the same as in 2005. The percent cover decreased in 2008 compared to 2007 and is likely primarily due to higher water levels causing a shift in species, but grazing by waterfowl cannot be discounted as affecting percent cover as well.

Table 2: 2005 – 2008 vegetation transect data summary.

Monitoring Year	2005	2006	2007	2008
Transect Length (feet)	530	530	530	530
# Vegetation Community Transitions along Transect	5	5	5	4
# Vegetation Communities along Transect	4	3	3	2
# Hydrophytic Vegetation Communities along Transect	2	2	1	1
Total Vegetative Species	31	31	31	19
Total Hydrophytic Species	13	15	15	16
Total Upland Species	18	16	16	3
Estimated % Total Vegetative Cover	30	45	55	30
% Transect Length Comprised of Hydrophytic Vegetation Communities	67	62	65	70
% Transect Length Comprised of Upland Vegetation Communities	7	6	5	0
% Transect Length Comprised of Unvegetated Open Water	4	31	30	30
% Transect Length Comprised of Bare Substrate	22	0	0	0

Chart 2: Transect maps showing vegetation types from the start of transect (0 feet) to the end of transect (530 feet) for each year monitored.





Chart 3: Length of vegetation communities within Transect 1 for each year monitored.



A total of 550 woody plantings were installed as part of the overall revegetation plan for the site. Observed mortality of planted woody vegetation species is summarized in **Table 3**. As of August 8, 2008, the overall survival rate is estimated at 45 percent, with a total of 305 individuals observed to be dead. This is down from the 92 percent survival rate reported in 2005, the 64 percent survival rate in 2006 and the 57 percent survival rate in 2007. Juniper plantings continue to do well; mortality of the other species is likely due to a lack of available water during the summer months.

Table 3: 2008 observed mortality of planted woody species for the Wagner Marsh Wetland Mitigation Site.

Plant Species	Number Originally Planted	Number Observed Alive	Number Observed Dead	Cause of Mortality
<i>Eleagnus commutata</i>	50	19	31	Mortality assumed to be due to lack of water.
<i>Juniperus scopulorum</i>	50	47	3	Mortality assumed to be due to lack of water.
<i>Populus deltoides</i>	50	27	23	Mortality assumed to be due to lack of water.
<i>Prunus virginiana</i>	100	63	37	Mortality assumed to be due to lack of water.
<i>Ribes aureum</i>	100	44	56	Mortality assumed to be due to lack of water.
<i>Rosa woodsii</i>	100	42	58	Mortality assumed to be due to lack of water.
<i>Shepherdia argentea</i>	100	3	97	Mortality assumed to be due to lack of water.
TOTAL	550	245	305	

3.3 Soils

Since the site was excavated and graded in Spring 2005, soils are highly disturbed throughout the site. Soils sampled in wetland areas were comprised of silty clay. The primary matrix color of the upper horizon was N 2.5/0. Much of the site was saturated.

3.4 Wetland Delineation

Delineated wetland boundaries are illustrated on **Figure 3 (Appendix A)**. Completed COE Wetland Delineation Forms are included in **Appendix B**. Soils, vegetation, and hydrology were discussed in preceding sections. Total aquatic habitat on the site in 2008 was 16.19 acres (**Figure 3 in Appendix A**). Wetlands comprised 7.38 acres of the 16.19-acre total, consisting of 2.12 acres of wetland originally created on the site by MDT plus 5.26 acres that have developed to date since implementation of the formal mitigation design in 2005. This is a slight decrease of 0.12 acre from the wetland extent in 2007.

Open water comprised 8.81 acres of the 16.19-acre total, an increase of 3.01 acre from the 5.8 acres of open water reported in 2007. Assuming water levels remain fairly constant, the shallow open water habitat observed in 2008 is expected to continue to become vegetated with emergent hydrophytic species over time. A 50-foot wetland buffer around wetlands on the site is approximately 5.19 acres in size. Credits that have developed to date are discussed below in **Section 3.10**.

3.5 Wildlife

Though only constructed in 2005, the wetland complex created on the site provides habitat for several wildlife species. One amphibian, six mammal and 21 bird species were observed at the site during 2008 monitoring (**Table 4**). The habitat value of the site is expected to increase as vegetation continues to establish and diversify. The site continues to be a favored resting/foraging area for birds, and especially waterfowl, with Mallards, Canada Geese, and Red-winged Blackbirds the most numerous bird species observed during the fall bird monitoring event (**Appendix B**).

Table 4: Fish and wildlife species observed at the Wagner Marsh Wetland Mitigation Site during 2005 to 2008.

AMPHIBIAN	
Western chorus frog (<i>Pseudacris triseriata</i>)	Woodhouse's toad (<i>Bufo woodhousii</i>)
REPTILE	
Western garter snake (<i>Thamnophis elegans</i>)	
BIRD	
American Black Duck (<i>Anas rubripes</i>)(?)	Killdeer (<i>Charadrius vociferous</i>)
American Coot (<i>Fulica americana</i>)	Lesser Scaup (<i>Aythya affinis</i>) ¹
American Crow (<i>Corvus brachyrhynchos</i>) ¹	Lesser Yellowlegs (<i>Tringa flavipes</i>) ¹
American Goldfinch (<i>Carduelis tristis</i>)	Mallard (<i>Anas platyrhynchos</i>)
American Robin (<i>Turdus migratorius</i>) ¹	Mourning Dove (<i>Zenaidura macroura</i>)
American Wigeon (<i>Anas americana</i>) ¹	Northern Harrier (<i>Circus cyaneus</i>)
Barn Swallow (<i>Hirundo rustica</i>)	Northern Pintail (<i>Anas acuta</i>) ¹
Black-billed Magpie (<i>Pica hudsonia</i>) ¹	Northern Shoveler (<i>Anas clypeata</i>) ¹
Blue-winged Teal (<i>Anas discors</i>) ¹	Pied-billed Grebe (<i>Podilymbus podiceps</i>)
California Gull (<i>Larus californicus</i>)	Red-tailed Hawk (<i>Buteo jamaicensis</i>)
Canada Goose (<i>Branta canadensis</i>)	Red-winged Blackbird (<i>Agelaius phoeniceus</i>)
Cinnamon Teal (<i>Anas cyanoptera</i>) ¹	Redhead (<i>Aythya americana</i>) ¹
Cliff Swallow (<i>Hirundo pyrrhonota</i>)	Ring-necked Pheasant (<i>Phasianus colchicus</i>)
Common Snipe (<i>Gallinago gallinago</i>)	Rock Dove (<i>Columba livia</i>) ¹
Eastern Kingbird (<i>Tyrannus tyrannus</i>)	Sandhill Crane (<i>Grus canadensis</i>)
Gadwall (<i>Anas strepera</i>)	Song Sparrow (<i>Melospiza melodia</i>)
Grasshopper Sparrow (<i>Ammodramus savannarum</i>)	Spotted Sandpiper (<i>Actitis macularia</i>) ¹
Great Blue Heron (<i>Ardea herodias</i>)	Tree Swallow (<i>Tachycineta bicolor</i>) ¹
Greater Yellowlegs (<i>Tringa melanoleuca</i>)	Vesper Sparrow (<i>Pooecetes gramineus</i>) ¹
Green-winged Teal (<i>Anas crecca</i>) ¹	Western Meadowlark (<i>Sturnella neglecta</i>) ¹
	Wilson's Phalarope (<i>Phalaropus tricolor</i>) ¹
MAMMAL	
Black-tailed jackrabbit (<i>Lepus californicus</i>) ¹	Red Fox (<i>Vulpes vulpes</i>) ¹
Eastern cottontail (<i>Sylvilagus floridanus</i>)	Vole (unidentified species)
Mule deer (<i>Odocoileus hemionus</i>)	White-tailed jackrabbit (<i>Lepus townsendi</i>)
Muskrat (<i>Ondatra zibethicus</i>) ¹	White-tailed deer (<i>Odocoileus virginiana</i>)
Raccoon (<i>Procyon lotor</i>)	

¹ Species observed by MDT staff


Bolded species represent those observed in 2008.

3.6 Macroinvertebrates

In 2005 macroinvertebrates were sampled within the emergent marsh complex on the east side of the site on the northern end of the crescent-shaped pond (**Figure 2 in Appendix A**). This site represented an area that had already been established prior to the construction of the mitigation site, and to some degree represented the site's potential after several years of establishment. That site had high taxa richness and an unusually high number of notonectid hemipterans (Bollman 2005). To better understand how the macroinvertebrate community changes over time, the sampling location was moved in 2006 to a portion of the mitigation site that was constructed in 2005. This site was much less developed in terms of the macroinvertebrate assemblage and was dominated by biting flies (Bollman 2006). The sample site was moved again in 2007 due to the 2006 sample site being dry during the mid-season visit; it had not been dry in the two preceding years. The new sample site is located in an area that was constructed in 2005, but appears to have a more stable water regime than the 2006 sample site. For this reason future changes in macroinvertebrate sample site locations is not expected. The 2008 sampling results are provided in **Appendix F** and were summarized by Rhithron Associates, Inc. in the italicized section below (Bollman 2008).

*Although the assessment score remained about the same, there were some significant changes in the invertebrate assemblage at this site in 2008 compared to 2007. The abundance of invertebrates increased, while the taxonomic richness remained low. While biting midges dominated the sample taken in 2007, this year mayflies (especially *Caenis* sp.) were the dominant organism. These findings, as well as the appearance of odonate nymphs (*Libellulidae* and *Enallagma* sp.) suggest that macrophytes were well-established in 2008. The thermal preference of the biota was calculated at 20.5°C, implying very warm water. Predators were collected, but the major functional components of the assemblage were gatherers and scrapers (*Physa* sp.), indicating some increase in functional complexity since 2007. Air-breathers and hemoglobin-bearers were much less prevalent in the invertebrate assemblage, suggesting an improvement in oxygenation since the previous year.*

Chart 4: Macroinvertebrate bioassessment scores for the Wagner Marsh Wetland Mitigation Site from 2005 to 2008.



NOTE: Direct comparisons can only be made between the 2007 and 2008 scores.

3.7 Functional Assessment

Completed functional assessment forms are presented in **Appendix B** and results are summarized in **Table 5**. Pre-construction conditions through 2007 conditions were assessed using the 1999 MDT MWAM, and 2008 conditions were assessed using the 2008 MDT MWAM. Although direct comparisons cannot be made, general trends in wetland development can still be determined (**Table 5**).

Table 5: Summary of 2001 and 2005 through 2008 wetland function/value ratings and functional points at the Wagner Marsh Wetland Mitigation Site.

Function and Value Parameters from the MDT Montana Wetland Assessment Method ¹	2001 ¹ Baseline Assessment	2005 ¹	2006 ¹	2007 ¹	2008 ²
Listed/Proposed T&E Species Habitat	Low (0.5)	Low (0.5)	Low (0.5)	Low (0.0)	Low (0.0)
MNHP Species Habitat	Low (0.2)	Low (0.2)	Low (0.2)	Low (0.2)	Mod (0.6)
General Wildlife Habitat	Low (0.3)	Mod (0.7)	Mod (0.7)	Mod (0.7)	High (0.9)
General Fish/Aquatic Habitat	N/A	N/A	N/A	N/A	N/A
Flood Attenuation	N/A	N/A	N/A	N/A	N/A
Short and Long Term Surface Water Storage	Mod (0.6)	High (1.0)	High (1.0)	High (1.0)	High (1.0)
Sediment/Nutrient/Toxicant Removal	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
Sediment/Shoreline Stabilization	N/A	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
Production Export/Food Chain Support	Mod (0.6)	High (0.8)	High (0.9)	High (0.9)	High (0.8)
Groundwater Discharge/Recharge	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)
Uniqueness	Low (0.2)	Mod (0.5)	Mod (0.5)	Mod (0.5)	Mod (0.5)
Recreation/Education Potential	Low (0.2)	Low (0.1)	Mod (0.5)	High (1.0)	Mod (0.1)
Actual Points / Possible Points	4.3/9	5.8/10	6.7/10	6.7/10	6.3/9
% of Possible Score Achieved	48%	58%	67%	67%	70%
Overall Category	IV	III	II	II	II
Total Acreage of Assessed Aquatic Habitat within AA Boundaries	3.87	11.84	11.49	13.30	16.19
Functional Units (acreage x actual points)	16.64	68.70	77.00	89.11	102.00
Net Acreage Gain	N/A	7.84	7.62	9.43	12.32
Net Functional Unit Gain	N/A	52.1	60.36-2001 8.30-2005	72.47-2001 12.11-2006	85.36-2001 12.89-2007

¹ Assessed using the 1999 MDT Montana Wetland Assessment Method.

² Assessed using the 2008 MDT Montana Wetland Assessment Method. The completed form is in **Appendix B**.

The created wetlands at Wagner Marsh were ranked as Category II wetlands in 2006, 2007 and 2008, as compared to Category IV in 2001. Functions that increased substantially over 2001 baseline conditions include MNHP species habitat, general wildlife habitat, short and long term surface water storage, production export, uniqueness, and recreation/education potential. The pre-project site provided about 16.6 functional units within the monitoring area, and the post-project site currently provides about 102 functional units, for a conservative gain of 85 functional units.

3.8 Photographs

Representative photographs taken from photo-points and transect ends are provided in **Appendix C**.

3.9 Maintenance Needs/Recommendations

A few salt cedar saplings were observed and removed during monitoring in 2008, others were shown to the weed sprayer on October 7, 2008. The presence of salt cedar on the site should continue to be monitored and individuals removed when encountered, but overall the threat of salt cedar invasion appears to be low.

In 2006 it was noted that spotted knapweed was well established on the berm on the east side of the site, and in upland communities and that Canada thistle was prevalent in the cattail area in the northwestern portion of the site. During mid-season visits in 2007 and again in 2008 it was noted that a comprehensive weed spraying program had been implemented at the site. This effort made significant progress toward eradicating these species from the site, however, spraying in subsequent years is still needed to fully address the severity of the problem.

Water levels continue to be variable, however it appears that the supplemental water being pumped into the site from the gravel mine west of the site is helping to maintain a somewhat less variable water regime when compared to previous years. Opportunities to create a more natural water regime on the site should be explored, as it could help the establishment and persistence of emergent vegetation on the site. For example water levels might be able to be maintained relatively high until the middle of July and then slowly decreased through the end of August, and then water levels might be able to be increased slightly in September and October.

The plant protectors used when planting woody species have started to greatly affect the growth of many of these plants. It is suggested that the plant protectors be removed.

3.10 Current Credit Summary

Based on documentation provided by MDT, approximately 2.12 acres of wetland and 1.75 acres of open water (3.87 acres total of aquatic habitat) were incidentally created on the site via pit excavation prior to formal mitigation project implementation in 2005 (*note: the April 1, 2004 MDT correspondence to the COE indicated 3.87 acres of wetlands and 1.75 acres of open water, which appears to have inadvertently double-counted the open water, adding 1.75 acres to the 2.12 wetland acres [see map in Appendix D]; the July 23, 2004 COE correspondence to MDT correctly indicated 2.12 acres of wetlands, but inadvertently provided an incorrect 1.92-acre figure for the actual 1.75 acres of open water*).

MDT is receiving credit for these wetlands as they were originally created in association with the 2000-2001 Shiloh Road interchange project and protected from disturbance by MDT (Urban pers. comm.). As of 2008, a total of approximately 16.19 acres of open water and wetland habitat (including the original 3.87 acres) occur within the monitoring area (**Table 6**). This is an increase of approximately 2.89 acres from 2007 totals (13.3 acres).

Of the 16.19-acre 2008 total, approximately 8.81 acres are currently open water habitat and the remaining 7.38 acres are vegetated wetland areas. Due to the variability in water levels at Wagner Marsh, it is unclear how much of the open water habitat will evolve into emergent wetland areas. Much of the 'disturbed-moist' vegetation type of previous monitoring years was

classified as emergent wetlands or open water in 2008. A 50 foot wetland buffer around wetlands on the site comprises approximately 5.19 acres (**Table 6**).

Table 6: Summary of open water and wetland acreages at the Wagner Marsh Wetland Mitigation Site for 2001, 2005, 2006, 2007, and 2008.

Year	Open Water (acre)	Wetland (acre)	Total Aquatic Habitat
2001 (pre-mitigation creation)	1.75	2.12	3.87
2005 (post-construction)	7.88	3.96	11.84
2006 (ongoing establishment)	4.96	6.53	11.49
2007 (ongoing establishment)	5.80	7.50	13.30
2008 (ongoing establishment)	8.81	7.38	16.19

The Corps of Engineers will determine which crediting ratios are applicable to the site. However, using the credit ratios listed, **Table 7** summarizes compensatory mitigation credits developed to date at the Wagner Marsh. Using these assumed credit ratios for wetlands, open water, and upland buffer, approximately 10.16 acres of credit are currently available, a decrease of approximately 0.14. This decrease is primarily attributed to water levels being higher and flooding out some of the wetland areas. However, if the water levels remain relatively constant, there is potential for a greater extent of emergent wetlands to establish at Wagner Marsh than in previous years.

Table 7: 2008 mitigation credit summary for the Wagner Marsh Wetland Mitigation Site.

Credit Category	Acre	Assumed Credit Ratio ¹	Credit ¹
Total Scrub/Shrub and Emergent Wetland	7.38	1:1	7.38
Total Open water	8.81	20% of wetland acreage ²	1.48
50-foot wide upland buffer	5.19	4:1	1.30
TOTAL	16.68		10.16

¹ The Corps of Engineers is the regulatory authority and will determine the actual mitigation ratios.

² According to July 23, 2004 correspondence from the Corps to MDT, "credit for open water will be limited to no more than 20% of the amount of actual wetland that develops at the site. For example, if 20 acres of wetland develops, up to 4 acres of additional acres of open water credit could be used as wetland mitigation credit."

The pre-project site provided about 16.6 functional units within the monitoring area, and in 2008 the mitigation site provides about 102 functional units, for a conservative gain of 85 functional units.

4.0 REFERENCES

- Berglund, J. 1999. *MDT Montana Wetland Assessment Method*. Prepared for: Montana Department of Transportation and Morrison-Maierle, Inc. Western EcoTech. Helena, Montana. 18 pp.
- Berglund, J., and R. McEldowney. 2008. *Montana Wetland Assessment Method*. Prepared for Montana Department of Transportation. Post, Buckley, Schuh and Jernigan (PBS&J). Helena, Montana. 42 p.
- Bollman, W. 2008. MDT Mitigated Wetland Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001-2008. Rhithron Associates, Inc. Missoula, Montana.
- Bollman, W. 2007. MDT Mitigated Wetland Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001-2007. Rhithron Associates, Inc. Missoula, Montana.
- Bollman, W. 2006. MDT Mitigated Wetland Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001-2006. Rhithron Associates, Inc. Missoula, Montana.
- Bollman, W. 2005. MDT Mitigated Wetland Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001-2005. Rhithron Associates, Inc. Missoula, Montana.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. US Army Corps of Engineers. Washington, DC.
- Reed, P.B. 1988. *National list of plant species that occur in wetlands: North West (Region 9)*. Biological Report 88(26.9), May 1988. U.S. Fish & Wildlife Service. Washington, D.C.
- Steinle, A. 2008. Montana Program Manager, U.S. Army Corps of Engineers, Helena, Montana. July 14th telephone conversation.
- U.S. Army Corps of Engineers (COE). 2008. *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region*, ed. J. S. Wakeley, R. W. Lichvar, and C. V. Noble. ERDC/EL TR-08-13. U.S. Army Engineer Research and Development Center, Vicksburg, Missouri.
- United States Department of Agriculture, Natural Resources Conservation Service (USDA-NRCS). 2006. *Field Indicators of Hydric Soils in the United States*, Version 6.0. G.W. Hurt and L.M. Vasilas (eds.). USDA-NRCS, in cooperation with the National Technical Committee for Hydric Soils.
- Western Regional Climate Center (WRCC). 2008. Evaporation pan data; precipitation data for Laurel, MT and Billings WSO, Montana. Obtained on November 22nd from the world wide web at: <http://www.wrcc.dri.edu/CLIMATEDATA.html>
- Western Regional Climate Center (WRCC). 2007. Monthly average pan evaporation rates for Montana. Huntley Experiment Station. Period of record 1911-2005. Obtained in Nov from the world wide web at: <http://www.wrcc.dri.edu/htmlfiles/westevap.final.html#MONTANA>

Appendix A

FIGURES 2 & 3

*MDT Wetland Mitigation Monitoring
Wagner Marsh
Billings, Montana*





Figure 3 Mapped Site Features 2008



Appendix B

2008 WETLAND MITIGATION SITE MONITORING FORMS

2008 BIRD SURVEY FORMS

2008 COE WETLAND DELINEATION FORMS

2008 FUNCTIONAL ASSESSMENT FORMS

MDT Wetland Mitigation Monitoring

Wagner Marsh

Billings, Montana

PBS&J / MDT WETLAND MITIGATION SITE MONITORING FORM

Project Name: **Wagner Marsh** Project Number: _____
Assessment Date: **August 8, 2008** Person(s) conducting the assessment: **R. McEldowney**
Location: _____ MDT District: **Billings** Milepost: **NA**
Legal Description: T **1S** R **25E** Section **28**
Weather Conditions: **Clear, calm, 70-95 deg F** Time of Day: **9 to 4 pm**
Initial Evaluation Date: **August 1, 2005** Monitoring Year: **4** # Visits in Year: **2**
Size of evaluation area: **39 acres** Land use surrounding wetland: **Rural/agricultural mostly, gravel pit being excavated immediately west of S. 56th St. West**

HYDROLOGY

Surface Water Source: **Groundwater and overland flow**
Inundation: **Present** Average Depth: 1 **feet** Range of Depths: **0-5 ft**
Percent of assessment area under inundation: **40%**
Depth at emergent vegetation-open water boundary: **Varies - 0 to 2 feet**
If assessment area is not inundated then are the soils saturated within 12 inches of surface: **Yes**
Other evidence of hydrology on the site (ex. – drift lines, erosion, stained vegetation, etc.):
Dried algal mats

Groundwater Monitoring Wells: **Present - monitored on 8/8/08**

Record depth of water below ground surface (in feet):

Well Number	Depth	Well Number	Depth	Well Number	Depth
8/8/08 #1	2.15 ft				
8/8/08 #2	Locked				
8/8/08 #3	5 ft				

Additional Activities Checklist:

- ☒ Map emergent vegetation-open water boundary on aerial photograph.
- ☒ Observe extent of surface water during each site visit and look for evidence of past surface water elevations (drift lines, erosion, vegetation staining, etc.)
- ☒ Use GPS to survey groundwater monitoring well locations, if present.

COMMENTS / PROBLEMS:

VEGETATION COMMUNITIES

Community Number: **1** Community Title (main spp): **Open water/aquatic bed**

Dominant Species	% Cover	Dominant Species	% Cover
Aquatic bed	5 = > 50%		

Comments / Problems: **Shallow ponds less than 5 feet deep that either contain submergent vegetation or are currently inundated but sparsely vegetated due to the relatively recent (2005) construction of the project and the dynamic fluctuations of water levels. Over time it is expected that some of these areas will become palustrine emergent wetlands. In some locations scattered individuals of emergent species occur.**

Community Number: **2** Community Title (main spp): **Salix exigua-Eleagnus angustifolia/Carex lanuginosa**

Dominant Species	% Cover	Dominant Species	% Cover
Eleagnus angustifolia	3 = 11-20%	Typha latifolia	2 = 6-10%
Salix exigua	4 = 21-50%	Carex lanuginosa	4 = 21-50%
Scirpus pungens	3 = 11-20%	Populus deltoides (sap)	2 = 6-10%
Cirsium arvense	3 = 11-20%		

Comments / Problems: **Palustrine scrub-shrub area on the northwest side of the site.**

Community Number: **3** Community Title (main spp): **Eleocharis palustris-Typha latifolia/Mixed graminoids**

Dominant Species	% Cover	Dominant Species	% Cover
Typha latifolia	2 = 6-10%	Eleocharis palustris	5 = > 50%
Typha angustifolia	2 = 6-10%	Juncus torreyi	4 = 21-50%
Scirpus acutus	2 = 6-10%	Agropyron repens	2 = 6-10%
Hordeum jubatum	3 = 11-20%	Polygonum lapathifolium	1 = 1-5%

Comments / Problems: **Palustrine emergent wetland.**

Community Number: **4** Community Title (main spp): **Polypogon monspeliensis**

Dominant Species	% Cover	Dominant Species	% Cover
Polypogon monspeliensis	5 = > 50%		
Typha latifolia	2 = 6-10%		
Scirpus acutus	1 = 1-5%		
Carex lanuginosa	1 = 1-5%		

Comments / Problems: **Not observed in 2007. Evolved into Community Number 10.**

Community Number: **5** Community Title (main spp): **Polygonum lapathifolium/Mixed graminoids**

Dominant Species	% Cover	Dominant Species	% Cover
Polygonum lapathifolium	5 = > 50%	Eleocharis palustris	2 = 6-10%
Juncus torreyi	1 = 1-5%		

Comments / Problems: **Not observed in 2007. Evolved into Community number 11.**

VEGETATION COMMUNITIES (continued)

Community Number: **6** Community Title (main spp): **Upland Grasses**

Dominant Species	% Cover	Dominant Species	% Cover
Festuca pratensis	5 = > 50%		
Bromus inermis	2 = 6-10%		
Bromus japonicus	3 = 11-20%		
Convolvulus arvensis	1 = 1-5%		
Sisymbrium altissimum	2 = 6-10%		

Comments / Problems: **Upland grassland community surrounding the constructed wetland area. The areas between wetland cells are primarily weedy, percent cover varies greatly and bare soil is prevalent throughout. These areas are dominated primarily by Chenopodium alba, Agropyron repens, Melilotus officinale, Convolvulus arvensis, Medicago sativa, Polygonum aviculare, and Agropyron smithii.**

Community Number: **7** Community Title (main spp): **Upland grasses – Drill Seeded**

Dominant Species	% Cover	Dominant Species	% Cover
Medicago sativa	1 = 1-5%		
Agropyron sp.	4 = 21-50%		
Chenopodium album	2 = 6-10%		
Agropyron smithii	1 = 1-5%		
Convolvulus arvensis	2 = 6-10%		
Centaurea maculosa	4 = 21-50%		

Comments / Problems: **Upland area - drill seeded berm on the east side of the site. Spotted knapweed is a problem in this area.**

Community Number: **8** Community Title (main spp): **Disturbed moist**

Dominant Species	% Cover	Dominant Species	% Cover
Melilotus officinale	3 = 11-20%		
Kochia scoparia	1 = 1-5%		
Hordeum jubatum	1 = 1-5%		

Comments / Problems: **Area is primarily bare ground with a variety of weedy and hydrophytic species. This community type may become dominated by hydrophytic vegetation over time if the hydroperiod and required duration of inundation occurs.**

Community Number: **9** Community Title (main spp): **Glyceria grandis**

Dominant Species	% Cover	Dominant Species	% Cover
Glyceria grandis	3 = 11-20%		

Comments / Problems: **Not observed in 2007 or 2008.**

VEGETATION COMMUNITIES (continued)

Community Number: **10** Community Title (main spp): **Mixed Graminoids**

Dominant Species	% Cover	Dominant Species	% Cover
Typha latifolia	1 = 1-5%	Phalaris arundinaceae	2 = 6-10%
Scirpus acutus	1 = 1-5%	Leptochloa fusca	3 = 11-20%
Carex lanuginosa	3 = 11-20%		

Comments / Problems: **New community in 2007. Evolved from Community Number 4.**

Community Number: **11** Community Title (main spp): **Phalaris arundinaceae**

Dominant Species	% Cover	Dominant Species	% Cover
Phalaris arundinaceae	5 = > 50%		
Polygonum lapathifolium	1 = 1-5%		

Comments / Problems: **New community in 2007. Evolved from Community Type 5.**

Community Number: **12** Community Title (main spp): **Scirpus acutus**

Dominant Species	% Cover	Dominant Species	% Cover
Scirpus acutus	5 = > 50%		
Echinochloa muricata	1 = 1-5%		

Comments / Problems: **New community in 2007. Located in the pond in NW portion of site where the PSS wetland is located.**

Community Number: **13** Community Title (main spp): **Carex lanuginosa/Hordeum jubatum**

Dominant Species	% Cover	Dominant Species	% Cover
Carex lanuginosa	4 = 21-50%	Cirsium arvense	2 = 6-10%
Hordeum jubatum	3 = 11-20%		
Phalaris arundinaceae	1 = 1-5%		
Festuca pratensis	1 = 1-5%		

Comments / Problems: **Was a new community in 2007, but not obs. in 2008. Site was classified as Community Type 3 in 2005 and 2006 and reverted to that community type in 2008.**

Community Number: **14** Community Title (main spp): **Echinochloa muricata/Hordeum jubatum**

Dominant Species	% Cover	Dominant Species	% Cover
Echinochloa muricata	4 = 21-50%		
Hordeum jubatum	3 = 11-20%		

Comments / Problems: **New community in 2007, but not observed in 2008.**

Additional Activities Checklist:

☒ Record and map vegetative communities on aerial photograph.

COMPREHENSIVE VEGETATION LIST

Plant Species	Vegetation Community Number (s)	Plant Species	Vegetation Community Number (s)
Asclepias sp.	6	Medicago lupulina	6,7,8
Agrostis alba	2,3	Medicago sativa	6,7,8
Agropyron cristatum	6	Melilotus officinale	8
Agropyron repens	3,6,7,8	Mustard sp.	8
Agropyron smithii	6,7	Nepeta cataria	13
Agropyron sp.	6,7	Onopordum acanthium	7
Alyssum sp.	6	Oenothera biennis	6
Aster brachyactis	3	Panicum capillare	8
Beckmannia syzigachne	8	Phalaris arundinaceae	11,13
Bromus inermis	6,7	Polygonum aviculare	3,6,7,8
Bromus japonicus	6,8	Polygonum lapathifolium	1,3,5,8
Carex lanuginosa	2,4,10,13	Polygonum pensylvanicum	1,3,8
Carex nebrascensis	2,3	Polypogon monspeliensis	4
Carex sp.	3	Populus deltoides	2
Centaurea maculosa	6,7,8	Potentilla anserina	1,8
Chenopodium album	6,7,8	Potentilla recta	6
Cirsium arvense	2,3,6	Rumex crispus	2
Convolvulus arvensis	6,7,8	Salix amygdaloides	2
Conyza canadensis	6,8	Salix exigua	2
Descurainia sophia	8	Salix lutea	3
Echinochloa muricata	1,12,14	Salsola iberica	6,8
Elaeagnus angustifolia	2	Scirpus acutus	3,10,12
Eleocharis palustris	1,3,8	Scirpus maritimus	3
Epilobium ciliatum	2,3,8	Scirpus pungens	2
Erodium cicutarium	6,8	Sisymbrium altissimum	6
Festuca idahoensis	6	Solidago canadensis	6
Festuca pratensis	6,13	Sonchus arvensis	6
Grindellia squarrosa	6	Tamarix ramosissima	2
Glyceria grandis	9	Taraxacum officinale	2,8
Hordeum jubatum	3,6,8,13,14	Thlaspi arvense	2
Juncus bufonius	3	Tragopogon dubius	6
Juncus torreyi	3	Typha angustifolia	3
Kochia scoparia	6	Typha latifolia	3,10
Lactuca serriola	6	Unidentified white aster	6
Leptochloa fusca	10	Unidentified shrub	3
Linum lewisii	6,8	Verbena bracteata	3,8
Lotus unifoliolatus	7		

Comments / Problems: Total number of species observed = 72 (excluding planted shrubs).

Application of herbicides on knapweed and Canada thistle appears to have been conducted in July. This was effective, but must be repeated in subsequent years if these noxious weeds are to be controlled.

PLANTED WOODY VEGETATION SURVIVAL

Plant Species	Number Originally Planted	Live Number Observed	Mortality Causes
Elaeagnus commutata	50	19	Mortality assumed to be due to lack of water.
Juniperus scopulorum	50	47	Mortality assumed to be due to lack of water.
Populus deltoides	50	27	Mortality assumed to be due to lack of water
Prunus virginiana	100	63	Mortality assumed to be due to lack of water
Ribes aureum	100	44	Mortality assumed to be due to lack of water
Rosa woodsii	100	42	Mortality assumed to be due to lack of water
Shepherdia argentea	100	3	Mortality assumed to be due to lack of water

Comments / Problems: The deer protector meshes appear to be inhibiting growth.

WILDLIFE

Birds

Were man-made nesting structures installed? No

If yes, type of structure: _____ How many? _____

Are the nesting structures being used? NA

Do the nesting structures need repairs? NA

Mammals and Herptiles

Mammal and Herptile Species	Number Observed	Indirect Indication of Use			
		Tracks	Scat	Burrows	Other
Mule and whitetail deer	6	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
Raccoon		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
Cottontail	3	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
Chorus Frog	1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	

Additional Activities Checklist:

Yes Macroinvertebrate Sampling (if required)

Comments / Problems:

PHOTOGRAPHS

Using a camera with a 50mm lens and color film take photographs of the following permanent reference points listed in the check list below. Record the direction of the photograph using a compass. When at the site for the first time, establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3 feet above ground. Survey the location with a resource grade GPS and mark the location on the aerial photograph.

Photograph Checklist:

- ☒ One photograph for each of the four cardinal directions surrounding the wetland.
- ☒ At least one photograph showing upland use surrounding the wetland. If more than one upland exists then take additional photographs.
- ☒ At least one photograph showing the buffer surrounding the wetland.
- ☒ One photograph from each end of the vegetation transect, showing the transect.

Location	Photograph Frame #	Photograph Description	Compass Reading (°)
Photopoint A	1	North side of site looking NNE toward WJH bird sanctuary.	22
Photopoint A	2	North side of site looking east across wetland creation area (and transect) toward berm on the east side of site and the canal beyond it.	105
Photopoint A	3	North side of site looking southeast across created wetlands and the south end of the transect.	162
Photopoint A	4	North side of site looking south at central area of the site.	214
Photopoint A	5	North side of site looking at cattail area and south end of the PSS area.	250
Photopoint A	6	North side looking at PSS area in NW corner of site.	310
Photopoint A	7	North side of site looking at pond in NW corner of site.	335
Photopoint B	1	West side of site looking north at the crescent shaped pond in the central portion of the west side of the site.	01
Photopoint B	2	West side of site looking east at a wetland creation area.	74
Photopoint B	3	West side of site looking south at wetland creation areas.	153
Photopoint C	1	South side of site looking NNE at drill seeding on the berm and wetland creation areas to the north.	24
Photopoint C	2	South side of site looking WSW at berm and wetland creation areas at southernmost tip of the site.	243
Photopoint C	3	South side of site looking WNW at wetland creation areas.	294
Photopoint C	4	South side of site looking NNW at wetland creation areas in the south side of the central portion of the site.	343
Photopoint D	1	East side of site looking WSW at beerm and wetland creation areas on the SE side of the site.	241
Photopoint D	2	East side of site looking WNW at the central portion of the site.	293
Photopoint D	3	East side of site looking NW at the transect area in a wetland creation area.	324
Photopoint D	4	East side of site looking north at the drill seeded berm and the north end of the transect.	356
Transect	1	West end of the transect looking ENE.	70
Transect	2	East end of the transect looking WSW.	250

Comments / Problems: Surrounding upland uses (agriculture) and buffer areas are shown in many of the photos listed in the table above.

GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points set at a 5 second recording rate. Record file numbers for site in designated GPS field notebook.

GPS Checklist:

- ☒ Jurisdictional wetland boundary.
- ☒ 4-6 landmarks that are recognizable on the aerial photograph.
- ☒ Start and End points of vegetation transect(s).
- ☒ Photograph reference points.
- ☒ Groundwater monitoring well locations.

Comments / Problems: **All GPS data listed above had been collected in previous years. The wetland boundaries were verified and modified where appropriate on the aerial photo.**

WETLAND DELINEATION

(attach COE delineation forms)

At each site conduct these checklist items:

- ☒ Delineate wetlands according to the 1987 Army COE manual.
- ☒ Delineate wetland – upland boundary onto aerial photograph.
- NA** Survey wetland – upland boundary with a resource grade GPS survey.

Comments / Problems: _____

FUNCTIONAL ASSESSMENT

(Complete and attach full MDT Montana Wetland Assessment Method field forms.)

(Also attach any completed abbreviated field forms, if used)

Comments / Problems: **None.**

MAINTENANCE

Were man-made nesting structure installed at this site? **NA**

If yes, do they need to be repaired? **NA**

If yes, describe the problems below and indicate if any actions were taken to remedy the problems.

Were man-made structures built or installed to impound water or control water flow into or out of the wetland? **NA**

If yes, are the structures working properly and in good working order? **NA**

If no, describe the problems below.

Comments / Problems: _____

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: **Wagner Marsh** Date: **8/8/2008** Examiner: **R. McEldowney (PBS&J)**

Transect Number: **1** Approximate Transect Length: **530 feet** Compass Direction from Start: **70°** Note: _____

Vegetation Type A: Eleocharis palustris-Typha sp./Mixed graminoids	
Length of transect in this type: 62 feet	
Plant Species	Cover
CARLAN	4 = 21-50%
AGRSTO	1 = 1-5%
JUNTOR	+ = < 1%
GLYGRA	+ = < 1%
ELEPAL	5 = > 50%
SCIPUN	+ = < 1%
JUNBAL, CARNEB	1 = 1-5%
TYPLAT	2 = 6-10%
POTANS	+ = < 1%
SCIMIC	+ = < 1%
SALEXI, SALLUT, ELEANG, CIRARV, CONCAN EACH	+ = < 1%
Total Vegetative Cover:	95%

Vegetation Type B: Eleocharis palustris-Typha sp./Mixed graminoids (ELEPAL/SCIPUN)	
Length of transect in this type: 80 feet	
Plant Species	Cover
ELEPAL	5 = > 50%
SCIPUN	3 = 11-20%
SCIMIC	2 = 6-10%
Total Vegetative Cover:	80%

Vegetation Type C: Open water (sparse veg)	
Length of transect in this type: 80 feet	
Plant Species	Cover
TYPANG	1 = 1-5%
ELEPAL	+ = < 1%
Total Vegetative Cover:	3%

Vegetation Type D: Eleocharis palustris-Typha sp./Mixed graminoids	
Length of transect in this type: 70 feet	
Plant Species	Cover
ELEPAL	4 = 21-50%
TYPLAT	1 = 1-5%
TYPANG	+ = < 1%
SCIMIC	+ = < 1%
SCIPUN	+ = < 1%
SCIMAR	+ = < 1%
Total Vegetative Cover:	25%

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: **Wagner Marsh** Date: **August 8, 2008** Examiner: **R. McEldowney (PBS&J)**

Transect Number: **1** Approximate Transect Length: **530 feet** Compass Direction from Start: **70°** Note: _____

Vegetation Type E: Eleocharis palustris-Typha latifolia./Mixed graminoids (ELEPAL/SCIMIC)	
Length of transect in this type: 152 feet	
Plant Species	Cover
ELEPAL	1 = 1-5%
SCIMAR	+ = < 1%
SCIMIC	1 = 1-5%
Total Vegetative Cover:	10%

Vegetation Type F: Open water (sparse veg)	
Length of transect in this type: 80 feet	
Plant Species	Cover
ELEPAL	+ = < 1%
Total Vegetative Cover:	0%

Vegetation Type G: Eleocharis palustris-Typha sp./Mixed graminoids	
Length of transect in this type: 6 feet	
Plant Species	Cover
ELEPAL	4 = 21-50%
SCIMIC	2 = 6-10%
END OF TRANSECT	
Total Vegetative Cover:	30%

Vegetation Type H:	
Length of transect in this type: feet	
Plant Species	Cover
Total Vegetative Cover:	%

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: _____ Date: _____ Examiner: _____

Transect Number: _____ Approximate Transect Length: _____ **feet** Compass Direction from Start: ____° Note: _____

Vegetation Type I:	
Length of transect in this type: _____ feet	
Plant Species	Cover
Total Vegetative Cover:	%

Vegetation Type J:	
Length of transect in this type: _____ feet	
Plant Species	Cover
Total Vegetative Cover:	%

Vegetation Type K:	
Length of transect in this type: _____ feet	
Plant Species	Cover
Total Vegetative Cover:	%

Vegetation Type L:	
Length of transect in this type: _____ feet	
Plant Species	Cover
Total Vegetative Cover:	%

MDT WETLAND MONITORING – VEGETATION TRANSECT

Cover Estimate

+ = < 1% 3 = 11-10%
1 = 1-5% 4 = 21-50%
2 = 6-10% 5 = > 50%

Indicator Class

+ = Obligate
- = Facultative/Wet
0 = Facultative

Source

P = Planted
V = Volunteer

Percent of perimeter developing wetland vegetation (excluding dam/berm structures): 50%

Establish transects perpendicular to the shoreline (or saturated perimeter). The transect should begin in the upland area. Permanently mark this location with a standard metal fencepost. Extend the imaginary transect line towards the center of the wetland, ending at the 3 foot depth (in open water), or at the point where water depths or saturation are maximized. Mark this location with another metal fencepost.

Estimate cover within a 10 foot wide "belt" along the transect length. At a minimum, establish a transect at the windward and leeward sides of the wetland. Remember that the purpose of this sampling is to monitor, not inventory, representative portions of the wetland site.

Comments: **Based on waterlines water levels were deeper earlier in the summer and is likely at least partially the cause of the shift in species composition and percent cover in some portions of the transect.**

BIRD SURVEY – FIELD DATA SHEET

Site: **Wagner** Date: 8/8/2008

Survey Time: **9 am** to **4 PM**

Bird Species	#	Behavior	Habitat	Bird Species	#	Behavior	Habitat
Sandhill cranes	3	F	MA				
Canada Goose	20	F	MA				
Killdeer	12	F	MA MF				
Mallard	22	F	MA				
Grtr Yellowlegs	5	F	MA MF				
Ring-necked Pheasant	1	N	UP				
Above data: 8/8/2008				Above Data:			

BEHAVIOR CODES

BP = One of a breeding pair

BD = Breeding display

F = Foraging

FO = Flyover

L = Loafing

N = Nesting

HABITAT CODES

AB = Aquatic bed

FO = Forested

I = Island

MA = Marsh

MF = Mud Flat

OW = Open Water

SS = Scrub/Shrub

UP = Upland buffer

WM = Wet meadow

US = Unconsolidated shore

Weather: **80+ degrees, clear, breezy**

Notes:

BIRD SURVEY – FIELD DATA SHEET

Site: **Wagner Marsh** Date: **10/7/08**
 Survey Time: **7:30 am** to **9:30 am**

Bird Species	#	Behavior	Habitat	Bird Species	#	Behavior	Habitat
Canada Goose	58	F L FO	MA AB				
Mallard	130	F L FO	OW AB MA MF				
N. Shoveller	4	FO					
Redwing Blackbirds	35	FO	UP				
Ring-necked Pheasant	6	F	UP				
Sandhill Cranes	3	FO					
Common Snipe	3	F FO	MA				
American Coot	1	F					

BEHAVIOR CODES

BP = One of a breeding pair
BD = Breeding display
F = Foraging
FO = Flyover
L = Loafing
N = Nesting

HABITAT CODES

AB = Aquatic bed
FO = Forested
I = Island
MA = Marsh
MF = Mud Flat
OW = Open Water
SS = Scrub/Shrub
UP = Upland buffer
WM = Wet meadow
US = Unconsolidated shore

Weather: **Clear, light breeze, 45 degrees F.**

Notes: **Sunrise occurred at approximately 7:28 am.**

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Wagner Marsh – Billings, MT</u> Applicant/Owner: <u>Montana Department of Transportation</u> Investigator: _____	Date: <u>8/8/2008</u> County: <u>Yellowstone</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: _____ Yes <u>X</u> No Is the site significantly disturbed (Atypical Situation)? <u>X</u> Yes _____ No Is the area a potential Problem Area?: _____ Yes <u>X</u> No (If needed, explain on reverse.)	Community ID: _____ Transect ID: _____ Plot ID: <u>SP-1</u>

Location: 682531 Easting, 5065131 Northing (UTM, WGS84, meters)

VEGETATION

Dominant Plant Species	Stratum	Indicator		Dominant Plant Species	Stratum	Indicator
1 TYPANG	H	OBL		9		
2 ELEPAL	H	OBL		10		
3 AGRREP	H	FACU		11		
4				12		
5				13		
6				14		
7				15		
8				16		

Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). 3/4 = 75%

Remarks: Area was disturbed from construction of mitigation site in 2005. Other species that are present but not dominant include HORJUB, SCIACU, and BECSZY.

HYDROLOGY

<p><u>X</u> Recorded Data (Describe in Remarks): _____ Stream, Lake, or Tide Gauge <u>X</u> Aerial Photographs _____ Other _____ No Recorded Data Available</p> <hr/> <p>Field Observations:</p> <p>Depth of Surface Water: _____ (in.)</p> <p>Depth to Free Water in Pit: _____ (in.)</p> <p>Depth to Saturated Soil: <u>0</u> (in.)</p>	<p>Wetland Hydrology Indicators:</p> <p>Primary Indicators:</p> <p><u>X</u> Inundated <u>X</u> Saturated in Upper 12 Inches _____ Water Marks _____ Drift Lines _____ Sediment Deposits _____ Drainage Patterns in Wetlands</p> <p>Secondary Indicators (2 or more required):</p> <p><u>X</u> Oxidized Root Channels in Upper 12 Inches _____ Water-Stained Leaves _____ Local Soil Survey Data _____ FAC-Neutral Test _____ Other (Explain in Remarks)</p>
<p>Remarks:</p> <p>Have dried algal mats indicating that this area was inundated previously. Soil was saturated to the surface at the time of the site visit.</p>	

SOILS

Map Unit Name (Series and Phase):		Ll- Larim gravelly loam, 15-35% slopes		Drainage Class: <u>Well to excessive</u>	
Taxonomy (Subgroup):		TYPIC USTORTHENTS, SANDY-SKELETAL, MIXED, FRIGID		Field Observations Confirm Mapped Type? <u> </u> Yes <u>X</u> No <u> </u>	

Profile Description:					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0-0.25	1	N 2.5/0			SILTY CLAY
0.25 - 5	2	N 4/0	5YR 4/4	Abundant, prominent	Silty clay

Hydric Soil Indicators:	
<input type="checkbox"/> Histosol <input type="checkbox"/> Histic Epipedon <input checked="" type="checkbox"/> Sulfidic Odor <input checked="" type="checkbox"/> Aquic Moisture Regime <input type="checkbox"/> Reducing Conditions <input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors	<input type="checkbox"/> Concretions <input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils <input type="checkbox"/> Organic Streaking in Sandy Soils <input type="checkbox"/> Listed on Local Hydric Soils List <input type="checkbox"/> Listed on National Hydric Soils List <input type="checkbox"/> Other (Explain in Remarks)

Remarks: Site was disturbed by wetland mitigation construction in 2005. Unable to dig below 5 inches due to cobbles.

WETLAND DETERMINATION

Hydrophytic Vegetation Present? <u>X</u> Yes <u> </u> No Wetland Hydrology Present? <u>X</u> Yes <u> </u> No Hydric Soils Present? <u>X</u> Yes <u> </u> No	Is this Sampling Point Within a Wetland? <u>X</u> Yes <u> </u> No
--	--

Remarks:
 Wetland sample point. Site is dominated by cattails, was saturated to the surface and inundated earlier in the summer, and has several hydric soil indicators.

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: <u>Wagner Marsh – Billings, MT</u> Applicant/Owner: <u>Montana Department of Transportation</u> Investigator: <u>PBS&J (RRM)</u>	Date: <u>8/8/2008</u> County: <u>Yellowstone</u> State: <u>MT</u>
Do Normal Circumstances exist on the site: <u> </u> Yes <u> X </u> No Is the site significantly disturbed (Atypical Situation)? <u> X </u> Yes <u> </u> No Is the area a potential Problem Area?: <u> </u> Yes <u> X </u> No (If needed, explain on reverse.)	Community ID: <u> </u> Transect ID: <u> </u> Plot ID: <u> SP-2 </u>

VEGETATION

Dominant Plant Species	Stratum	Indicator		Dominant Plant Species	Stratum	Indicator
1 AGRREP	H	FACU		9		
2 POTANS	H	OBL		10		
3 CONARV	H	NL		11		
4				12		
5				13		
6				14		
7				15		
8				16		

Percent of Dominant Species that are OBL, FACW, or FAC (excluding FAC-). 1/3 = 33%

Remarks: Area was disturbed from construction of mitigation site in 2005. NL = not listed, assumed to be upland species. Non-dominant species include western wheatgrass, foxtail barley, and birdsfoot trefoil.

HYDROLOGY

<p><u> X </u> Recorded Data (Describe in Remarks):</p> <p style="padding-left: 40px;"><u> </u> Stream, Lake, or Tide Gauge</p> <p style="padding-left: 40px;"><u> X </u> Aerial Photographs</p> <p style="padding-left: 40px;"><u> </u> Other</p> <p><u> </u> No Recorded Data Available</p> <hr/> <p>Field Observations:</p> <p style="padding-left: 40px;">Depth of Surface Water: <u> -- </u> (in.)</p> <p style="padding-left: 40px;">Depth to Free Water in Pit: <u> -- </u> (in.)</p> <p style="padding-left: 40px;">Depth to Saturated Soil: <u> -- </u> (in.)</p>	<p>Wetland Hydrology Indicators:</p> <p>Primary Indicators:</p> <p style="padding-left: 20px;"><u> </u> Inundated</p> <p style="padding-left: 20px;"><u> </u> Saturated in Upper 12 Inches</p> <p style="padding-left: 20px;"><u> </u> Water Marks</p> <p style="padding-left: 20px;"><u> </u> Drift Lines</p> <p style="padding-left: 20px;"><u> </u> Sediment Deposits</p> <p style="padding-left: 20px;"><u> </u> Drainage Patterns in Wetlands</p> <p>Secondary Indicators (2 or more required):</p> <p style="padding-left: 20px;"><u> </u> Oxidized Root Channels in Upper 12 Inches</p> <p style="padding-left: 20px;"><u> </u> Water-Stained Leaves</p> <p style="padding-left: 20px;"><u> </u> Local Soil Survey Data</p> <p style="padding-left: 20px;"><u> </u> FAC-Neutral Test</p> <p style="padding-left: 20px;"><u> </u> Other (Explain in Remarks)</p>
<p>Remarks:</p> <p>Site appears to have been inundated briefly earlier in the summer, but does not have strong indications of wetland hydrology.</p>	

SOILS

Map Unit Name		Le- Larim Loam, 0-4% slopes		Drainage Class: <u>Well to excessive</u>	
(Series and Phase):				Field Observations	
Taxonomy (Subgroup):		<u>TYPIC ARGIBOROLLS, LOAMY-SKELETAL, MIXED</u>		Confirm Mapped Type? <u> </u> Yes <u>X</u> No	

Profile Description:					
Depth inches	Horizon	Matrix Color (Munsell Moist)	Mottle Colors (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc.
0-10	1	10YR 4/2			Silty clay

Hydric Soil Indicators:	
<input type="checkbox"/> Histosol <input type="checkbox"/> Histic Epipedon <input type="checkbox"/> Sulfidic Odor <input type="checkbox"/> Aquic Moisture Regime <input type="checkbox"/> Reducing Conditions <input type="checkbox"/> Gleyed or Low-Chroma Colors	<input type="checkbox"/> Concretions <input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils <input type="checkbox"/> Organic Streaking in Sandy Soils <input type="checkbox"/> Listed on Local Hydric Soils List <input type="checkbox"/> Listed on National Hydric Soils List <input type="checkbox"/> Other (Explain in Remarks)

Remarks:
 No hydric soil indicators observed. Soil was moist throughout the soil profile.

WETLAND DETERMINATION

Hydrophytic Vegetation Present? <u> </u> Yes <u>X</u> No Wetland Hydrology Present? <u> </u> Yes <u>X</u> No Hydric Soils Present? <u> </u> Yes <u>X</u> No	Is this Sampling Point Within a Wetland? <u> </u> Yes <u>X</u> No
---	---

Remarks: The site was disturbed by mitigation construction in 2005. Site lacks dominance by hydrophytic vegetation and does not exhibit hydric soil indicators. Site hydrology is in flux. This area appears to have been inundated at least briefly prior to the field visit, but not long enough to promote the establishment and dominance of hydrophytic vegetation.

MDT MONTANA WETLAND ASSESSMENT FORM (revised March 2008)

1. Project Name: Wagner Marsh Wetland Mitigation Site 2. MDT Project #: _____ 3. Control #: _____

3. Evaluation Date: 8/8/2008 4. Evaluator(s): RRM (PBS&J) 5. Wetland/Site #(s): Wagner Marsh

6. Wetland Location(s): Township 1 S, Range 25 E, Section 28; Township _____ N, Range _____ E, Section _____

Approximate Stationing or Roadposts: _____

Watershed: 13 - Upper Yellowstone County: Yellowstone

7. Evaluating Agency: PBS&J

8. Wetland Size (acre): _____ (visually estimated)

Purpose of Evaluation:

7.38 (measured, e.g. GPS)

☐ Wetland potentially affected by MDT project

☐ Mitigation wetlands; pre-construction

☒ Mitigation wetlands; post-construction

☐ Other _____

9. Assessment Area (AA) Size (acre): _____ (visually estimated)

(see manual for determining AA) 16.19 (measured, e.g. GPS)

10. CLASSIFICATION OF WETLAND AND AQUATIC HABITATS IN AA (See manual for definitions.)

HGM Class (Brinson)	Class (Cowardin)	Modifier (Cowardin)	Water Regime	% OF AA
Depressional	Emergent Wetland	Excavated	Seasonal / Intermittent	36
Depressional	Aquatic Bed	Excavated	Permanent / Perennial	34
Depressional	Scrub-Shrub Wetland	Excavated	Seasonal / Intermittent	10
Depressional	Unconsolidated Bottom	Excavated	Permanent / Perennial	20

Comments: A mitigation site created in an old MDT gravel pit.

11. ESTIMATED RELATIVE ABUNDANCE (of similarly classified sites within the same Major Montana Watershed Basin; see manual.)
abundant

12. GENERAL CONDITION OF AA

i. Disturbance: Use matrix below to select the appropriate response; see manual for Montana listed noxious weed and aquatic nuisance vegetation species lists.

Conditions within AA	Predominant Conditions Adjacent to (within 500 feet of) AA		
	Managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings; and noxious weed or ANVS cover is ≤15%.	Land not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to minor clearing; contains few roads or buildings; noxious weed or ANVS cover is ≤30%.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >30%.
AA occurs and is managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings; and noxious weed or ANVS cover is ≤15%.	---	---	moderate disturbance
AA not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings; noxious weed or ANVS cover is ≤30%.	---	---	---
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >30%.	---	---	---

Comments (types of disturbance, intensity, season, etc.): Wetland mitigation site constructed in 2005. Disturbance within the AA has been high in the past, but with the creation of the wetland mitigation site the disturbance has ceased. No further disturbances are expected onsite. Immediately west of the site a new gravel pit has been constructed by a private company.

ii. Prominent noxious, aquatic nuisance, and other exotic vegetation species: Some Russian olive occurs scattered around the site. Salt cedar does occur, but is being managed effectively and is becoming less and less. Some limited amounts of Canada thistle occurs in wetland areas; spotted knapweed is being controlled in the uplands. Japanese brome also occurs in the uplands.

iii. Provide brief descriptive summary of AA and surrounding land use/habitat: AA is an old gravel pit converted into a groundwater dependent wetland complex. Surrounding land has a rolling topography and land use is predominantly agricultural - hay and livestock production. A gravel pit has been constructed on the west side of S. 56th St. W.

13. STRUCTURAL DIVERSITY (Based on number of "Cowardin" vegetated classes present [do not include unvegetated classes]; see #10 above.)

Existing # of "Cowardin" Vegetated Classes in AA	Initial Rating	Is current management preventing (passive) existence of additional vegetated classes?	Modified Rating
≥3 (or 2 if one is forested) classes	high	NA	NA
2 (or 1 if forested) classes	---	NA	NA
1 class, but not a monoculture	---	←NO	---
1 class, monoculture (1 species comprises ≥90% of total cover)	---	NA	NA

Comments: PSS, PEM, PAB. Some scattered cottonwoods.

Wetland/Site #(s): Wagner Marsh**14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS OR ANIMALS****i. AA is Documented (D) or Suspected (S) to contain:** Check box based on definitions in manual.

Primary or critical habitat (**list species**) ☐ D ☐ S _____
 Secondary habitat (**list species**) ☐ D ☐ S _____
 Incidental habitat (**list species**) ☐ D ☐ S _____
 No usable habitat ☒ S

ii. Rating: Based on the strongest habitat chosen in 14A(i) above, select the corresponding functional point and rating.

Highest Habitat Level	Doc/Primary	Sus/Primary	Doc/Secondary	Sus/Secondary	Doc/Incidental	Sus/Incidental	None
Functional Point/Rating	---	---	---	---	---	---	0L

Sources for documented use (e.g. observations, records): _____

14B. HABITAT FOR PLANTS OR ANIMALS RATED S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM

Do not include species listed in 14A above.

i. AA is Documented (D) or Suspected (S) to contain: Check box based on definitions in manual.

Primary or critical habitat (**list species**) ☐ D ☐ S _____
 Secondary habitat (**list species**) ☒ D ☐ S Sandhill crane (S2N), migrating raptors
 Incidental habitat (**list species**) ☐ D ☐ S _____
 No usable habitat ☐ S

ii. Rating: Based on the strongest habitat chosen in 14A(i) above, select the corresponding functional point and rating.

Highest Habitat Level	Doc/Primary	Sus/Primary	Doc/Secondary	Sus/Secondary	Doc/Incidental	Sus/Incidental	None
S1 Species Functional Point/Rating	---	---	---	---	---	---	---
S2 and S3 Species Functional Point/Rating	---	---	.6M	---	---	---	---

Sources for documented use (e.g. observations, records): Observed during site visits enough to believe that sandhill cranes are using the site regularly.**14C. GENERAL WILDLIFE HABITAT RATING****i. Evidence of Overall Wildlife Use in the AA:** Check substantial, moderate, or low based on supporting evidence.☒ **Substantial:** Based on any of the following [check].

- ☒ observations of abundant wildlife #s or high species diversity (during any period)
- ☒ abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ presence of extremely limiting habitat features not available in the surrounding area
- ☐ interview with local biologist with knowledge of the AA

☐ **Minimal:** Based on any of the following [check].

- ☐ few or no wildlife observations during peak use periods
- ☐ little to no wildlife sign
- ☐ sparse adjacent upland food sources
- ☐ interview with local biologist with knowledge of AA

☐ **Moderate:** Based on any of the following [check].

- ☐ observations of scattered wildlife groups or individuals or relatively few species during peak periods
- ☐ common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ adequate adjacent upland food sources
- ☐ interview with local biologist with knowledge of the AA

ii. Wildlife Habitat Features: Working from top to bottom, check appropriate AA attributes in matrix to arrive at rating. Structural diversity is from #13. For class cover to be considered evenly distributed, the most and least prevalent **vegetated** classes must be within 20% of each other in terms of their percent composition of the AA (see #10). Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; and A = absent [see manual for further definitions of these terms].

Structural Diversity (see #13)	<input checked="" type="checkbox"/> High								<input type="checkbox"/> Moderate								<input type="checkbox"/> Low			
Class Cover Distribution (all vegetated classes)	<input type="checkbox"/> Even				<input checked="" type="checkbox"/> Uneven				<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even			
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
<input type="checkbox"/> Low Disturbance at AA (see #12i)	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<input type="checkbox"/> Moderate Disturbance at AA (see #12i)	---	---	---	---	H	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<input type="checkbox"/> High Disturbance at AA (see #12i)	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

iii. Rating: Use the conclusions from i and ii above and the matrix below to select the functional point and rating.

Evidence of Wildlife Use (i)	Wildlife Habitat Features Rating (ii)			
	<input type="checkbox"/> Exceptional	<input checked="" type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
<input checked="" type="checkbox"/> Substantial	---	.9H	---	---
<input type="checkbox"/> Moderate	---	---	---	---
<input type="checkbox"/> Minimal	---	---	---	---

Comments: Site is well used by migrating waterfowl.

Wetland/Site #(s): _____

14D. GENERAL FISH HABITAT ☒ **NA** (proceed to 14E)

If the AA is not used by fish, fish use is not restorable due to habitat constraints, or is not desired from a management perspective [such as fish entrapped in a canal], then check the NA box and proceed to 14E.

Assess this function if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [i.e., fish use is precluded by perched culvert or other barrier].

Type of Fishery: ☐ Cold Water (CW) ☐ Warm Water (WW) Use the CW or WW guidelines in the manual to complete the matrix.

i. Habitat Quality and Known / Suspected Fish Species in AA: Use matrix to select the functional point and rating.

Duration of Surface Water in AA	<input type="checkbox"/> Permanent / Perennial						<input type="checkbox"/> Seasonal / Intermittent						<input type="checkbox"/> Temporary / Ephemeral					
Aquatic Hiding / Resting / Escape Cover	<input type="checkbox"/> Optimal		<input type="checkbox"/> Adequate		<input type="checkbox"/> Poor		<input type="checkbox"/> Optimal		<input type="checkbox"/> Adequate		<input type="checkbox"/> Poor		<input type="checkbox"/> Optimal		<input type="checkbox"/> Adequate		<input type="checkbox"/> Poor	
Thermal Cover: optimal / suboptimal	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S
FWP Tier I fish species	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
FWP Tier II or Native Game fish species	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
FWP Tier III or Introduced Game fish	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
FWP Non-Game Tier IV or No fish species	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

Sources used for identifying fish spp. potentially found in AA: _____

ii. Modified Rating: NOTE: Modified score cannot exceed 1.0 or be less than 0.1.

a) Is fish use of the AA significantly reduced by a culvert, dike, or other man-made structure or activity, **or** is the waterbody included on the current final MDEQ list of waterbodies in need of TMDL development with listed "Probable Impaired Uses" including cold or warm water fishery or aquatic life support, **or** do aquatic nuisance plant or animal species (see **Appendix E**) occur in fish habitat? ☐ **YES**, reduce score in i by 0.1 = ____ or ☐ **NO**

b) Does the AA contain a documented spawning area or other critical habitat feature (i.e., sanctuary pool, upwelling area; specify in comments) for native fish or introduced game fish? ☐ **YES**, add to score in i or iia 0.1 = ____ or ☐ **NO**

iii. Final Score and Rating: **Comments:** Though the Biological Resources Rpt states that black-nosed dace and carp can be found within the ponds no fish have been observed during site visits in 2005, 2006, 2007, or 2008 and no surface inlet or outlet exists. The ponds are relatively shallow and so provide poor overwintering habitat for fish.

14E. FLOOD ATTENUATION ☒ **NA** (proceed to 14F)


Applies only to wetlands that are subject to flooding via in-channel or overbank flow.

If wetlands in AA are not flooded from in-channel or overbank flow, check the NA box and proceed to 14F.

Entrenchment Ratio (ER) Estimation (see manual for additional guidance). Entrenchment ratio = (flood-prone width) / (bankfull width).

Flood-prone width = estimated horizontal projection of where 2 X maximum bankfull depth elevation intersects the floodplain on each side of the stream.

_____ / _____ = _____
flood prone width / bankfull width = entrenchment ratio



Slightly Entrenched ER ≥ 2.2			Moderately Entrenched ER = 1.41 – 2.2		Entrenched ER = 1.0 – 1.4		
C stream type	D stream type	E stream type	B stream type		A stream type	F stream type	G stream type

i. Rating: Working from top to bottom, use the matrix below to select the functional point and rating.

Estimated or Calculated Entrenchment (Rosgen 1994, 1996)	<input type="checkbox"/> Slightly Entrenched C, D, E stream types			<input type="checkbox"/> Moderately Entrenched B stream type			<input type="checkbox"/> Entrenched A, F, G stream types		
Percent of Flooded Wetland Classified as Forested and/or Scrub/Shrub	<input type="checkbox"/> 75%	<input type="checkbox"/> 25-75%	<input type="checkbox"/> <25%	<input type="checkbox"/> 75%	<input type="checkbox"/> 25-75%	<input type="checkbox"/> <25%	<input type="checkbox"/> 75%	<input type="checkbox"/> 25-75%	<input type="checkbox"/> <25%
AA contains no outlet or restricted outlet	---	---	---	---	---	---	---	---	---
AA contains unrestricted outlet	---	---	---	---	---	---	---	---	---

ii. Are ≥10 acres of wetland in the AA subject to flooding AND are man-made features which may be significantly damaged by floods located within 0.5 mile downstream of the AA? ☐ **YES** ☐ **NO** **Comments:** _____

Wetland/Site #(s): Wagner Marsh**14F. SHORT AND LONG TERM SURFACE WATER STORAGE** ☐ NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow.
If no wetlands in the AA are subject to flooding or ponding, then check the NA box and proceed to 14G.

- i. **Rating:** Working from top to bottom, use the matrix below to select the functional point and rating. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral [see manual for further definitions of these terms].

Estimated Maximum Acre Feet of Water Contained in Wetlands within the AA that are Subject to Periodic Flooding or Ponding	<input checked="" type="checkbox"/> >5 acre feet			<input type="checkbox"/> 1.1 to 5 acre feet			<input type="checkbox"/> ≤1 acre foot		
Duration of Surface Water at Wetlands within the AA	<input checked="" type="checkbox"/> P/P	<input type="checkbox"/> S/I	<input type="checkbox"/> T/E	<input type="checkbox"/> P/P	<input type="checkbox"/> S/I	<input type="checkbox"/> T/E	<input type="checkbox"/> P/P	<input type="checkbox"/> S/I	<input type="checkbox"/> T/E
Wetlands in AA flood or pond ≥ 5 out of 10 years	1H	---	---	---	---	---	---	---	---
Wetlands in AA flood or pond < 5 out of 10 years	---	---	---	---	---	---	---	---	---

Comments: _____

14G. SEDIMENT / NUTRIENT / TOXICANT / RETENTION AND REMOVAL ☐ NA (proceed to 14H)

Applies to wetland with potential to receive sediments, nutrients, or toxicants through influx of surface or ground water or direct input.
If no wetlands in the AA are subject to such input, check the NA box and proceed to 14H.

- i. **Rating:** Working from top to bottom, use the matrix below to select the functional point and rating.

Sediment, Nutrient, and Toxicant Input Levels within AA	AA receives or surrounding land use has potential to deliver sediments, nutrients, or compounds at levels such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody is on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% Cover of Wetland Vegetation in AA	<input type="checkbox"/> ≥ 70%		<input checked="" type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
Evidence of Flooding / Ponding in AA	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
AA contains no or restricted outlet	---	---	.7M	---	---	---	---	---
AA contains unrestricted outlet	---	---	---	---	---	---	---	---

Comments: _____

14H. SEDIMENT / SHORELINE STABILIZATION ☐ NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action.
If 14H does not apply, check the NA box and proceed to 14I.

% Cover of <u>Wetland</u> Streambank or Shoreline by Species with Stability Ratings of ≥6 (see Appendix F).	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input checked="" type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
<input type="checkbox"/> ≥ 65%	---	---	---
<input checked="" type="checkbox"/> 35-64%	.7M	---	---
<input type="checkbox"/> < 35%	---	---	---

Comments: Shoreline vegetation continues to become established.**14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT**

- i. **Level of Biological Activity:** Synthesis of wildlife and fish habitat rates (select).

General Fish Habitat Rating (14Di)ii)	General Wildlife Habitat Rating (14Ci)ii)		
	<input checked="" type="checkbox"/> E/H	<input type="checkbox"/> M	<input type="checkbox"/> L
<input type="checkbox"/> E/H	---	---	---
<input type="checkbox"/> M	---	---	---
<input type="checkbox"/> L	---	---	---
<input checked="" type="checkbox"/> NA	H	---	---

- ii. **Rating:** Working from top to bottom, use the matrix below to select the functional point and rating. Factor A = acreage of vegetated wetland component in the AA; Factor B = level of biological activity rating from above (14Ii); Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to the duration of surface water in the AA, where P/P, S/I, and T/E were previously defined, and A = "absent" [see manual for further definitions of these terms].

A	<input checked="" type="checkbox"/> Vegetated Component >5 acres						<input type="checkbox"/> Vegetated Component 1-5 acres						<input type="checkbox"/> Vegetated Component <1 acre					
B	<input checked="" type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low	
C	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
P/P	---	.7M	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
S/I	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
T/E/A	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

Wetland/Site #(s): Wagner Marsh**14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT** (continued)iii. **Modified Rating:** Note: Modified score cannot exceed 1.0 or be less than 0.1.**Vegetated Upland Buffer:** Area with $\geq 30\%$ plant cover, $\leq 15\%$ noxious weed or ANVS cover, AND that is not subjected to periodic mechanical mowing or clearing (unless for weed control).Is there an average ≥ 50 -foot wide vegetated upland buffer around $\geq 75\%$ of the AA's perimeter? ☒ **YES**, add 0.1 to score in ii = 0.80 ☐ **NO**iv. **Final Score and Rating:** .8H **Comments:** _____**14J. GROUNDWATER DISCHARGE / RECHARGE**

Check the appropriate indicators in i and ii below.

i. Discharge Indicators

- ☐ The AA is a slope wetland.
☐ Springs or seeps are known or observed.
☒ Vegetation growing during dormant season/drought.
☐ Wetland occurs at the toe of a natural slope.
☐ Seeps are present at the wetland edge.
☒ AA permanently flooded during drought periods.
☐ Wetland contains an outlet, but no inlet.
☐ Shallow water table and the site is saturated to the surface.
☐ Other: _____

ii. Recharge Indicators

- ☐ Permeable substrate present without underlying impeding layer.
☐ Wetland contains inlet but no outlet.
☐ Stream is a known 'losing' stream. Discharge volume decreases.
☐ Other: _____

iii. **Rating:** Use the information from i and ii above and the table below to select the functional point and rating.

Criteria	Duration of Saturation at AA Wetlands <i>FROM GROUNDWATER DISCHARGE</i> or <i>WITH WATER THAT IS RECHARGING THE GROUNDWATER SYSTEM</i>			
	<input checked="" type="checkbox"/> P/P	<input type="checkbox"/> S/I	<input type="checkbox"/> T	<input type="checkbox"/> None
<input checked="" type="checkbox"/> Groundwater Discharge or Recharge	1H	---	---	---
<input type="checkbox"/> Insufficient Data/Information	---			

Comments: _____**14K. UNIQUENESS**i. **Rating:** Working from top to bottom, use the matrix below to select the functional point and rating.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland OR plant association listed as "S1" by the MTNHP			AA does not contain previously cited rare types AND structural diversity (#13) is high OR contains plant association listed as "S2" by the MTNHP			AA does not contain previously cited rare types OR associations AND structural diversity (#13) is low-moderate		
	<input type="checkbox"/> Rare	<input type="checkbox"/> Common	<input type="checkbox"/> Abundant	<input type="checkbox"/> Rare	<input type="checkbox"/> Common	<input checked="" type="checkbox"/> Abundant	<input type="checkbox"/> Rare	<input type="checkbox"/> Common	<input type="checkbox"/> Abundant
<input checked="" type="checkbox"/> Low Disturbance at AA (#12i)	---	---	---	---	---	.5M	---	---	---
<input type="checkbox"/> Moderate Disturbance at AA (#12i)	---	---	---	---	---	---	---	---	---
<input type="checkbox"/> High Disturbance at AA (#12i)	---	---	---	---	---	---	---	---	---

Comments: The site currently has a low level of disturbance.**14L. RECREATION / EDUCATION POTENTIAL**☐ NA (proceed to Overall Summary and Rating page)

Affords 'bonus' points if AA provides a recreational or educational opportunity.

i. **Is the AA a known or potential recreational or educational site?** ☒ **YES**, go to ii. ☐ **NO**, check the NA box.ii. **Check categories that apply to the AA:** ☒ Educational/Scientific Study ☐ Consumptive Recreational ☐ Non-consumptive recreational
☐ Other: _____iii. **Rating:** Use the matrix below to select the functional point and rating.

Known or Potential Recreational or Educational Area		Known	Potential
Public ownership or public easement with general public access (no permission required)		---	---
Private ownership with general public access (no permission required)		---	---
Private or public ownership without general public access, or requiring permission for public access		.1M	---

Comments: The site receives educational use through the WJH Bird Facility that is north-adjacent to the mitigation area.**15. GENERAL SITE NOTES:** _____

Wetland/Site #(s): _____

Function & Value Variables	Rating – Actual Functional Points	Possible Functional Points	Functional Units: Actual Points x Estimated AA Acreage	Indicate the Four Most Prominent Functions with an Asterisk
A. Listed / Proposed T&E Species Habitat	low 0.00	1.00	0	
B. MT Natural Heritage Program Species Habitat	mod 0.60	1.00	9.71	
C. General Wildlife Habitat	high 0.90	1.00	14.57	*
D. General Fish Habitat	NA	---		
E. Flood Attenuation	NA	---		
F. Short and Long Term Surface Water Storage	high 1.00	1.00	16.19	*
G. Sediment / Nutrient / Toxicant Removal	mod 0.70	1.00	11.33	
H. Sediment / Shoreline Stabilization	mod 0.70	1.00	11.33	
I. Production Export / Food Chain Support	high 0.80	1.00	12.95	*
J. Groundwater Discharge / Recharge	high 1.00	1.00	16.19	
K. Uniqueness	mod 0.50	1.00	8.10	
L. Recreation / Education Potential (bonus point)	mod 0.10		1.62	*
Total Points	6.3	9	102 Total Functional Units	
Percent of Possible Score 70% (round to nearest whole number)				

Category I Wetland: (must satisfy **one** of the following criteria; otherwise go to Category II)

- ☐ Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**
☐ Score of 1 functional point for Uniqueness; **or**
☐ Score of 1 functional point for Flood Attenuation **and** answer to Question 14E.ii is "yes"; **or**
☐ Percent of possible score > 80% (round to nearest whole #).

Category II Wetland: (Criteria for Category I not satisfied **and** meets any **one** of the following criteria; otherwise go to Category IV)

- ☐ Score of 1 functional point for MT Natural Heritage Program Species Habitat; **or**
☒ Score of .9 or 1 functional point for General Wildlife Habitat; **or**
☐ Score of .9 or 1 functional point for General Fish Habitat; **or**
☐ "High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish/Aquatic Habitat; **or**
☐ Score of .9 functional point for Uniqueness; **or**
☒ Percent of possible score > 65% (round to nearest whole #).

☐ **Category III Wetland:** (Criteria for Categories I, II, or IV not satisfied)**Category IV Wetland:** (Criteria for Categories I or II are not satisfied and all of the following criteria are met; if not go to Category III)

- ☐ "Low" rating for Uniqueness; **and**
☐ Vegetated wetland component < 1 acre (do not include upland vegetated buffer); **and**
☐ Percent of possible score < 35% (round to nearest whole #).

OVERALL ANALYSIS AREA (AA) RATING: Check the appropriate category based on the criteria outlined above.
☐ I ☒ II ☐ III ☐ IV

Appendix C

2008 REPRESENTATIVE PHOTOGRAPHS

MDT Wetland Mitigation Monitoring
Wagner Marsh
Billings, Montana

Wagner Marsh Wetland Mitigation Site 2008



Photo Point A – Photo 1 Location: North Side
Compass bearing: 22 degrees



Photo Point A – Photo 2 Location: North Side
Compass bearing: 105 degrees



Photo Point A – Photo 3 Location: North Side
Compass bearing: 162 degrees



Photo Point A – Photo 4 Location: North Side
Compass bearing: 214 degrees



Photo Point A – Photo 5 Location: North Side
Compass bearing: 250 degrees



Photo Point A – Photo 6 Location: North Side
Compass bearing: 310 degrees



Photo Point A – Photo 7 Location: North Side
Compass bearing: 335 degrees



Photo Point B – Photo 1 Location: West Side
Compass bearing: 01 degrees

Wagner Marsh Wetland Mitigation Site 2008



Photo Point B – Photo 2 Location: West Side
Compass bearing: 74 degrees



Photo Point B – Photo 3 Location: West Side
Compass bearing: 153 degrees



Photo Point C – Photo 1 Location: South Side
Compass bearing: 24 degrees



Photo Point C – Photo 2 Location: South Side
Compass bearing: 243 degrees



Photo Point C – Photo 3 Location: South Side
Compass bearing: 294 degrees



Photo Point C – Photo 4 Location: South Side
Compass bearing: 343 degrees



Photo Point D – Photo 1 Location: East Side
Compass bearing: 241 degrees



Photo Point D – Photo 2 Location: East Side
Compass bearing: 293 degrees

Wagner Marsh Wetland Mitigation Site 2008



Photo Point D – Photo 3 Location: East Side
Compass bearing: 324 degrees



Photo Point D – Photo 4 Location: East Side
Compass bearing: 356 degrees



Transect Photo Point #1 Location: West end
Compass bearing: 70 degrees



Transect Photo Point #2 Location: East end
Compass bearing: 250 degrees



2008 macroinvertebrate sampling location

Appendix D

CONCEPTUAL SITE LAYOUT

*MDT Wetland Mitigation Monitoring
Wagner Marsh
Billings, Montana*

WETLAND - 1.16 AC

POND - 1.03 AC

WETLAND - 2.71 AC

STAGING AREA

A-7

991.30

B-2

B-1A

B-1

A-6

A-5

A-3


A-4

POND - 0.72 AC

NEW WETLANDS
17.72 AC

- New Created Wetlands
- Old Created Wetlands
- Open Water

WAGNER PIT WETLAND AREAS



Appendix E

BIRD SURVEY PROTOCOL GPS PROTOCOL

*MDT Wetland Mitigation Monitoring
Wagner Marsh
Billings, Montana*

BIRD SURVEY PROTOCOL

This protocol was developed by the Montana Department of Transportation (MDT) to monitor bird use within their Wetland Mitigation Sites. Though each wetland mitigation site is vastly different, the bird survey data collection methods were standardized to order to increase repeatability. The protocol uses an "area search within a restricted time frame" to collect data on bird species, density, behavior, and habitat-type use.

Survey Area

Sites that can be entirely walked: Sites where the entire perimeter or area can be walked include, but are not limited to: small ponds, enhanced historic river channels, and wet meadows. If the wetland is not uncomfortably inundated, walk several meandering transects to sufficiently cover the wetland. Meandering transects can be used, even if a small portion of the area is inaccessible (e.g. cannot cross due to inundation). Use binoculars to identify the bird species, to count the number of individuals, and to identify their behavior and habitat type. Data can be recorded directly onto the bird survey form or into a field notebook. The number of meandering transects and their direction (or location) should be recorded in the field notebook and/or drawn onto the aerial photograph or topographic map. Meandering transects are not formal and should not be staked. Each site should be walked and surveyed to the fullest extent within the set time limit.

Sites than cannot be entirely walked: Sites where the entire perimeter or area cannot be walked include, but are not limited to: very large sites (i.e. perimeter of 2-3 miles), and large-bodied waters (i.e. reservoirs), where deep water habitat (> 6 feet) is close to shore. For large-bodied waters where only one area was graded to create or enhance the development of wetland, bird surveys should be walked along meandering transects within or around the graded area (see above.). For sites that cannot be walked, bird surveys should be conducted from many lookout posts, established at key vantage points. The general location of lookout posts should be recorded in the field notebook or drawn onto the aerial photograph or topographic map. Lookout post locations do not need to be staked. Both binoculars and spotting scopes may be used in order to accurately identify and count the birds. Depending upon the size of the open water, more time may be spent viewing the mitigation area from lookout posts than is spent traveling between posts.

Survey Time

Ideally, bird surveys should be conducted in the morning hours when bird activity is often greatest (i.e. sunrise to no later than 11:00 am). Surveys can be completed before 11am if all transects have been walked or all lookout posts have been viewed with no new bird activity observed. For some sites bird surveys may need to be performed in the late afternoon or evening due to traveling constraints or weather. The overall limiting time factor will be the number of budgeted hours for the project.

Data Recording

Bird Species List: Record each bird species observed onto the Bird Survey-Field Data Sheet (or field notebook). Record the bird's common name using the appropriate 4-letter code. The 4-letter code uses the first two letters of the first two word's of the bird's common name or if one name, the first four letters. For example, Mourning Dove is coded as MODO while Mallard is coded as MALL. If an unknown individual is observed, use the 4-letter protocol, but define your

BIRD SURVEY PROTOCOL (continued)

abbreviation at the bottom of the field data sheet. For example, unknown shorebird is UNSB; unknown brown bird is UNBR; unknown warbler is UNWA; and unknown waterfowl is UNWF. For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parenthesis; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded as UNBB / FO (25).

Bird Density: For each observation record the actual or estimated number of individuals observed per species and per behavior. Totals can be tallied in the office and entered onto the Bird Survey-Field Data Sheet.

Bird Behavior: Bird behavior must be identified by what is known. When a species is observed, the behavior that is immediately exhibited is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair (BP); foraging (F); flyover (FO); loafing (L), which is defined as sleeping, roosting, or floating with head tucked under wing; and nesting (N). If other behaviors that have a specific descriptive word are observed then it can be used and should later be added to the protocol. Descriptive words or phrases such as "migrating" or "living on site" are unknown behaviors.

Bird Species Habitat Use: When a species is observed, the habitat is also recorded. The following broad habitat categories are used:

- ◆ aquatic bed (AB), defined as rooted-floating, floating-leaved, or submergent vegetation.
- ◆ marsh (MA), defined as emergent (e.g. cattail, bulrush) vegetation with surface water.
- ◆ wet meadow (WM), defined as grasses, sedges, or rushes with little to no surface water.
- ◆ scrub-shrub (SS), defined as shrub covered wetland.
- ◆ forested (FO), defined as tree covered wetland.
- ◆ open water (OW), defined as unvegetated surface water.
- ◆ upland (UP), defined as the upland buffer.

Other categories can be used and defined on the data sheet and should later be added to the protocol.

Other Fields

Bird Visit: Each bird survey (i.e. spring, fall, and mid-season) should be completed on separate Bird Survey-Field Data Sheets.

Time: Record the start time and end time on the Bird Survey-Field Data Sheet.

Date: Record the date of the bird survey.

Weather: Record the weather conditions (i.e. temperature, wind, condition).

Notes: Note if a particular individual bird is using a constructed nest box and note the condition of constructed nest box(es). Also record any comments about the site, wildlife, wetland conditions, etc.

GPS MAPPING AND AERIAL PHOTO REFERENCING PROCEDURE

From 2001 through 2006, PBS&J mapped the vegetation community boundaries, photograph points, and other sampling locations in the field using the resource-grade Trimble GEO III GPS (Global Positioning System) unit. The data were collected with a minimum of three positions per feature using Course/Acquisition code. The collected data were then transferred to a personal computer (PC) and differentially corrected to the nearest operating Community Base Station. The corrected data were then exported to ACAD drawings in Montana State Plane Coordinates NAD 83 international feet. The Trimble GEO III GPS unit was also used for some sites in 2007.

The collected and processed Trimble Geo III GPS positions had a 68% accuracy of 7 feet except in isolated areas where accuracy fell to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

In 2007 and 2008 sites were mapped using the resource-grade Magellan MobileMapper Office GPS unit. The Magellan GPS unit has a comparable accuracy level to the Trimble Geo III unit.

Each year, MDT photographs each mitigation site from the air. These aerial photographs are not geo-referenced, but serve as a visual aid to map wetland development and vegetation communities, and to show approximate locations for various monitoring activities (i.e. photograph points, transects, or macroinvertebrate sampling). Reference points that are observable on the aerial photo (i.e. road, stream channel, or fence) were also marked with the GPS unit in order to better position the aerial photograph. This positioning did not remove any of the distortion inherent to all photos. All mapped features and community boundaries were reviewed by the wetland biologist, to increase the figure's accuracy.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

Appendix F

2008 MACROINVERTEBRATE SAMPLING PROTOCOL AND DATA

*MDT Wetland Mitigation Monitoring
Wagner Marsh
Billings, Montana*

AQUATIC INVERTEBRATE SAMPLING PROTOCOL

Equipment List

- D-frame sampling net with 1 mm mesh.
- 1-liter, wide-mouth, plastic sample jars provided by Rhithron Associates, Inc. (Quart sized, wide-mouthed canning jars can be substituted.)
- 95% ethanol (alternatively isopropyl alcohol).
- Pre-printed sample labels (printed on rite-in-the-rain paper); two labels per sample.
- Pencil.
- Clear packaging tape.
- 3-5 gallon plastic pail.
- Large tea strainer or framed screen.
- Cooler with ice for storing sample.

Site Selection

Select a site that is accessible with hip waders or rubber boots. If the substrate is too soft, place a wide board down to walk on. Choose a site that is representative of the overall condition of the wetland. Annual sampling should occur at the same site within the wetland.

Sampling Procedure

Wetland invertebrates (macroinvertebrates) inhabit the substrate, the water column, the stems and leaves of aquatic vegetation, and the water surface. At the given location, each habitat type is sampled and combined into a single 1-liter sample jar. Pre-cautions are made to minimize disturbing the sample site in order to maximize the number of animals collected.

Fill the pail with approximately 1 gallon of wetland water. Ideally, sample the water column from near-shore outward to a depth of 3 feet. Sample the water column using a long sweep of the net, keeping the net at about half the depth of the water. Sample the water surface with a long sweep of the net. Aquatic vegetation is sampled by pulling the net beneath the water surface, for at least a meter in distance. The substrate is sampled by pulling the net along the bottom, bumping it against the substrate several times as you pull. Be sure to place some muck, mud, and/or vegetation into the jar. After sampling a habitat, rinse the net in the bucket and look for insects, crustaceans, and other aquatic invertebrates. It is not necessary to sample habitats in any specific order, but all habitats, if present, are to be sampled. Habitats can be sampled more than once.

Fill about 1 cup of ethanol into the sample jar. Sieve the contents of the bucket through the straining device and pour or carefully scrape the contents of the strainer into the sample jar. Top off the jar with enough ethanol to cover all the material and leave as little headroom as possible. Alternatively, sampled materials can be lifted out of the net and put directly into the jar. Be sure to include some muck, mud, and/or vegetation into the jar. Each macroinvertebrate sampling site should have only one sampling jar.

Using pencil, complete two labels with the required information: project name, project number, date, collector's name, and habitats sampled. Do not complete the label with ink as it will dissolve in ethanol. For wetlands with at least two macroinvertebrate sampling sites, number the site consecutively followed by the total number of sites (e.g. Sample 2 of 3 sites). Place one label into the jar and seal the jar. Dry the jar off, if necessary, and tape the second label to the outside of the jar.

Photograph each macroinvertebrate sampling site.

Sample Handling/Delivery

In the field, keep sample jars cool by placing in a cooler with a small amount of ice.

Deliver samples to the PBS&J office in Missoula, where they will be inventoried and delivered to Rhithron Associates, Inc.

**MDT Mitigated Wetland Monitoring Project: Aquatic Invertebrate Monitoring
Summary 2001 – 2008**

Prepared for Post, Buckley, Schuh, and Jernigan (PBS&J)
Prepared by W. Bollman, Rhithron Associates, Inc.

INTRODUCTION

This report summarizes data generated from eight years of mitigated wetland monitoring from sites throughout the State of Montana. Over all years of sampling, a total of 210 invertebrate samples have been collected. Table 1 lists the currently monitored sites at which aquatic invertebrates were collected in 2008, and summarizes the sampling history of each.

METHODS

Sample processing

Aquatic invertebrate samples were collected at mitigated wetland sites in the summer months of 2001, 2002, 2003, 2004, 2005, 2006, 2007, and 2008 by personnel of PBS&J (Table 1). Sampling procedures were based on the protocols developed by the Montana Department of Environmental Quality (MDEQ) for wetland sampling. Sampling consisted of D-frame net sweeps through emergent vegetation (when present), the water column, and over the water surface, and included disturbing and scraping substrates at each sampled site. These sample components were composited and preserved in ethanol at each wetland site. Samples were delivered to Rhithron Associates, Inc. for processing, taxonomic determinations, and data analysis.

Standard sorting protocols were applied to achieve representative subsamples of a minimum of 100 organisms. Caton sub-sampling devices (Caton 1991), divided into 30 grids, each approximately 5 cm by 6 cm, were used. Grid contents were examined under stereoscopic microscopes using 10x-30x magnification. All aquatic invertebrates from each selected grid were sorted from the substrate, and placed in 95% ethanol for subsequent identification. Grid selection, examination, and sorting continued until at least 100 organisms were sorted. A large/rare search was conducted to collect any taxa not found in the subsampling procedure.

Organisms were individually examined using 10x – 80x stereoscopic dissecting scopes (Leica S8E and S6E) and identified to the lowest practical taxonomic levels using appropriate published taxonomic references. Identification, counts, life stages, and information about the condition of specimens were recorded on bench sheets. To obtain accuracy in richness measures, organisms that could not be identified to the target level specified in MDEQ protocols were designated as “not unique” if other specimens from the same group could be taken to target levels. Organisms designated as “unique” were those that could be definitively distinguished from other organisms in the sample. Identified organisms were preserved in 95% ethanol in labeled vials, and archived at the Rhithron laboratory. Midges were morphotyped using 10x – 80x stereoscopic dissecting microscopes (Leica S8E and S6E) and representative specimens were slide mounted and examined at 200x – 1000x magnification using an Olympus BX 51 compound microscope. Slide mounted organisms were also archived at the Rhithron laboratory.

Assessment

The method employed to assess these wetlands is based on an index incorporating a battery of 12 bioassessment metrics or attributes (Table 2) tested and recommended by Stribling et al. (1995) in a report to the Montana Department of Health and Environmental Science. In that study, it was determined that some of the metrics were of limited use in some geographic regions, and for some wetland types. Despite that finding, all 12 metrics are used in this evaluation of mitigated wetlands, since detailed geographic information and wetland classifications were unavailable. Scoring criteria for the 12 metrics were developed specifically for this project, since mitigated wetlands were not included in original criteria development.

Scoring criteria for wetland metrics were developed by generally following the tactic used by Stribling et al. (1995). Boxplots were generated using a statistical software package (Statistica™), and distributions, median values, ranges, and quartiles for each metric were examined. For the wetland sites, “good” scores were generally

those that fell above the 75th percentile (for those metrics that decrease in value in response to stress) or below the 25th percentile (for metrics that respond to stress by an increase in value) of all scores. Additional scoring ranges were established by bisecting the range below the 75th percentile for decreasing scores (or above the 25th percentile for increasing scores) into “sub-optimal” and “poor” assessment categories. A score of 5, 3, or 1 was assigned to good, sub-optimal, and poor metric performance, respectively. In this way, metric values were translated into normalized metric scores, and scores for all metrics were summed to produce a total bioassessment score, which is expressed as a percentage of the maximum possible score (60). Total bioassessment scores were classified according to a similar process, using the ranges and distributions of total scores for all sites studied in all years. Data from a total of 167 samples were used to develop criteria.

Six sites in this study supported aquatic fauna characteristic of lotic habitats rather than lentic wetland habitats; these sites were excluded from mitigated wetland scoring criteria development, and were evaluated with a metric battery specific to flowing water habitats. In 2008, the lotic sites were Camp Creek (2 sites), Cloud Ranch stream, Jack Creek – McKee Spring, and Jocko Spring Creek (2 sites). Invertebrate assemblages at these sites were generally characteristic of montane or foothill stream conditions and were assessed using the tested metric battery developed for montane streams of Western Montana (MVFP index: Bollman 1998).

The purpose of constructing an index from biological attributes or metrics is to provide a means of integrating information to facilitate the determination of whether management action is needed. However, the nature of the action needed is not determined solely by the index score or impairment classification, but by consideration of an analysis of the component metrics, the taxonomic composition of the assemblages, and other issues. The diagnostic functions of the metrics and taxonomic data need more study since our understanding of the interrelationships of natural environmental factors and anthropogenic disturbances is tentative. Thus, the further interpretive remarks accompanying the raw taxonomic and metric data in this summary are offered cautiously. Year-to-year comparisons depend on an assumption that specific sites were revisited in each year, and that equivalent sampling methods were utilized at each site revisit.

Bioassessment metrics – wetlands

An index based on the performance of 12 metrics was constructed, as described above. Table 2 lists those metrics, describes their calculation and the expected response of each to increased degradation or impairment of the wetland.

In addition to the summed scores of each metric and the associated impairment classification described above, each individual metric informs the bioassessment to some degree. The four richness metrics (Total taxa, POET, Chironomidae taxa, and Crustacea taxa + Mollusca taxa) can be interpreted to express habitat complexity as well as water quality. Complex, diverse habitats consist of variable substrates, emergent vegetation, variable water depths and other factors, and are potential features of long-established stable wetlands with minimal human disturbance. In the study conducted by Stribling et al. (1995), all four richness metrics were found to be significantly associated with water quality parameters including conductance, salinity, and total dissolved solids.

Four composition metrics (%Chironomidae, %Orthocladiinae of Chironomidae, %Crustacea + %Mollusca, and %Amphipoda) measure the relative contributions of certain taxonomic groups that may have significant responses to habitat and/or water quality impacts. For example, amphipods have been demonstrated to increase in abundance in alkaline conditions. Short-lived, relatively mobile taxa such as chironomids dominate ephemeral environments; many are hemoglobin-bearers capable of tolerating de-oxygenated conditions.

Two tolerance metrics (Hilsenhoff Biotic Index and %Dominant taxon) were included in the bioassessment battery. The HBI indicates the overall invertebrate assemblage tolerance to nutrient enrichment, warm water, and/or low dissolved oxygen conditions. The percent abundance of the dominant taxon has been demonstrated to be strongly associated with pH, conductance, salinity, total organic carbon, and total dissolved solids.

Two trophic measures (%Collector-gatherers and %Filterers) may be helpful in expressing functional integrity of the invertebrate assemblage, which can be impacted by poor water quality or habitat degradation. High proportions of filtering organisms suggest nutrient and/or organic enrichment, while abundant collectors suggest

more positive functional conditions and well-developed wetland morphology. These organisms graze periphyton growing on stable surfaces such as macrophytes.

Summary metric values and scores for the 2008 samples are given in Tables 4a-4c and 5. Thermal preference of invertebrate assemblages was calculated using Brandt 2001.

Bioassessment metrics – lotic habitats

For sites supporting rheophilic invertebrate assemblages, bioassessment was based on a metric battery and scoring criteria developed for montane regions of Montana (MVFP index: Bollman 1998). The six metrics constituting the bioassessment index used for MVFP sites in this study were selected because, both individually and as an integrated metric battery, they are robust at distinguishing impaired sites from relatively unimpaired sites (Bollman 1998). They have been demonstrated to be more variable with anthropogenic disturbance than with natural environmental gradients (Bollman 1998). Each of the six metrics, and their expected responses to various stressors is described below.

1. Ephemeroptera (mayfly) taxa richness. The number of mayfly taxa declines as water quality diminishes. Impairments to water quality which have been demonstrated to adversely affect the ability of mayflies to flourish include elevated water temperatures, heavy metal contamination, increased turbidity, low or high pH, elevated specific conductance and toxic chemicals. Few mayfly species are able to tolerate certain disturbances to instream habitat, such as excessive sediment deposition.
2. Plecoptera (stonefly) taxa richness. Stoneflies are particularly susceptible to impairments that affect a stream on a reach-level scale, such as loss of riparian canopy, streambank instability, channelization, and alteration of morphological features such as pool frequency and function, riffle development and sinuosity. Just as all benthic organisms, they are also susceptible to smaller scale habitat loss, such as by sediment deposition, loss of interstitial spaces between substrate particles, or unstable substrate.
3. Trichoptera (caddisfly) taxa richness. Caddisfly taxa richness has been shown to decline when sediment deposition affects habitat. In addition, the presence of certain case-building caddisflies can indicate good retention of woody debris and lack of scouring flow conditions.
4. Number of sensitive taxa. Sensitive taxa are generally the first to disappear as anthropogenic disturbances increase. The list of sensitive taxa used here includes organisms sensitive to a wide range of disturbances, including warmer water temperatures, organic or nutrient pollution, toxic pollution, sediment deposition, substrate instability and others. Unimpaired streams of western Montana typically support at least four sensitive taxa (Bollman 1998).
5. Percent filter feeders. Filter-feeding organisms are a diverse group; they capture small particles of organic matter, or organically enriched sediment material, from the water column by means of a variety of adaptations, such as silken nets or hairy appendages. In forested montane streams, filterers are expected to occur in insignificant numbers. Their abundance increases when canopy cover is lost and when water temperatures increase and the accompanying growth of filamentous algae occurs. Some filtering organisms, specifically the Arctopsyche caddisflies (*Arctopsyche* spp. and *Parapsyche* spp.) build silken nets with large mesh sizes that capture small organisms such as chironomids and early-instar mayflies. Here they are considered predators, and, in this study, their abundance does not contribute to the percent filter feeders metric.
6. Percent tolerant taxa. Tolerant taxa are ubiquitous in stream sites, but when disturbance increases, their abundance increases proportionately. The list of taxa used here includes organisms tolerant of a wide range of disturbances, including warmer water temperatures, organic or nutrient pollution, toxic pollution, sediment deposition, substrate instability and others.

Table 1. Montana Department of Transportation Mitigated Wetlands Monitoring Project sites: sampling history. Only those sites sampled in 2008 are included. An asterisk indicates lotic sites.

Site Identifier	2001	2002	2003	2004	2005	2006	2007	2008
Roundup	+	+	+	+	+	+	+	+
Hoskins Landing MS-1		+	+	+	+	+	+	+
Peterson Ranch Pond 2		+		+	+	+	+	+
Peterson Ranch Pond 4		+	+	+	+	+	+	+
Perry Ranch		+			+			+
Camp Creek MS-1*		+	+	+	+	+	+	+
Camp Creek MS-2*						+	+	+
Cloud Ranch Pond				+	+		+	+
Cloud Ranch Stream*				+			+	+
Jack Creek – Pond				+	+	+	+	+
Jack Creek – McKee*							+	+
Norem				+	+	+	+	+
Rock Creek Ranch					+	+	+	+
Wagner Marsh					+	+	+	+
Alkali Lake 1						+	+	+
West Fork of Charley Creek							+	+
Woodson Pond MI 1							+	+
Woodson Stream MI 2*							+	+
Little Muddy Creek							+	+
Selkirk Ranch							+	+
DH Ranch							+	+
Jocko Spring Creek MS-1								+
Jocko Spring Creek MS-2								+
Sportsman's Campground Site #1								+
Sportsman's Campground Site #2								+
Sportsman's Campground Site #3								+
Lonepine #1								+
Lonepine #2								+

Table 2. Aquatic invertebrate metrics employed for wetland (lentic) invertebrate assemblages in the MDT mitigated wetlands study, 2001 – 2008.

Metric	Metric Calculation	Expected response to degradation or impairment
Total taxa	Count of unique taxa identified to lowest recommended taxonomic level	Decrease
POET	Count of unique Plecoptera, Trichoptera, Ephemeroptera, and Odonata taxa identified to lowest recommended taxonomic level	Decrease
Chironomidae taxa	Count of unique midge taxa identified to lowest recommended taxonomic level	Decrease
Crustacea taxa + Mollusca taxa	Count of unique Crustacea taxa and Mollusca taxa identified to lowest recommended taxonomic level	Decrease
% Chironomidae	Percent abundance of midges in the subsample	Increase
Orthocladiinae / Chironomidae	Number of individual midges in the sub-family Orthocladiinae / total number of midges in the subsample.	Decrease
% Amphipoda	Percent abundance of amphipods in the subsample	Increase
% Crustacea + % Mollusca	Percent abundance of crustaceans in the subsample plus percent abundance of molluscs in the subsample	Increase
HBI	Relative abundance of each taxon multiplied by that taxon's modified Hilsenhoff Biotic Index (tolerance) value. These numbers are summed over all taxa in the subsample.	Increase
%Dominant taxon	Percent abundance of the most abundant taxon in the subsample	Increase
%Collector-Gatherers	Percent abundance of organisms in the collector-gatherer functional group	Decrease
%Filterers	Percent abundance of organisms in the filterer functional group	Increase

RESULTS

(Note: Individual site discussions were removed from this report by PBS&J and are included in the macroinvertebrate sections of individual monitoring reports. Summary tables for lentic (4a – 4c) and lotic (5) sites and project specific taxa listing(s) and metrics report(s) are provided on the following pages.)

Table 4a. Metric values and scores for wetland (lentic) sites in the MDT mitigated wetland study – 2008 sampling.

METRIC	Roundup	Hoskins Landing MS 1	Peterson Ranch Pond 2	Peterson Ranch Pond 4	Perry Ranch	Cloud Ranch Pond	Jack Creek Pond	Norem
Total taxa	9	18	13	25	11	27	21	14
POET	0	2	1	3	0	5	2	0
Chironomidae taxa	4	5	3	6	5	14	7	6
Crustacea + Mollusca	3	6	3	5	2	4	6	2
% Chironomidae	80.37%	17.00%	3.70%	13.21%	88.79%	49.53%	42.86%	34.69%
Orthocladiinae/Chir	0.63	0.18	1.50	0.21	0.82	0.66	0.40	0.53
% Amphipoda	0.00%	8.00%	0.00%	0.00%	0.00%	6.54%	15.24%	0.00%
% Crustacea + % Mollusca	15.89%	48.00%	86.11%	43.40%	6.54%	10.28%	30.48%	26.53%
HBI	8.01	7.62	7.85	7.40	7.37	5.94	8.17	7.61
% Dominant taxon	50.47%	27.00%	84.26%	25.47%	62.62%	13.08%	19.05%	26.53%
% Collector-Gatherers	31.78%	54.00%	87.96%	20.75%	20.56%	56.07%	65.71%	44.90%
% Filterers	2.80%	10.00%	0.00%	1.89%	0.00%	3.74%	1.90%	0.00%
Total taxa	1	3	1	5	1	5	5	1
POET	1	1	1	3	1	5	1	1
Chironomidae taxa	3	3	3	3	3	5	5	3
Crustacea + Mollusca	1	5	1	3	1	3	5	1
% Chironomidae	1	5	5	5	1	1	1	3
Orthocladiinae/Chir	5	1	5	3	5	5	3	5
% Amphipoda	5	3	5	5	5	3	3	5
% Crustacea + % Mollusca	5	3	1	3	5	5	5	5
HBI	1	1	1	3	3	5	1	1
% Dominant taxon	1	5	1	5	1	5	5	5
% Collector-Gatherers	1	3	5	1	1	3	3	1
% Filterers	3	1	3	3	3	3	3	3
Total Score	28	34	32	42	30	48	40	34
Percent of Maximum Score	46.67%	56.67%	53.33%	70.00%	50.00%	80.00%	66.67%	56.67%
Impairment Classification	poor	sub-optimal	sub-optimal	good	poor	good	sub-optimal	sub-optimal

Table 4b. Metric values and scores for wetland (lentic) sites in the MDT mitigated wetland study – 2008 sampling.

METRIC	Rock Creek Ranch	Wagner Marsh	Alkali Lake	West Fork of Charley Creek	Woodson Pond	Woodson Stream	Little Muddy Creek	Selkirk Ranch
Total taxa	23	11	10	9	13	7	14	17
POET	1	4	0	0	1	3	1	1
Chironomidae taxa	5	2	2	1	7	0	2	8
Crustacea + Mollusca	5	2	3	3	2	2	3	5
% Chironomidae	28.97%	2.83%	5.41%	0.91%	60.00%	0.00%	55.00%	23.38%
Orthoclaadiinae/Chir	0.97	0.00	0.00	0.00	0.52	0	0.64	0.33
% Amphipoda	0.00%	0.00%	0.00%	67.27%	0.00%	7.69%	0.00%	5.19%
% Crustacea + % Mollusca	28.97%	39.62%	32.43%	70.91%	25.45%	15.38%	17.00%	48.05%
HBI	6.91	7.45	8.57	8.19	8.14	4.62	6.97	7.76
% Dominant taxon	22.43%	48.11%	48.65%	67.27%	25.45%	30.77%	35.00%	32.47%
% Collector-Gatherers	30.84%	52.83%	21.62%	68.18%	86.36%	23.08%	29.00%	16.88%
% Filterers	1.87%	0.00%	0.00%	0.00%	0.00%	30.77%	0.00%	32.47%
Total taxa	5	1	1	1	1	1	1	3
POET	1	5	1	1	1	3	1	1
Chironomidae taxa	3	1	1	1	5	1	1	5
Crustacea + Mollusca	3	1	1	1	1	1	1	3
% Chironomidae	3	5	5	5	1	5	1	3
Orthoclaadiinae/Chir	5	1	1	1	5	Not Scored	5	3
% Amphipoda	5	5	5	1	5	3	5	3
% Crustacea + % Mollusca	5	3	5	1	5	5	5	3
HBI	3	3	1	1	1	5	3	1
% Dominant taxon	5	3	3	1	5	5	3	5
% Collector-Gatherers	1	3	1	3	5	1	1	1
% Filterers	3	3	3	3	3	1	3	1
Total Score	42	34	28	20	38	31	30	32
Percent of Maximum Score	70.00%	56.67%	46.67%	33.33%	63.33%	56.36%	50.00%	53.33%
Impairment Classification	good	sub-optimal	poor	poor	sub-optimal	sub-optimal	poor	sub-optimal

Table 4c. Metric values and scores for wetland (lentic) sites in the MDT mitigated wetland study – 2008 sampling.

METRIC	DH Ranch	Sportsman's Campground Site # 1	Sportsman's Campground Site # 2	Sportsman's Campground Site # 3	Lonepine # 1	Lonepine # 2
Total taxa	15	16	9	12	18	4
POET	1	1	0	0	2	0
Chironomidae taxa	6	6	3	7	12	3
Crustacea + Mollusca	2	5	3	4	1	1
% Chironomidae	52.29%	10.91%	41.18%	69.09%	81.82%	57.14%
Orthocladiinae/Chir	0.09	0.17	0.00	0.25	0.13	0.00
% Amphipoda	0.00%	24.55%	5.88%	27.27%	0.00%	0.00%
% Crustacea + % Mollusca	30.28%	83.64%	23.53%	29.09%	7.27%	42.86%
HBI	7.33	7.55	8.76	7.55	7.60	8.14
% Dominant taxon	33.03%	56.36%	29.41%	25.45%	25.45%	42.86%
% Collector-Gatherers	49.54%	20.91%	11.76%	57.27%	55.45%	28.57%
% Filterers	0.92%	63.64%	11.76%	25.45%	22.73%	42.86%
Total taxa	3	3	1	1	3	1
POET	1	1	1	1	1	1
Chironomidae taxa	3	3	3	5	5	3
Crustacea + Mollusca	1	3	1	3	1	1
% Chironomidae	1	5	3	1	1	1
Orthocladiinae/Chir	1	1	1	3	1	1
% Amphipoda	5	1	3	1	5	5
% Crustacea + % Mollusca	5	1	5	5	5	3
HBI	3	3	1	3	3	1
% Dominant taxon	5	1	5	5	5	3
% Collector-Gatherers	3	1	1	3	3	1
% Filterers	3	1	1	1	1	1
Total Score	34	24	26	32	34	22
Percent of Maximum Score	56.67%	40.00%	43.33%	53.33%	56.67%	36.67%
Impairment Classification	sub-optimal	poor	poor	sub-optimal	sub-optimal	poor

Table 5. Metric values and scores for stream (lotic) sites in the MDT mitigated wetland study – 2008 sampling.

METRIC	Camp Creek MS-1	Camp Creek MS-2	Cloud Ranch Stream	Jack Creek – McKee Spring	Jocko Spring Creek MS-1	Jocko Spring Creek MS-2
E Richness	7	5	4	1	0	1
P Richness	2	2	0	0	0	1
T Richness	4	6	5	3	2	5
Pollution Sensitive Richness	0	1	0	0	0	0
Filterer Percent	29.00%	37.00%	5.00%	40.00%	15.00%	11.00%
Pollution Tolerant Percent	5.00%	3.00%	28.00%	1.00%	62.00%	15.00%
E Richness	3	2	2	0	0	0
P Richness	2	2	0	0	0	1
T Richness	2	3	3	2	1	3
Pollution Sensitive Richness	0	1	0	0	0	0
Filterer Percent	1	0	3	0	1	1
Pollution Tolerant Percent	3	3	0	3	0	1
Total score	11	11	8	5	2	6
Percent of maximum score	61%	61%	44%	28%	11%	33%
Impairment classification	slight	slight	moderate	moderate	severe	moderate

LITERATURE CITED

- Bollman, W. 1998. Montana Valleys and Foothill Prairies Ecoregion. Master's Thesis. (M.S.) University of Montana, Missoula, Montana.
- Brandt, D. 2001. Temperature Preferences and Tolerances for 137 Common Idaho Macroinvertebrate Taxa. Report to the Idaho Department of Environmental Quality, Coeur d' Alene, Idaho.
- Caton, L. W. 1991. Improving subsampling methods for the EPA's "Rapid Bioassessment" benthic protocols. Bulletin of the North American Benthological Society, 8(3): 317-319.
- Stribling, J.B., J. Lathrop-Davis, M.T. Barbour, J.S. White, and E.W. Leppo. 1995. Evaluation of environmental indicators for the wetlands of Montana: the multimetric approach using benthic macroinvertebrates. Report to the Montana Department of Health and Environmental Science, Helena, Montana.

Taxa Listing

Project ID: MDT08PBSJ
RAI No.: MDT08PBSJ012

RAI No.: MDT08PBSJ012

Sta. Name: Wagner Marsh

Client ID:

Date Coll.: 8/8/2008

No. Jars: 1

STORET ID:

Taxonomic Name	Count	PRA	Unique	Stage	Qualifier	BI	Function
Non-Insect							
Acari	1	0.94%	Yes	Unknown		5	PR
Copepoda	1	0.94%	Yes	Unknown		8	CG
Physidae							
Physidae	41	38.68%	Yes	Unknown		8	SC
Odonata							
Coenagrionidae							
<i>Enallagma</i> sp.	4	3.77%	Yes	Larva		7	PR
Libellulidae							
Libellulidae	1	0.94%	Yes	Larva	Damaged	9	PR
Ephemeroptera							
Baetidae							
<i>Callibaetis</i> sp.	2	1.89%	Yes	Larva		9	CG
Caenidae							
<i>Caenis</i> sp.	51	48.11%	Yes	Larva		7	CG
Heteroptera							
Corixidae							
Corixidae	1	0.94%	Yes	Adult	Damaged	10	PH
Notonectidae							
Notonectidae	1	0.94%	Yes	Larva		10	PR
Chironomidae							
Chironomidae							
<i>Pseudochironomus</i> sp.	2	1.89%	Yes	Larva		5	CG
Tanypodinae	1	0.94%	Yes	Larva	Early Instar	7	PR
Sample Count	106						

Metrics Report

Project ID: MDT08PBSJ
RAI No.: MDT08PBSJ012
Sta. Name: Wagner Marsh
Client ID:
STORET ID:
Coll. Date: 8/8/2008


Abundance Measures

Sample Count: 106
Sample Abundance: 1,272.00 8.33% of sample used

Coll. Procedure:
Sample Notes:

Taxonomic Composition

Category	R	A	PRA
Non-Insect	3	43	40.57%
Odonata	2	5	4.72%
Ephemeroptera	2	53	50.00%
Plecoptera			
Heteroptera	2	2	1.89%
Megaloptera			
Trichoptera			
Lepidoptera			
Coleoptera			
Diptera			
Chironomidae	2	3	2.83%




Dominant Taxa

Category	A	PRA
Caenis	51	48.11%
Physidae	41	38.68%
Enallagma	4	3.77%
Pseudochironomus	2	1.89%
Callibaetis	2	1.89%
Tanypodinae	1	0.94%
Notonectidae	1	0.94%
Libellulidae	1	0.94%
Corixidae	1	0.94%
Copepoda	1	0.94%
Acari	1	0.94%

Functional Composition

Category	R	A	PRA
Predator	5	8	7.55%
Parasite			
Collector Gatherer	4	56	52.83%
Collector Filterer			
Macrophyte Herbivore			
Piercer Herbivore	1	1	0.94%
Xylophage			
Scraper	1	41	38.68%
Shredder			
Omnivore			
Unknown			



Metric Values and Scores

Metric	Value	BIBI	MTP	MTV	MTM
<i>Composition</i>					
Taxa Richness	11	1	0		0
Non-Insect Percent	40.57%				
E Richness	2	1		1	
P Richness	0	1		0	
T Richness	0	1		0	
EPT Richness	2		0		0
EPT Percent	50.00%		2		1
Oligochaeta+Hirudinea Percent					
Baetidae/Ephemeroptera	0.038				
Hydropsychidae/Trichoptera	0.000				
<i>Dominance</i>					
Dominant Taxon Percent	48.11%		1		0
Dominant Taxa (2) Percent	86.79%				
Dominant Taxa (3) Percent	90.57%	1			
Dominant Taxa (10) Percent	99.06%				
<i>Diversity</i>					
Shannon H (loge)	1.257				
Shannon H (log2)	1.813		1		
Margalef D	2.144				
Simpson D	0.378				
Evenness	0.134				
<i>Function</i>					
Predator Richness	5		2		
Predator Percent	7.55%	1			
Filterer Richness	0				
Filterer Percent	0.00%			3	
Collector Percent	52.83%		3		3
Scraper+Shredder Percent	38.68%		3		1
Scraper/Filterer	0.000				
Scraper/Scraper+Filterer	0.000				
<i>Habit</i>					
Burrower Richness	1				
Burrower Percent	1.89%				
Swimmer Richness	2				
Swimmer Percent	2.83%				
Clinger Richness	0	1			
Clinger Percent	0.00%				
<i>Characteristics</i>					
Cold Stenotherm Richness	0				
Cold Stenotherm Percent	0.00%				
Hemoglobin Bearer Richness	2				
Hemoglobin Bearer Percent	2.83%				
Air Breather Richness	0				
Air Breather Percent	0.00%				
<i>Voltinism</i>					
Univoltine Richness	5				
Semivoltine Richness	1	1			
Multivoltine Percent	6.60%		3		
<i>Tolerance</i>					
Sediment Tolerant Richness	0				
Sediment Tolerant Percent	0.00%				
Sediment Sensitive Richness	0				
Sediment Sensitive Percent	0.00%				
Metals Tolerance Index	3.020				
Pollution Sensitive Richness	0	1		0	
Pollution Tolerant Percent	89.62%	1		0	
Hilsenhoff Biotic Index	7.453		0		0
Intolerant Percent	0.00%				
Supertolerant Percent	44.34%				
CTQa	96.000				

Bioassessment Indices

BioIndex	Description	Score	Pct	Rating
BIBI	B-IBI (Karr et al.)	10	20.00%	
MTP	Montana DEQ Plains (Bukantis 1998)	15	50.00%	Moderate
MTV	Montana Revised Valleys/Foothills (Bollman 1998)	4	22.22%	Moderate
MTM	Montana DEQ Mountains (Bukantis 1998)	5	23.81%	Moderate

