

---

# MONTANA DEPARTMENT OF TRANSPORTATION WETLAND MITIGATION MONITORING REPORT: YEAR 2015

---

*Easton Ranch  
Park County, Montana*



Prepared for:

**MONTANA**  
**MDT**  
**DEPARTMENT OF TRANSPORTATION**  
2701 Prospect Ave  
Helena, MT 59620-1001

Prepared by:



PO Box 1133  
Bozeman, MT 59771-1133

October 2015

# **MONTANA DEPARTMENT OF TRANSPORTATION**

## **WETLAND MITIGATION MONITORING REPORT:**

**YEAR 2015**

*Easton Ranch*  
*Park County, Montana*  
Constructed: 2009

MDT Project Number STPX-0034(14)  
Control Number 4866

MFWP: SPA MDT R3-56-2008  
USACE: NWO-2006-90370-MTB

Prepared for:

**MONTANA DEPARTMENT OF TRANSPORTATION**  
2701 Prospect Ave  
Helena, MT 59620-1001

Prepared by:

**Confluence Consulting, Inc.**  
P.O. Box 1133  
Bozeman, MT 59771

October 2015

CCI Project No: MDT.006

“MDT attempts to provide accommodations for any known disability that may interfere with a person participating in any service, program, or activity of the Department of Transportation. Alternative accessible formats of this information will be provided upon request. For further information, call 406-444-7228, TTY at 800-335-7592, or Montana Relay at 711.”

## TABLE OF CONTENTS

1.	INTRODUCTION.....	1
2.	METHODS .....	6
2.1.	Hydrology .....	6
2.2.	Vegetation .....	7
2.3.	Soil .....	8
2.4.	Wetland Delineation .....	8
2.5.	Wildlife .....	9
2.6.	Functional Assessment.....	9
2.7.	Photo Documentation .....	9
2.8.	GPS Data .....	9
2.9.	Maintenance Needs.....	9
3.	RESULTS.....	10
3.1.	Hydrology .....	10
3.2.	Vegetation .....	11
3.3.	Soil .....	23
3.4.	Wetland Delineation .....	23
3.5.	Wildlife .....	24
3.6.	Functional Assessment.....	27
3.7.	Photo Documentation .....	29
3.8.	Maintenance Needs.....	29
3.9.	Current Credit Summary .....	30
4.	REFERENCES.....	37

## TABLES

Table 1. Wetland Credit Determination for the Easton Ranch Wetland Mitigation Site. ....	3
Table 2. Vegetation species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	12
Table 3. Data summary for Transect 1 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	17
Table 4. Data summary for Transect T-2 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	19
Table 5. Data summary for Transect T-3 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	21
Table 6. Total wetland acres delineated from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	24
Table 7. Wildlife species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	25
Table 8. Functions and Values of the Easton Ranch Wetland Mitigation Site from 2010 to 2015. ....	27
Table 9. Summary of wetland credits at the Easton Ranch Wetland Mitigation Site from 2010 to 2015. ....	31
Table 10. Summary of mitigation goals for Easton Ranch wetland mitigation site. ....	33
Table 11. Summary of performance standards and success criteria for Easton Ranch wetland mitigation site. ....	34

## CHARTS

Chart 1. Transect maps showing community types on Transect T-1 from start (0 feet) to finish (1072 feet in 2010 and 1376 feet from 2011 to 2015) at the Easton Ranch Wetland Mitigation Site. ....	18
Chart 2. Length of habitat types within Transect T-1 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	18
Chart 3. Transect maps showing community types on Transect T-2 from 2010 to 2015 from start (0 feet) to finish (1333 feet) at the Easton Ranch Wetland Mitigation Site. ....	19
Chart 4. Length of habitat types within Transect T-2 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	20
Chart 5. Transect maps showing community types on Transect T-3 from 2010 to 2015 from start (0 feet) to finish (751 feet in 2010 through 2014 and 732 feet in 2015) at the Easton Ranch Wetland Mitigation Site. ....	21
Chart 6. Length of habitat types within Transect T-3 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site. ....	22



## FIGURES

Figure 1. Project location of Easton Ranch Wetland Mitigation Site. ....	2
Figure 2. 2015 Monitoring Activity Locations.....	Appendix A
Figure 3. 2015 Mapped Site Features.....	Appendix A

## APPENDICES

Appendix A	Project Area Maps – Figures 2 and 3
Appendix B	2015 MDT Wetland Mitigation Site Monitoring Form 2015 USACE Wetland Determination Data Forms 2015 MDT Montana Wetland Assessment Methods Forms
Appendix C	Project Area Photographs
Appendix D	Project Plan Sheet

Cover: View looking east at transect 3 across vegetation community Type 1 at the Easton Wetland Mitigation Site.

## 1. INTRODUCTION

The 2015 Easton Ranch Wetland Mitigation Monitoring Report presents the results of the sixth year of post-construction monitoring at the Easton Ranch mitigation area. This Montana Department of Transportation (MDT) wetland mitigation site is located within Watershed 13 – Upper Yellowstone River Basin. Wetlands were developed at this location to provide compensatory mitigation for wetland impacts associated with transportation projects in the Butte District. The Easton Ranch site was selected after an extensive search of potential wetland and stream restoration sites by MDT within the Shields River Valley in cooperation with personnel from the Park Conservation District and the US Department of Agriculture (USDA) Natural Resource Conservation Service Center (NRCS) in Livingston. Figures 2 and 3 in Appendix A show the site Monitoring Activity Locations and Mapped Site Features, respectively. The MDT Mitigation Site Monitoring Form, US Army Corps of Engineers (USACE) Wetland Determination Data Forms, and the 2008 MDT Montana Wetland Assessment Forms are included in Appendix B. Project area photographs are included in Appendix C and the Project Plan Sheet is included in Appendix D.

The Easton Ranch wetland mitigation site encompasses approximately 32.65 fenced acres (this is a reduction from 34 acres previously reported as a result of aerial photography and GIS rectification) within a conservation easement area located east of the Shields River within the boundaries of the larger Easton Family Ranch, the previous landowner. The wetland mitigation project at the Easton Ranch is located in the northwest quarter of Section 32, Township 4 North, Range 9 East, Park County, Montana. The site is located approximately three miles east of US Highway 89 and four miles northeast of Wilsall (Figure 1).

Construction entailed the excavation of a series of wetland cells and a flood channel that bisects the 32.65-acre mitigation area. The primary source of wetland hydrology is groundwater supplemented by surface water from high flows associated with the Shields River. An existing irrigation diversion and delivery system was maintained to provide supplemental water to the eastern portion of the site in a flow through system. Revegetation tasks included planting cuttings and containerized shrubs, seeding wetland herbaceous species within the excavated wetland areas, and transplanting wetland plants and soils from existing wetlands to excavated areas. The wetland project was designed to increase flood storage, improve wildlife habitat, and restore riparian and wetland habitat impacted by past agricultural practices within the Shields River watershed. The project objectives include:

- Re-establish a previously existing, relic floodplain channel and associated riparian and floodplain wetland areas totaling 1.56 acres;




Figure 1. Project location of Easton Ranch Wetland Mitigation Site.



- Create approximately 25 acres of emergent, scrub/shrub and riparian wetlands by replacing existing hay fields with a variety of wetland communities that mimic habitats found in bio-referenced wetland areas located north and south of the project;
- Preserve 1.1 acres of existing scrub/shrub, forested, and palustrine emergent communities at several locations within the project area;
- Mimic old meander scars and relic flood channels within the wetland mitigation site;
- Improve water storage capacity and increase the amount of floodplain area across the site;
- Increase the amount of wildlife habitat in this reach of the Shields River.

The project credit ratios approved by the USACE are shown in Table 1.

**Table 1. Wetland Credit Determination for the Easton Ranch Wetland Mitigation Site.**

Proposed Mitigation Features	Compensatory Mitigation Type	USACE Mitigation Ratios	Acres	Final Credit Estimate (Acres)
Creation of palustrine emergent wetland via shallow excavation.	Creation	1:1	24.95	24.95
Re-establishment of relic flood channel.	Restoration (Re-establishment)	1:1	1.56	1.56
Preservation of existing shrub/scrub and palustrine emergent wetland.	Preservation	4:1	1.10	0.275
Establish a 50-foot wide upland buffer.	Upland Buffer	5:1	6.43	1.29
Project Impacts	Debit	--	--	(0.67)
<b>Total</b>	<b>Total</b>			<b>27.41</b>

The USACE approved performance standards are listed below.

1. **Wetland Characteristics:** All restored, created, enhanced, and preserved wetlands within the project limits will meet the three parameter criteria for hydrology, vegetation, and soils established for determining wetland areas as outlined in the *1987 Corps of Engineers Wetland Delineation Manual* (Environmental Laboratory 1987) and the *2010 Regional Supplement to the Corps of Engineers Manual: Western Mountains, Valleys, and Coast Region* (USACE 2010).
  - a) **Wetland Hydrology Success** will be achieved where wetland hydrology is present as per the technical guidelines in the 1987 Manual and the 2010 Regional Supplement to the Corps of Engineers Wetland Delineation Manual.
    - (i) Soil saturation will be present for at least 12.5 percent of the growing season.

- (ii) Groundwater wells will be left undisturbed within the site for the purpose of monitoring groundwater elevations during the growing season.
  - (iii) Depressional wetlands excavated into the upland areas will be monitored to determine if groundwater hydrology is filling sites and establishing vegetation communities.
  - (iv) Hydrologic success will also require that the constructed stream channel be stable in the wetlands.
- b) **Hydric Soil Success** will be achieved where hydric soil conditions are present (per the most recent Natural Resource Conservation Service (NRCS) definitions for hydric soil) or appear to be forming, the soil is sufficiently stable to prevent erosion, and the soil is able to support plant cover. Soil sampling will be conducted during the course of the monitoring period to determine if wetland areas are exhibiting characteristics of hydric soils per the 1987 Wetland Manual. Since typical hydric soil indicators may require long periods to form, a lack of distinctive hydric soil features will not be considered a failure if hydrologic and vegetation success is achieved.
- c) **Hydrophytic Vegetation Success** will be achieved through the delineation of developing wetlands utilizing the technical guidelines established in the 1987 Wetland Manual and the 2010 Regional Supplement. The following concept of “dominance”, as defined in the 1987 Manual, will be applied during future routine wetland determinations in created/restored wetlands: *“Subjectively determine the dominant species by estimating those having the largest relative basal area (woody overstory), greatest height (woody understory), greatest percentage of aerial cover (herbaceous understory), and/or greatest number of stems (woody vines).”*
  - i. **Woody Plants** – Trees and shrubs are to be installed at various locations to provide structural diversity within the site at the direction of the MDT Reclamation Specialist. Survival of woody plant species planted within the site will be evaluated to determine survival rates and success of the planting each year of the monitoring period. Success of these planted species will be determined by stem counts each year to determine survival rates of the various planted woody species and will also include the evaluation of naturally recruited woody plant species within the site. *“Scrub/shrub wetland habitat will be achieved where 30 percent absolute cover by cuttings, planted and volunteer woody plants is reached within the defined monitoring period or the site is showing signs of progression (e.g. by approximating stem densities and estimating future canopy coverage, or using other appropriate methods) towards that goal at the end of the defined monitoring period.”*



- ii. **Herbaceous Plants** – At the conclusion of the monitoring period, ocular coverage of desirable hydrophytic vegetation (wetland plants listed as OBL, FACW and FAC) will be at least 80 percent. A wetland seed mix was prepared for this site that included tufted hairgrass (*Deschampsia caespitosa*), Northwest Territory sedge (*Carex utriculata*), Baltic rush (*Juncus balticus*), American sloughgrass (*Beckmannia syzigachne*), American mannagrass (*Glyceria grandis*), and bluejoint reedgrass (*Calamagrostis canadensis*).
2. **Wetland Acreage Development** will provide 27.41 net credit acres for the project site (Table 1 and Project Plan Sheet, Appendix D).
- a) Emergent wetlands will comprise approximately 70 to 75 percent of the site.
  - b) Scrub/shrub wetland and riparian areas will comprise 15 to 20 percent of the site primarily along the proposed stream corridor and between created wetlands.
  - c) Open water will comprise approximately less than 5 percent of the total wetland area within the site after final monitoring.
3. **Floodplain Channel Restoration Success** will be evaluated in terms of revegetation and bank stability success.
- a) The floodplain channel corridor will be considered stable when banks are vegetated with a majority of deep-rooting riparian and wetland plant species.
  - b) Bank pins will be established at appropriate locations along the new relic floodplain channel to monitor channel stability and to measure channel movement.
  - c) Bank stability success will be evaluated by utilizing the bio-reference reaches to the north and south of the project area as comparisons due to their relatively undisturbed and vegetated mixture of woody and herbaceous riparian and wetland plant species.
  - d) Vegetation transects will be monitored along the relic floodplain channel corridor to determine root stability indices of the riparian and wetland plant species as it develops.
4. **Bank Stabilization Success** along the Shields River in the northwestern corner of the site will be evaluated in terms of revegetation and bank stability success.
- a) Bank stability will be achieved when the banks are vegetated with a majority of deep-rooting riparian and wetland plant species.
  - b) This area will be visually inspected and photo documented for incorporation into the annual monitoring reports to outline the success of the bank stabilization.

- c) If annual monitoring determines that the banks are eroding, the USACE and Fish, Wildlife, and Parks (FWP) will be contacted to coordinate a field meeting for joint evaluation and consultation on remediation.
- 5. **Upland Buffer Success** will be achieved when noxious weeds do not exceed 10 percent of cover within the buffer areas on site. Any area within the creditable buffer zone disturbed by project construction must have at least 50 percent aerial cover of non-weed species by the end of the monitoring period.
- 6. **Weed Control** will be based upon annual monitoring of the site to determine weed species and degree of infestation within the site, and control measures based upon the monitoring results will be implemented by MDT to minimize and/or eliminate the intrusion of State Listed Noxious weed species within the site. The MDT will manage the wetland conservation easement area to meet a goal of having less than 5 percent absolute cover of state listed noxious weed species across the site.
- 7. **Fencing** of the proposed mitigation site has been installed along the easement boundaries to protect the integrity of the wetland from disturbance that may be detrimental to the site. Fencing installed along the perimeter of the site has been designed to be “wildlife friendly” to allow for wildlife movement into and out of the wetland complex.
- 8. **Monitoring** of this MDT mitigation site will be based upon the MDT standard monitoring protocols utilized for all MDT wetland mitigation sites for a minimum period of five years or longer as determined by the US Army Corps, Montana Regulatory Office’s review of annual monitoring reports for the site and whether or not the site has met the wetland success criteria.

## 2. METHODS

The sixth year of monitoring was completed on June 26, 2015. Information for the Mitigation Monitoring Form and Wetland Determination Data Form was collected in the field (Appendix B). Monitoring activity sites were located with a global positioning system (GPS) as shown on Figure 2 (Appendix A). Information collected included a wetland delineation, vegetation community mapping, vegetation transect monitoring, soil and hydrology data collection, bird and wildlife use documentation, photographic documentation, and a non-engineering examination of the infrastructure established within the mitigation project area.

### 2.1. Hydrology

The presence of hydrological indicators as outlined on the Wetland Determination Data Form was assessed at four data points established within the project area. The hydrologic indicators were evaluated according to features observed during the site visit. The data were recorded on the electronic Wetland Determination Data Form (Appendix B). Hydrologic assessments allow evaluation of mitigation criteria addressing inundation/saturation requirements.

Technical criteria for wetland hydrology guidelines have been established as “permanent or periodic inundation, or soil saturation within 12 inches of the ground surface for a significant period (12.5 percent of the growing season) during the growing season” (USACE 2010). Systems with continuous inundation or saturation for greater than 12.5 percent of the growing season are considered jurisdictional wetlands. The growing season is defined for purposes of this report as the number of days when there is a 50 percent probability that the minimum daily temperature is greater than or equal to 28 degrees Fahrenheit (Environmental Laboratory 1987). Temperature data recorded for the meteorological station at Wilsall 8 ENE, Montana (249023) has a median (5 years in 10) growing season length of 120 days. Areas defined as wetlands would require 15 days of inundation or saturation within 12 inches of the ground surface to meet the hydrology criteria. Soil pits excavated during the wetland delineation were used to evaluate groundwater levels within 18 inches of the ground surface. The data were recorded on the Wetland Determination Data Form (Appendix B).

## **2.2. Vegetation**

The boundaries of the dominant vegetation communities were determined in the field during the active growing season and subsequently delineated on the 2015 aerial photograph. Percent cover of dominant species within a community type was visually estimated and recorded using the following classes: 0 (less than 1 percent), 1 (1 to 5 percent), 2 (6 to 10 percent), 3 (11 to 20 percent), 4 (21 to 50 percent), and 5 (greater than 50 percent) (Appendix B). Community types were named based on the dominant vegetation species that characterized each mapped polygon (Figure 3, Appendix A).

Temporal changes in vegetation were evaluated through annual assessments of static belt transects established in June, 2010 (Figure 2, Appendix A). Vegetation composition was assessed and recorded along three vegetation belt transects (T-1, T-2, T-3) approximately 10 feet wide and 1376, 1333, and 751 feet long, respectively (Figure 2, Appendix A). Transects T-2 and T-3 traverse the floodplain channel corridor and banks to provide an assessment of root stability indices of the developing riparian and wetland plant species (Figure 2, Appendix A).

The transect locations were recorded with a resource-grade GPS unit. Spatial changes in the dominant vegetation communities were recorded along the stationed transect. The percent aerial cover of each vegetation species within the belt transect was estimated using the same values and cover ranges used for the polygon data on the 2015 aerial photograph (Figure 3, Appendix B). Photographs were taken at the endpoints of each transect during the monitoring event (Appendix C).

The survival of woody species planted onsite was recorded during monitoring. Survival rates are evaluated annually. Additionally, natural recruitment of woody

species will be noted. The Montana State Noxious Weed List (July 2015), prepared by the Montana Department of Agriculture, was used to categorize weeds identified within the site. The location of noxious weeds was noted in the field and mapped on the aerial photo (Figure 3, Appendix A). The noxious weed species identified are color-coded. The locations are denoted with the symbol “x”, “▲”, or “■” representing 0 to 0.1 acre, 0.1 to 1 acre, or greater than 1 acre in extent, respectively. Cover classes are represented by T, L, M, or H, for less than 1 percent, 1 to 5 percent, 6 to 25 percent, and 26 to 100 percent, respectively.

### **2.3. Soil**

Soil information was obtained from the *Soil Survey for Park County Area* (USDA 2010) and *in situ* soil descriptions. Soil cores were excavated using a hand auger and evaluated according to procedures outlined in the 1987 Manual and the 2010 Regional Supplement. A description of the soil profile, including hydric soil indicators when present, was recorded on the Wetland Determination Data Form for each profile (Appendix B).

### **2.4. Wetland Delineation**

Waters of the U.S. including special aquatic sites and jurisdictional wetlands were delineated throughout the project area in accordance with criteria established in the 1987 Manual and the 2010 Regional Supplement to the Corps of Engineers Manual: Western Mountains, Valleys, and Coast Region (USACE 2010). In order to delineate a representative area as wetland, the technical criteria for hydrophytic vegetation, hydric soil, and wetland hydrology, as described in the 1987 USACE Manual and the 2010 Regional Supplement, must be satisfied. The name and indicator status of plant species was derived from the 2014 National Wetland Plant List (NWPL) (Lichvar *et al.* 2014). A Routine Level-2 On-site Determination Method (Environmental Laboratory 1987) was used to delineate jurisdictional areas within the project boundaries. The information was recorded on the Wetland Determination Data Form (Appendix B).

The wetland boundary was determined in the field based on changes in plant communities and/or hydrology, and changes in soil characteristics. Topographic relief boundaries within the project area were also examined and cross referenced with soil and vegetation communities as supportive information for this delineation. Vegetation composition, soil characteristics, and hydrology were assessed at likely wetland and adjacent upland locations. If all three parameters met the criteria, the area was designated as wetland and mapped by vegetation community type. If any one of the parameters did not exhibit positive wetland indicators, the area was determined to be upland unless the site was classified as an atypical situation, potential problem area, or special aquatic site, i.e., mudflat. The wetland boundary was identified on the 2015 aerial photograph. Wetland areas were GPS surveyed and calculated using geographic information system (GIS) methods.

## **2.5. Wildlife**

Observations of use by mammal, reptile, amphibian, and bird use were recorded on the Mitigation Monitoring form during the site visit. Indirect use indicators including tracks, scat, burrow, eggshells, skins, and bones were also recorded. These signs were recorded while traversing the site for other required activities. Direct sampling methods such as snap traps, live traps, and pitfall traps, were not used. A comprehensive species list of wildlife observed from 2010 through 2015 during the annual monitoring periods has been compiled.

## **2.6. Functional Assessment**

The 2008 MDT Montana Wetland Assessment Method (Berglund and McEldowney 2008) was used to evaluate functions and values on the site from 2010 to 2015. This method provides an objective means of assigning wetlands an overall rating and provides regulators a means of assessing mitigation success based on wetland functions. Functions are self-sustaining properties of a wetland ecosystem that exist in the absence of society and relate to ecological significance without regard to subjective human values (Berglund and McEldowney 2008). Field data for this assessment were collected during the site visit. Wetland Assessment Forms were completed for three separate assessment areas (AA) within mitigation site (Appendix B).

## **2.7. Photo Documentation**

Monitoring at photo points provided supplemental information documenting wetland, upland, and vegetation transect conditions; site trends; and current land uses surrounding the site. Photographs were taken at established photo points throughout the mitigation area during the site visit (Appendix C). Photo point locations were recorded with a resource grade GPS unit (Figure 2, Appendix A).

## **2.8. GPS Data**

Site features and survey points were collected with a resource grade Thales Pro Mark III GPS unit during the 2015 monitoring season. Points were collected using WAAS-enabled differential correction satellites, typically improving resolution to sub-meter accuracy. The collected data were then transferred to a personal computer, imported into GIS, and presented in Montana State Plane Single Zone NAD 83 meters. Site features and survey points that were located with GPS included fence boundaries, photograph points, transect endpoints, wetland/upland boundaries, and wetland data points.

## **2.9. Maintenance Needs**

Channels, engineered structures, fencing, and other features were examined during the site visit for obvious signs of breaching, damage, or other problems. This was a cursory examination and did not constitute an engineering-level structural inspection.



### 3. RESULTS

#### 3.1. Hydrology

Climate data from the meteorological station at Wilsall 8 ENE, Montana (249023), recorded an average annual precipitation rate of 20.2 inches from April 1957 to August 2015 (Western Region Climate Center {WRCC} 2010). The recorded annual precipitation rate was 24.15 inches (2010), 18.03 inches (2011), 16.34 inches (2012), 21.43 inches (2013), and 20.10 inches (2014). This data indicates 2010 and 2013 received above-average precipitation with 2011 and 2012 exhibiting below-average precipitation. The historic precipitation average from January to August was 15.01 inches. The precipitation totals for this same period was 17.56 inches (2010), 13.36 inches (2011), 12.41 inches (2012), 13.41 inches (2013), 18.14 inches (2014), and 13.71 inches (2015). This data set corroborates that 2010 and 2014 received above-average precipitation while indicating that precipitation prior to and during the growing season for 2011 through 2013 was below average. Precipitation during the January to August period in 2015 was also below average. The extent of surface water across the site fluctuates seasonally and is moderately driven by direct precipitation and surface runoff. Although precipitation contributes to hydrology within this site, elevated seasonal groundwater levels and overbank flows from the Shields River appear to be the principal contributors to wetland hydrology at this site. The lower than average precipitation levels over the past several years have undoubtedly decreased groundwater levels in the mitigation site and the entire Shields Valley.

The irrigation diversion system located upgradient of the wetland cells was closed during the 2010 - 2015 investigations. Per MDT communication, the area was flood irrigated in June and July of 2013. Approximately ten percent of the site was inundated with surface water during the 2015 investigation at depths ranging from 0 to 2.0 feet. The average depth was 0.2 feet and the depth at the emergent vegetation/open water boundary was 0.5 feet. Inundated areas were located within the lowest contour of the excavated depressions. Unlike the 2011 monitoring event at this site, which revealed recent scour holes, sediment deposits, wrack lines, water marks, and other signs of contemporary inundation, there have been no signs of overbank flooding from the Shields River or activation of the flood channel since 2011.

Three data points were sampled to determine the wetland/upland boundaries. There were no hydrologic indicators observed at SP-1. Data points SP-2 and SP-3 were located in areas that met the wetland criteria. No primary wetland hydrology indicators were observed at wetland data point SP-3, which was located within the preserved channel. Positive wetland hydrology indicators at this data point included the FAC-neutral test and geomorphic position. One positive primary hydrologic indicator (oxidized rhizospheres along living roots) was documented at data point SP-2, located in an excavated wetland cell in the south-central part of the site. Additional hydrological indicators observed in various wetland areas of the Easton Ranch site included surface water,

saturation, water stained leaves, drift and sediment deposits, and dry season water table. Site wide saturation and inundation levels were less than what was observed in 2013 and 2014. Decreased saturation and inundation levels in 2015 were likely a result of decreased regional precipitation rates prior to the site investigation.

The 2011 spring runoff levels and duration were high as a result of an above-average snowpack in the mountains and above average spring precipitation. The constructed flood channel through the mitigation site was activated for the first time since construction during the early part of the 2011 growing season. Fluvial geomorphic processes resulted in the initial development of scour holes, riffles, and point bars. Surface water was not present in the channel during the June 2015 site visit, likely a result of low groundwater and below-average snowpack. No areas of bank erosion were noted.

### 3.2. Vegetation

Monitoring year 2015 marked the sixth year of monitoring on the Easton Ranch wetland mitigation site. One hundred and fifty plant species have been observed site-wide since 2010 (Table 2). Vegetation plant communities were mapped and named by plant composition and dominance. The composition of each community is listed on the Mitigation Monitoring Form (Appendix B). The community boundaries are shown on Figure 3 in Appendix A.

Three upland and six wetland community types were observed on the site in 2015. The upland communities were Type 1 – *Phleum pratense*/*Poa pratensis*, Type 13 – *Bromus* spp./*Trifolium* spp., Type 10 – *Bromus inermis*/*Populus tremuloides* and the wetland communities include Type 3 – *Carex* spp., Type 4 – *Salix drummondiana*, Type 5 – *Populus balsamifera*, Type 7 – Aquatic Macrophytes, Type 11 – *Juncus* spp., and Type 12 – *Eleocharis palustris*/*Typha latifolia*. These communities are discussed below.

Upland community Type 1 – *Phleum pratense*/*Poa pratensis* was identified on 8.25 acres of higher elevation areas that surround the constructed wetland cells and channel (Figure 3, Appendix A). The 0.5 acre decrease in area was due to the update of the project boundary to correspond with the most recent rectified aerial imagery. The community was dominated by herbaceous species including common Timothy (*Phleum pratense*), Kentucky bluegrass (*Poa pratensis*), smooth brome (*Bromus inermis*), caraway (*Carum carvi*), orchard grass (*Dactylis glomerata*), and common dandelion (*Taraxacum officinale*).

Wetland community Type 3 – *Carex* species (spp.) encompassed 0.44 acres in the pre-existing emergent wetlands located at the north and southwest boundaries of the site. The community included a diverse mix of wetland species including Northwest Territory sedge (beaked sedge, *Carex utriculata*), leafy tussock sedge (*Carex aquatilis*), lamp rush (*Juncus effusus*), narrow-leaf willow (*Salix exigua*), and red-tinged bulrush (*Scirpus microcarpus*).

**Table 2. Vegetation species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Scientific Names	Common Names	WMVC Indicator Status <sup>1</sup>
<i>Achillea millefolium</i>	Common Yarrow	FACU
<i>Agrostis gigantea</i>	Black Bent	FAC
<i>Agrostis stolonifera</i>	Spreading Bent	FAC
<i>Algae, green</i>	Algae, green	NL
<i>Alisma gramineum</i>	Narrow-Leaf Water-Plantain	OBL
<i>Alnus incana</i>	Speckled Alder	FACW
<i>Alopecurus geniculatus</i>	Marsh Meadow-Foxtail	OBL
<i>Alopecurus pratensis</i>	Field Meadow-Foxtail	FAC
<i>Alyssum alyssoides</i>	Pale Alyssum	NL
<i>Amaranthus retroflexus</i>	Red-Root	FACU
<b><i>Asclepias speciosa</i></b>	<b>Showy Milkweed</b>	<b>FAC</b>
<i>Avena fatua</i>	Wild Oats	NL
<i>Bare Ground</i>	Bare Ground	NL
<i>Bassia scoparia</i>	Mexican-Fireweed	FAC
<i>Beckmannia syzigachne</i>	American Slough Grass	OBL
<i>Brassica kaber</i>	Brassica kaber	NL
<b><i>Brassica napus</i></b>	<b>Turnip</b>	<b>NL</b>
<i>Bromus arvensis</i>	Field Brome	UPL
<i>Bromus carinatus</i>	California Brome	NL
<i>Bromus ciliatus</i>	Fringed Brome	FAC
<i>Bromus inermis</i>	Smooth Brome	FAC
<i>Bromus tectorum</i>	Cheatgrass	NL
<i>Calamagrostis canadensis</i>	Bluejoint	FACW
<i>Carduus nutans</i>	Nodding Plumeless-Thistle	UPL
<i>Carex aquatilis</i>	Leafy Tussock Sedge	OBL
<b><i>Carex aurea</i></b>	<b>Golden-Fruit Sedge</b>	<b>FACW</b>
<b><i>Carex limosa</i></b>	<b>Mud Sedge</b>	<b>OBL</b>
<i>Carex nebrascensis</i>	Nebraska Sedge	OBL
<b><i>Carex parryana</i></b>	<b>Parry's Sedge</b>	<b>FACW</b>
<i>Carex praegracilis</i>	Clustered Field Sedge	FACW
<i>Carex rostrata</i>	Swollen Beaked Sedge	OBL
<b><i>Carex scoparia</i></b>	<b>Pointed Broom Sedge</b>	<b>FACW</b>
<i>Carex sp.</i>	Sedge	NL
<i>Carex stipata</i>	Stalk-Grain Sedge	OBL
<i>Carex utriculata</i>	Northwest Territory Sedge	OBL
<i>Carex vesicaria</i>	Lesser Bladder Sedge	OBL
<i>Carum carvi</i>	Caraway	FACU
<i>Cassiope mertensiana</i>	Western Moss-Heather	FACU
<i>Chenopodium album</i>	Lamb's-Quarters	FACU
<i>Chenopodium leptophyllum</i>	Narrow-Leaf Goosefoot	FACU
<i>Cirsium arvense</i>	Canadian Thistle	FAC
<i>Cirsium douglasii</i>	Douglas' Thistle	OBL

<sup>1</sup> 2014 NWPL (Lichvar et al.)

New species identified in 2015 are bolded.

**Table 2. (Continued). Vegetation species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Scientific Names	Common Names	WMVC Indicator Status <sup>1</sup>
<i>Cirsium vulgare</i>	Bull Thistle	FACU
<i>Convolvulus arvensis</i>	Field Bindweed	NL
<i>Cornus alba</i>	Red Osier	FACW
<i>Cynoglossum officinale</i>	Gypsy-Flower	FACU
<i>Dactylis glomerata</i>	Orchard Grass	FACU
<i>Dasiphora fruticosa</i>	Golden-Hardhack	FAC
<i>Deschampsia caespitosa</i>	Tufted Hair Grass	FACW
<i>Descurainia sophia</i>	Herb Sophia	NL
<i>Dracocephalum sp.</i>	Dragonhead	NL
<i>Eleocharis palustris</i>	Common Spike-Rush	OBL
<i>Elodea sp.</i>	Waterweed	NL
<i>Elymus cinereus</i>	Great Basin Wildrye	NL
<i>Elymus repens</i>	Creeping Wild Rye	FAC
<i>Elymus sp.</i>	Wild Rye	NL
<i>Elymus trachycaulus</i>	Slender Wild Rye	FAC
<i>Epilobium ciliatum</i>	Fringed Willowherb	FACW
<i>Equisetum arvense</i>	Field Horsetail	FAC
<i>Equisetum hyemale</i>	Tall Scouring-Rush	FACW
<i>Festuca arundinacea</i>	Tall fescue	NL
<i>Festuca pratensis</i>	Meadow Fescue	NL
<i>Fragaria virginiana</i>	Virginia Strawberry	FACU
<i>Galium palustre</i>	Common Marsh Bedstraw	OBL
<i>Geum macrophyllum</i>	Large-Leaf Avens	FAC
<i>Glyceria elata</i>	Tall Manna Grass	FACW
<i>Glyceria grandis</i>	American Manna Grass	OBL
<i>Glyceria striata</i>	Fowl Manna Grass	OBL
<i>Glycyrrhiza lepidota</i>	American Licorice	FAC
<i>Helianthus annuus</i>	Common Sunflower	FACU
<i>Hordeum jubatum</i>	Fox-Tail Barley	FAC
<i>Juncus balticus</i>	Baltic Rush	FACW
<i>Juncus bufonius</i>	Toad Rush	FACW
<i>Juncus effusus</i>	Lamp Rush	FACW
<i>Juncus ensifolius</i>	Dagger-Leaf Rush	FACW
<i>Juncus nevadensis</i>	Sierran Rush	FACW
<i>Juncus sp.</i>	Rush	NL
<i>Juncus tenuis</i>	Lesser Poverty Rush	FAC
<i>Juncus torreyi</i>	Torrey's Rush	FACW
<i>Lappula occidentalis</i>	Flatspine Stickseed	NL
<i>Larix occidentalis</i>	Western Larch	FACU
<b><i>Lepidium perfoliatum</i></b>	<b>Clasping Pepperwort</b>	<b>FACU</b>
<i>Leymus cinereus</i>	Great Basin Lyme Grass	FAC
<i>Lotus corniculatus</i>	Garden Bird's-Foot-Trefoil	FAC

<sup>1</sup> 2014 NWPL (Lichvar et al.)

New species identified in 2015 are bolded.

**Table 2. (Continued). Vegetation species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Scientific Names	Common Names	WMVC Indicator Status <sup>1</sup>
<i>Lycopus asper</i>	Rough Water-Horehound	OBL
<i>Medicago lupulina</i>	Black Medick	FACU
<i>Medicago sativa</i>	Alfalfa	UPL
<i>Medicago sp.</i>	Medick/burclover	NL
<i>Melilotus officinalis</i>	Yellow Sweet-Clover	FACU
<i>Mentha arvensis</i>	American Wild Mint	FACW
<i>Mimulus guttatus</i>	Seep Monkey-Flower	OBL
<i>Myriophyllum sp.</i>	Water-Milfoil	NL
Open Water	Open Water	NL
<i>Pascopyrum smithii</i>	Western-Wheat Grass	FACU
<i>Persicaria lapathifolia</i>	Dock-Leaf Smartweed	FACW
<i>Persicaria maculosa</i>	Spotted Lady's-Thumb	FACW
<i>Phalaris arundinacea</i>	Reed Canary Grass	FACW
<i>Phleum pratense</i>	Common Timothy	FAC
<i>Plantago major</i>	Great Plantain	FAC
<i>Poa palustris</i>	Fowl Blue Grass	FAC
<i>Poa pratensis</i>	Kentucky Blue Grass	FAC
<i>Polypogon monspeliensis</i>	Annual Rabbit's-Foot Grass	FACW
<i>Populus angustifolia</i>	Narrow-Leaf Cottonwood	FACW
<i>Populus balsamifera</i>	Balsam Poplar	FAC
<i>Populus tremuloides</i>	Quaking Aspen	FACU
<b><i>Potamogeton gramineus</i></b>	<b>Grassy Pondweed</b>	<b>OBL</b>
<b><i>Potamogeton praelongus</i></b>	<b>White-Stem Pondweed</b>	<b>OBL</b>
<b><i>Potentilla anserina</i></b>	<b>Silverweed</b>	<b>OBL</b>
<i>Potentilla gracilis</i>	Graceful Cinquefoil	FAC
<i>Prunus virginiana</i>	Choke Cherry	FACU
<i>Ranunculus aquatilis</i>	White Water-Crowfoot	OBL
<i>Ranunculus sp.</i>	Buttercup	NL
<i>Rhamnus alnifolia</i>	Alder-Leaf Buckthorn	FACW
<i>Ribes lacustre</i>	Bristly Black Gooseberry	FAC
<i>Rosa woodsii</i>	Woods' Rose	FACU
<i>Rumex crispus</i>	Curly Dock	FAC
<i>Ruppia maritima</i>	Beaked Ditch-Grass	OBL
<i>Salix amygdaloides</i>	Peach-Leaf Willow	FACW
<i>Salix bebbiana</i>	Gray Willow	FACW
<i>Salix drummondiana</i>	Drummond's Willow	FACW
<i>Salix exigua</i>	Narrow-Leaf Willow	FACW
<i>Salix lasiandra</i>	Pacific Willow	FACW
<i>Salix lutea</i>	Yellow Willow	OBL
<i>Salix sp.</i>	Willow	NL
<b><i>Schedonorus pratensis</i></b>	<b>Meadow False Rye Grass</b>	<b>FACU</b>
<i>Scirpus microcarpus</i>	Red-Tinge Bulrush	OBL
<i>Scirpus pallidus</i>	Pale Bulrush	OBL
<i>Scutellaria galericulata</i>	Hooded Skullcap	OBL

<sup>1</sup> 2014 NWPL (Lichvar et al.)

New species identified in 2015 are bolded.



**Table 2. (Continued). Vegetation species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Scientific Names	Common Names	WMVC Indicator Status <sup>1</sup>
<i>Scutellaria lateriflora</i>	Mad Dog Skullcap	FACW
<i>Sinapis arvensis</i>	Corn Mustard	NL
<i>Sisymbrium altissimum</i>	Tall Hedge-Mustard	FACU
<i>Sisyrinchium idahoense</i>	Idaho Blue-Eyed-Grass	FACW
<i>Solidago canadensis</i>	Canadian Goldenrod	FACU
<b><i>Sonchus arvensis</i></b>	<b>Field Sow-Thistle</b>	<b>FACU</b>
<i>Stellaria graminea</i>	Grass-Leaf Starwort	FACU
<b><i>Symphyotrichum</i> sp.</b>	<b>Aster</b>	<b>NL</b>
<i>Taraxacum officinale</i>	Common Dandelion	FACU
<i>Thlaspi arvense</i>	Field Pennycress	UPL
<i>Tragopogon dubius</i>	Meadow Goat's-beard	NL
<i>Trifolium arvense</i>	Rabbit-foot Clover	NL
<i>Trifolium hybridum</i>	Alsike Clover	FAC
<i>Trifolium pratense</i>	Red Clover	FACU
<i>Trifolium repens</i>	White Clover	FAC
<i>Trifolium</i> sp.	Clover	NL
<i>Triglochin maritima</i>	Seaside Arrow-Grass	OBL
<i>Typha latifolia</i>	Broad-Leaf Cat-Tail	OBL
<i>Urtica dioica</i>	Stinging Nettle	FAC
<i>Verbascum thapsus</i>	Great Mullein	FACU
<i>Vicia americana</i>	American Purple Vetch	FAC
<b><i>Xanthium strumarium</i></b>	<b>Rough Cocklebur</b>	<b>FAC</b>

<sup>1</sup> 2014 NWPL (Lichvar *et al.*)

New species identified in 2015 are bolded.

Wetland community Type 4 – *Salix drummondiana* was identified in a 0.14 acres area in the northwest corner of the site near the bank of the Shields River. The area encompassed a pre-existing scrub-shrub wetland associated with an old flood channel meander scar of the Shields River. Dominant species included Drummond's willow (*Salix drummondiana*), western-wheatgrass (*Pascopyrum smithii*), and Nebraska sedge (*Carex nebrascensis*). Other wetland species identified in this community include American sloughgrass (*Beckmannia syzigachne*), bristly black gooseberry (*Ribes lacustre*), red-tinge bulrush, American mannagrass (*Glyceria grandis*), stinging nettle (*Urtica dioica*), orchard grass, and common Timothy.

Wetland community Type 5 – *Populus balsamifera* was a pre-existing undisturbed forested, scrub/shrub wetland located on 0.63 acres south of the construction area. The vegetation community was dominated by balsam poplar (*Populus balsamifera*), narrow-leaf cottonwood (*Populus angustifolia*), smooth brome, fowl mannagrass (*Glyceria striata*), gray willow (*Salix bebbiana*), red tinge bulrush, Pacific willow (*Salix lasiandra*), and blue skullcap (*Scutellaria lateriflora*).

Wetland community Type 7 – Aquatic Macrophytes was found in excavated depressions with the longest duration of surface water and supports semi-permanent open water. Two depressions were identified as Aquatic Macrophytes community across the site and were located within the southern half of the site where a higher seasonal groundwater table is sustained. The community characterized approximately 0.78 acres of the site. The wetland has been classified as an aquatic bed community since 2011, generally defined as a wetland vegetation class dominated by plants “that grow principally on or below the surface of the water for most of the growing season in almost all years (Cowardin *et al.* 1979).” The dominant species were green algae (protist), sedges (*Carex* sp.) and American sloughgrass, with lower covers of water-milfoil (*Myriophyllum* sp.), American mannagrass, and beaked ditch-grass (*Ruppia maritima*). A third depression (approximately 0.14 acres) that existed in 2014 was replaced with community Type 12.

Upland community Type 13 – *Bromus* spp./*Phleum pratense* was identified on 12.17 acres of upland located within the excavated footprint. This community replaced Community Type 8 – *Bromus* spp./*Trifolium* spp. in 2015 as primary colonizing species decreased dominance and more persistent, perennial plants increased in cover. The vegetation cover increased notably within this community between 2013 and 2015. This community increased in size by 0.4 acres since 2014, primarily a result of the contraction of adjacent wetland habitat into the lower elevations of this community. The community was dominated by smooth brome, common Timothy, Kentucky bluegrass, and creeping wild rye (*Elymus repens*). Sixty-two other species were observed at five percent or less in this community.

Upland community Type 10 – *Bromus inermis*/*Populus tremuloides* was identified on 0.22 acres of upland located along the northern boundary. This community was distinguished from Upland Type 1- *Phleum pratense*/*Poa pratensis* by an increase in smooth brome and regeneration of quaking aspen (*Populus tremuloides*) species observed in 2015. Other species observed were creeping wildrye, common dandelion, and Kentucky bluegrass.

Wetland community Type 11 – *Juncus* spp./*Glyceria* spp. was identified on 9.9 acres of the constructed depressions and floodplain channel which is a 0.5 acre decline since it was first characterized in 2014. This community replaced Community Type 6 – *Beckmannia syzigachne* as primary colonizing species decreased in dominance and more persistent, perennial plants increased in cover. The base elevation of a majority of the depressions in this community contained surface water or signs of recent inundation in 2015. This diverse community type was dominated by fowl mannagrass, Baltic rush (*Juncus balticus*), Torrey’s rush (*Juncus torreyi*), field meadow foxtail (*Alopecurus pratensis*), field horsetail (*Equisetum arvense*), and lamp rush. Sixty six other species were identified at five percent or less cover in this community.

Wetland community Type 12 – *Eleocharis palustris*/*Typha latifolia* was identified in 2015 in one 0.11 acre excavated depression within Community Type 11 – *Juncus* spp., replacing a small area of Community Type 7 – Aquatic Macrophytes as species dominance transitioned from aquatic macrophytes to common spike-rush (*Eleocharis palustris*) and broad-leaf cat-tail (*Typha latifolia*). This newly characterized community, observed in 2015, was dominated by common spike-rush, broad-leaf cat-tail, American sloughgrass, and Northwest Territory sedge.

In general, the site had a decline in hydrophytic vegetation since initial monitoring in 2010. Much of the project area has experienced reduced hydrology and is transitioning to upland vegetation communities. Community Type 11 – *Juncus* spp. (previously Type 6 – *Beckmannia syzigachne*), decreased in size from 10.43 acres in 2014 to 9.9 acres in 2015. The overall percent cover of hydrophytic vegetation in the constructed floodplain declined in 2015, reducing soil stability and protection from erosion when the channel is activated during high flows in the Shields River.

Vegetation cover was measured along three transects at the Easton Ranch Mitigation Site in 2015 (Figure 2, Appendix A). The data recorded on Transect 1 (Monitoring Forms, Appendix B) are summarized in tabular and graphical formats in Table 3 and Charts 1 and 2, respectively. The transect ends were photographed (Pages C-1 and C-2 in Appendix C). Transect T-1 extends 1,376 feet (1,072 feet in 2010 due to field error during survey) from south to north across several constructed cells east of the constructed channel. The transect intervals alternated between upland community Types 1 – *Phleum pratense*/*Poa pratensis*, 13 – *Bromus* spp./*Phleum pratense* and 10 – *Bromus inermis*/*Populus tremuloides* and wetland communities Types 7 – Aquatic macrophytes and 11 – *Juncus* spp.. Wetland community 11 replaced wetland community 6 in 2014 due to the continued development of wetland vegetation within these excavated depressions. Hydrophytic vegetation communities comprised 22.2 percent of T-1 in 2015, a decrease of 1.3 percent since 2014.

**Table 3. Data summary for Transect 1 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Monitoring Year	2010	2011	2012	2013	2014	2015
Transect Length (feet)	1072	1376	1376	1376	1376	1376
Vegetation Community Transitions along Transect	11	11	12	12	14	14
Vegetation Communities along Transect	3	4	4	4	5	5
Hydrophytic Vegetation Communities along Transect	1	2	2	2	2	2
Total Vegetative Species	33	18	34	44	53	57
Total Hydrophytic Species	15	19	20	29	33	37
Total Upland Species	18	19	14	15	20	20
Estimated % Total Vegetative Cover	65	70	80	85	85	85
Estimate % Unvegetated	35	30	20	15	15	15
% Transect Length Comprising Hydrophytic Vegetation Communities	28.0	17.0	14.7	17.0	23.5	22.2
% Transect Length Comprising Upland Vegetation Communities	70.0	83.0	82.5	83.0	76.5	77.8
% Transect Length Comprising Unvegetated Open Water	2.5	0.0	2.8	0.0	0.0	0.0
% Transect Length Comprising Mudflat	0.0	0.0	0.0	0.0	0.0	0.0




Chart 1. Transect maps showing community types on Transect T-1 from start (0 feet) to finish (1072 feet in 2010 and 1376 feet from 2011 to 2015) at the Easton Ranch Wetland Mitigation Site.





Chart 2. Length of habitat types within Transect T-1 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.

Data collected on Transect T-2 (Monitoring Form, Appendix B) are summarized in tabular and graphic formats (Table 4, Charts 3 and 4, respectively). The endpoints of Transect T-2 were photographed (Pages C-3 and C-4 in Appendix C). Wetland Types 3 and 11 and upland Types 1 and 13 were identified along this transect. Hydrophytic vegetation communities comprised 40 percent of T-2 in 2015, a slight decrease from 42 percent in 2014. An increase of one hydrophytic species, for a total of 36 species, was documented along T-2 in 2015.


**Table 4. Data summary for Transect T-2 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Monitoring Year	2010	2011	2012	2013	2014	2015
<b>Transect Length (feet)</b>	<b>1333</b>	<b>1333</b>	<b>1333</b>	<b>1333</b>	<b>1333</b>	<b>1333</b>
Vegetation Community Transitions along Transect	11	8	7	7	7	8
Vegetation Communities along Transect	4	4	4	4	4	4
Hydrophytic Vegetation Communities along Transect	2	2	2	2	2	2
Total Vegetative Species	35	38	42	45	52	54
Total Hydrophytic Species	17	22	29	32	35	36
Total Upland Species	18	16	13	13	17	18
Estimated % Total Vegetative Cover	65	75	80	85	85	85
Estimated % Unvegetated	35	25	20	15	15	15
% Transect Length Comprising Hydrophytic Vegetation Communities	38.7	41.0	39.5	38.9	41.9	40.0
% Transect Length Comprising Upland Vegetation Communities	61.3	59.0	60.5	61.1	58.1	60.0
% Transect Length Comprising Unvegetated Open Water	0.0	0.0	0.0	0.0	0.0	0.0
% Transect Length Comprising Mudflat	0.0	0.0	0.0	0.0	0.0	0.0



**Chart 3. Transect maps showing community types on Transect T-2 from 2010 to 2015 from start (0 feet) to finish (1333 feet) at the Easton Ranch Wetland Mitigation Site.**






**Chart 4. Length of habitat types within Transect T-2 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**


Transect T-3 was established west to east across the constructed cells and channel in the south half of the site (Figure 2, Appendix A). During the 2015 field survey an error in the starting point of T-3 was identified and later confirmed using GIS. The length of T-3 was corrected to 732 feet to correspond with field observations and the most recent rectified aerial imagery and the new project boundary. Transect T-3 data (Monitoring Form, Appendix B) are summarized in tabular and graphic formats (Table 5 and Charts 5 and 6, respectively). Photographs of the endpoints of Transect T-3 are located on Page C-5 in Appendix C. The transect intervals intercepted upland community Types 1 and 13 and wetland community Type 11. Hydrophytic vegetation comprised 39.6 percent of Transect T-3 in 2015, a decrease from 48.9 percent in 2014. Hydrophytic vegetation was not observed in the restored stream channel at the point that T-3 transected the channel. The loss of hydrology and the continuing drying trend for the area has resulted in a transition to an upland vegetation community within the channel.

**Table 5. Data summary for Transect T-3 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Monitoring Year	2010	2011	2012	2013	2014	2015
Transect Length (feet)	751	751	751	751	751	732
Vegetation Community Transitions along Transect	11	9	9	8	8	4
Vegetation Communities along Transect	3	3	3	3	3	3
Hydrophytic Vegetation Communities along Transect	1	1	1	1	1	1
Total Vegetative Species	24	35	33	34	39	39
Total Hydrophytic Species	11	17	20	20	24	23
Total Upland Species	13	18	13	14	15	16
Estimated % Total Vegetative Cover	65	70	80	85	85	85
Estimated % Unvegetated	35	30	20	15	15	15
% Transect Length Comprising Hydrophytic Vegetation Communities	45.0	50.0	49.1	48.9	48.9	43.3
% Transect Length Comprising Upland Vegetation Communities	55.0	50.0	50.9	51.1	51.1	56.7
% Transect Length Comprising Unvegetated Open Water	0.0	0.0	0.0	0.0	0.0	0.0
% Transect Length Comprising Mudflat	0.0	0.0	0.0	0.0	0.0	0.0



**Chart 5. Transect maps showing community types on Transect T-3 from 2010 to 2015 from start (0 feet) to finish (751 feet in 2010 through 2014 and 732 feet in 2015) at the Easton Ranch Wetland Mitigation Site.**



**Chart 6. Length of habitat types within Transect T-3 from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Thirty one infestations of Canadian thistle (*Cirsium arvense*), a Priority 2B noxious weed, were identified on site, primarily in uplands and along the site perimeter (Figure 3, Appendix A). The infestations ranged in area from less than 0.1 acres to between 0.1 and 1.0 acre. The cover classes ranged from a trace (less than 1 percent) to moderate (6 to 25 percent) cover. Canadian thistle was observed in communities 1, 3, 5, 13, 10, and 11. Thirteen infestations of gypsy-flower (*Cynoglossum officinale*) were observed on site, primarily in uplands. The size of the infestations was less than 0.1 acres with a trace (less than one percent) to five percent cover.

Several hundred cuttings and containerized vegetative materials were planted along the constructed flood channel to increase channel stability. A thorough survey of the project area was conducted in 2015 to locate and identify surviving cuttings and containerized saplings. The plants that were thriving in 2014 exhibited good vigor during the 2015 site visit. Approximately 12 red-osier dogwood (*Cornus alba*), 35 sandbar willow, 43 thin-leaf alder, and 75 willow cuttings were identified as surviving in 2015. There was an increase in the amount of woody volunteer species, primarily quaking aspen (*Populus tremuloides*) along the northern and southern project boundaries in 2015. A trace amount of narrow-leaf cottonwood (*Populus angustifolia*) was observed within the constructed cells.

### 3.3. Soil

The project site was mapped in the *Park County Soil Survey* (USDA 2010) within the Meadowcreek and rarely-flooded Nesda complexes, found on 0 to 2 percent slopes (155A). The Meadowcreek series is a somewhat poorly drained clay loam soil located on floodplains within valleys. The map unit is listed on the Montana Hydric soil list and is classified as a frigid Fluvaquentic Haplustoll. The Nesda loam (600B) is mapped in a small area at the south end of the project. The loam is a well-drained, frigid Fluventic Haplustoll that is listed on the Montana hydric soil list.

Soil test pits were excavated at three locations, all within what was originally mapped as the Meadowcreek soil series (SP-1 through SP-3, Figure 2, Appendix A). Data points SP-2 and SP-3 were located within wetland Community 11. Data point SP-1 was located within upland Community 13. The soil profile at SP-1 revealed a very dark gray (10YR 3/2) gravelly loamy sand. The soils observed at data point SP-1 had no hydric soil indicators. Soil profile SP-2 was a dark grayish brown (10YR 4/2) sandy clay loam with twenty percent red (2.5YR 4/8) redox concentrations along pore linings. This soil met the criteria for depleted below dark surface (A11), depleted matrix (F3), and classification as a hydric soil. The profile at SP-3 revealed a very dark brown (10 YR 2/2) silt loam soil to a depth of 15 inches and cobbles below that. This soil did not meet exhibit any hydric soil indicators. This may be because it is within a constructed overflow flood channel and the indicators have not developed yet. However, it is unclear how site hydrology may be influencing this site. Irrigation water had not been released onto the site as of the 2015 survey.

### 3.4. Wetland Delineation

Three data points were used to support the wetland boundary (SP-1 to SP-3, Figure 2, Appendix A and Wetland Determination Data Forms, Appendix B). Data points SP-2 and SP-3 were located within community Type 11, with data point SP-1 in community Type 13. The total wetland acreage, including pre-existing wetland, was 12.01 acres in 2015, a decrease of 0.60 acres since 2014 (Table 6). Water from the irrigation system at the northeast boundary had yet to be diverted to the site during the June 2015 site visit. The frequency and duration of surface and ground water does not appear to be sufficient to support a dominance of hydrophytic vegetation in a majority of the excavated and pre-existing wetland areas. Several wetlands within the project area decreased in size due to lack of hydrology. Wetland area was corrected along the southern boundary to reflect the most recent rectified aerial imagery. The delineation mapped 1.1 acres of pre-existing emergent and shrub/scrub wetland within the mitigation boundaries in 2015 (Figure 3, Appendix A). The pre-existing wetlands were originally defined during the baseline investigation completed in August 2001 (MDT 2008). The 2015 delineated wetland acres include 1.56 acres of the re-established flood channel (Community 11, Figure 3, Appendix A) and 9.34 acres of created wetland. Uplands account for 20.64 acres of the mitigation site.

The vegetation cover in the depressions characterized by Community 11 (wetland) decreased in 2015.

**Table 6. Total wetland acres delineated from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

Habitat	2001 (acres)	2010 (acres)	2011 (acres)	2012 (acres)	2013 (acres)	2014 (acres)	2015 (acres)
Pre-existing Wetland Area	1.10	1.10	1.10	1.10	1.10	1.10	1.10
Created Wetland Area	---	10.43	10.54	10.54	11.30	11.54	10.91
<b>Total Wetland Habitat</b>	<b>1.10</b>	<b>11.53</b>	<b>11.64</b>	<b>11.64</b>	<b>12.40</b>	<b>12.64</b>	<b>12.01</b>

### 3.5. Wildlife

A comprehensive list of bird and other wildlife species observed directly or indirectly from 2010 to 2015 is presented in Table 7. Twenty-four bird species were identified in 2015. The behaviors and habitats of all birds observed in 2015 are listed on the Mitigation Monitoring Form (Appendix B). Three white-tailed deer (*Odocoileus virginianus*) and tracks, one mountain cottontail (*Sylvilagus nuttallii*), and northern pocket gopher (*Thomomys talpoides*) burrows were observed on site in 2015. Because there is an active bald eagle (*Haliaeetus leucocephalus*) nest directly southwest of the site on the west side of the Shields River the project site is within the primary habitat zone for bald eagles (L. Urban, MDT, personal communications, 2015). Adult and juvenile bald eagles frequent the project site and were observed during the 2015 site visit.



**Table 7. Wildlife species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

COMMON NAME	SCIENTIFIC NAME
<b>AMPHIBIAN</b>	
Columbia Spotted Frog	<i>Rana luteiventris</i>
Woodhouse's Toad	<i>Bufo woodhousii</i>
<b>MAMMAL</b>	
Coyote	<i>Canis latrans</i>
Deer Mouse	<i>Peromyscus maniculatus</i>
Deer Sp.	<i>Odocoileus</i> sp.
Long-tailed Vole	<i>Microtus longicaudus</i>
Meadow Vole	<i>Microtus pennsylvanicus</i>
Moose	<i>Alces americanus</i>
<b>Mountain Cottontail</b>	<b><i>Sylvilagus nuttallii</i></b>
<b>Northern Pocket Gopher</b>	<b><i>Thomomys talpoides</i></b>
Porcupine	<i>Erethizon dorsatum</i>
Pronghorn	<i>Antilocapra americana</i>
Raccoon	<i>Procyon lotor</i>
Richardson's Ground Squirrel	<i>Spermophilus richardsonii</i>
Striped Skunk	<i>Mephitis mephitis</i>
White-footed Mouse	<i>Peromyscus leucopus</i>
<b>White-tailed Deer</b>	<b><i>Odocoileus virginianus</i></b>
<b>REPTILE</b>	
Plains Gartersnake	<i>Thamnophis radix</i>

Species identified in 2015 are listed in **bold** type.

**Table 7 (continued). Wildlife species observed from 2010 to 2015 at the Easton Ranch Wetland Mitigation Site.**

COMMON NAME	SCIENTIFIC NAME
<b>BIRD</b>	
American Coot	<i>Fulica americana</i>
American Crow	<i>Corvus brachyrhynchos</i>
American Goldfinch	<i>Spinus tristis</i>
American Kestrel	<i>Falco sparverius</i>
<b>American Robin</b>	<b><i>Turdus migratorius</i></b>
<b>American White Pelican</b>	<b><i>Pelecanus erythrorhynchos</i></b>
American Wigeon	<i>Anas americana</i>
<b>Bald Eagle</b>	<b><i>Haliaeetus leucocephalus</i></b>
Band-tailed Pigeon	<i>Patagioenas fasciata</i>
Bank Swallow	<i>Riparia riparia</i>
<b>Barn Swallow</b>	<b><i>Hirundo rustica</i></b>
Belted Kingfisher	<i>Megasceryle alcyon</i>
<b>Black-billed Magpie</b>	<b><i>Pica hudsonia</i></b>
Black-capped Chickadee	<i>Poecile atricapillus</i>
<b>Bullock's Oriole</b>	<b><i>Icterus bullockii</i></b>
Canada Goose	<i>Branta canadensis</i>
Cedar Waxwing	<i>Bombycilla cedrorum</i>
<b>Common Nighthawk</b>	<b><i>Chordeiles minor</i></b>
Eastern Kingbird	<i>Tyrannus tyrannus</i>
Golden Eagle	<i>Aquila chrysaetos</i>
Gray Catbird	<i>Dumetella carolinensis</i>
<b>Gray Partridge</b>	<b><i>Perdix perdix</i></b>
<b>Great Blue Heron</b>	<b><i>Ardea herodias</i></b>
Great Horned Owl	<i>Bubo virginianus</i>
House Wren	<i>Troglodytes aedon</i>
<b>Killdeer</b>	<b><i>Charadrius vociferus</i></b>
Lesser Yellowlegs	<i>Tringa flavipes</i>
Mallard	<i>Anas platyrhynchos</i>
<b>Marsh Wren</b>	<b><i>Cistothorus palustris</i></b>
Mountain Bluebird	<i>Sialia currucoides</i>
<b>Mourning Dove</b>	<b><i>Zenaida macroura</i></b>
<b>Northern Flicker</b>	<b><i>Colaptes auratus</i></b>
Northern Harrier	<i>Circus cyaneus</i>
Osprey	<i>Pandion haliaetus</i>
<b>Pacific Wren</b>	<b><i>Troglodytes pacificus</i></b>
<b>Red-tailed Hawk</b>	<b><i>Buteo jamaicensis</i></b>
<b>Red-winged Blackbird</b>	<b><i>Agelaius phoeniceus</i></b>
<b>Sandhill Crane</b>	<b><i>Grus canadensis</i></b>
<b>Savannah Sparrow</b>	<b><i>Passerculus sandwichensis</i></b>
<b>Song Sparrow</b>	<b><i>Melospiza melodia</i></b>
<b>Spotted Sandpiper</b>	<b><i>Actitis macularius</i></b>
<b>Tree Swallow</b>	<b><i>Tachycineta bicolor</i></b>
Turkey Vulture	<i>Cathartes aura</i>
<b>Vesper Sparrow</b>	<b><i>Poocetes gramineus</i></b>
Western Bluebird	<i>Sialia mexicana</i>
<b>Western Meadowlark</b>	<b><i>Sturnella neglecta</i></b>
Willet	<i>Tringa semipalmata</i>
<b>Wilson's Snipe</b>	<b><i>Gallinago delicata</i></b>
<b>Yellow Warbler</b>	<b><i>Dendroica petechia</i></b>
Yellow-rumped Warbler	<i>Dendroica coronata</i>

Species identified in 2015 are listed in **bold** type.

### 3.6. Functional Assessment

The 2008 MDT Montana Wetland Assessment Method (MWAM) (Berglund and McElDowney 2008) has been used to evaluate three assessment areas (AA) (Table 8 and Appendix B). The AAs were separated by Creation, Restoration, and Preservation areas of the mitigation site and are discussed below.

The Creation AA encompassed 9.34 acres of constructed palustrine, emergent wetland cells and has generated 52.77 functional units. The decrease of 0.6 acres of created wetland was due to both reduced hydrology in the system and a correction of the project boundary to correspond with the most recent rectified aerial imagery. The overall rating for the Creation AA remained at a Category III wetland characterized by low disturbance in 2015. The ratings were high for short and long term surface water storage, sediment/nutrient/toxicant removal, and production export/food chain support. The number of units and acreage are expected to increase as some areas of upland in the excavated areas (Community 13) transition to wetland habitat, provided sufficient wetland hydrology exists within the site. However, the current lack of hydrology on the area has resulted in a reduction of wetland acres and will not be sufficient enough to result in the expansion of wetland acreage.

The Restoration AA consisted of 1.56 acres of re-established flood channel. The Restoration AA (flood channel) received a Category III rating with 58.5 percent of the total possible points. There was an increase from moderate to high rating for sediment/shoreline stabilization between 2012 and 2013. Ratings were high for sediment/nutrient/toxicant removal and moderate for general wildlife habitat, flood attenuation, short and long term surface water storage, production export/food chain support, groundwater discharge/recharge, and uniqueness. The Restoration AA achieved a total of 9.13 functional units in 2015.

**Table 8. Functions and Values of the Easton Ranch Wetland Mitigation Site from 2010 to 2015.**

Function and Value Parameters from the 2008 MDT Montana Wetland Assessment Method	2010 Creation	2011 Creation	2012 Creation	2013 Creation	2014 Creation	2015 Creation
Listed/Proposed T&E Species Habitat	Low (0.0)	Low (0.1)	Low (0.1)	Low (0.1)	Low (0.0)	Low (0.0)
MTNHP Species Habitat	Mod (0.6)	Mod (0.6)	Low (0.2)	Low (0.2)	Low (0.2)	Low (0.2)
General Wildlife Habitat	Mod (0.5)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
General Fish/Aquatic Habitat	NA	NA	NA	NA	NA	NA
Flood Attenuation	Mod (0.6)	Mod (0.5)	Mod (0.5)	Mod (0.5)	Mod (0.5)	Mod (0.5)
Short and Long Term Surface Water Storage	High ( 0.9)	High ( 0.8)	High ( 0.8)	High ( 0.8)	High (0.8)	High (0.8)
Sediment/Nutrient/Toxicant Removal	Mod (0.7)	Mod (0.7)	High (0.9)	High (0.9)	High (0.9)	High (0.9)
Sediment/Shoreline Stabilization	Low (0.2)	Low (0.2)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)
Production Export/ Food Chain Support	Mod (0.5)	High (0.8)	High (0.8)	High (0.8)	High (0.8)	High (0.8)
Groundwater Discharge/Recharge	High (1.0)	High (1.0)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
Uniqueness	Low (0.2)	Low (0.3)	Mod (0.4)	Mod (0.4)	Mod (0.4)	Mod (0.4)
Recreation/Education Potential (bonus points)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)
<b>Actual Points / Possible Points</b>	<b>5.25 / 10</b>	<b>5.75 / 10</b>	<b>5.75 / 10</b>	<b>5.75 / 10</b>	<b>5.65 / 10</b>	<b>5.65 / 10</b>
<b>% of Possible Score Achieved</b>	<b>52.5%</b>	<b>57.5%</b>	<b>57.5%</b>	<b>57.5%</b>	<b>56.5%</b>	<b>56.5%</b>
<b>Overall Category</b>	<b>III</b>	<b>III</b>	<b>III</b>	<b>III</b>	<b>III</b>	<b>III</b>
<b>Acreage of Assessed Aquatic Habitats within Easement</b>	<b>8.98</b>	<b>9.09</b>	<b>9.09</b>	<b>9.74</b>	<b>9.98</b>	<b>9.34</b>
<b>Functional Units (acreage x actual points)</b>	<b>47.15</b>	<b>52.27</b>	<b>52.27</b>	<b>56.01</b>	<b>56.39</b>	<b>52.77</b>

**Table 8 (Continued). Functions and Values of the Easton Ranch Wetland Mitigation Site from 2010 to 2015.**

Function and Value Parameters from the 2008 MDT Montana Wetland Assessment Method	2010 Restoration	2011 Restoration	2012 Restoration	2013 Restoration	2014 Restoration	2015 Restoration
Listed/Proposed T&E Species Habitat	Low (0.0)	Low (0.1)	Low (0.1)	Low (0.1)	Low (0.0)	Low (0.0)
MTNHP Species Habitat	Mod (0.6)	Mod (0.6)	Low (0.2)	Low (0.2)	Low (0.2)	Low (0.2)
General Wildlife Habitat	Low (0.3)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
General Fish/Aquatic Habitat	NA	NA	NA	NA	NA	NA
Flood Attenuation	Mod (0.5)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)
Short and Long Term Surface Water Storage	Mod ( 0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)
Sediment/Nutrient/Toxicant Removal	Mod (0.6)	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)
Sediment/Shoreline Stabilization	Mod (0.6)	Mod (0.6)	Mod (0.6)	High (0.9)	High (0.9)	High (0.9)
Production Export/ Food Chain Support	Mod (0.5)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
Groundwater Discharge/Recharge	High (1.0)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)	Mod (0.7)
Uniqueness	Low (0.2)	Low (0.3)	Mod (0.4)	Mod (0.4)	Mod (0.4)	Mod (0.4)
Recreation/Education Potential (bonus points)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)
<b>Actual Points / Possible Points</b>	<b>4.95 / 10</b>	<b>5.95 / 10</b>	<b>5.65 / 10</b>	<b>5.95 / 10</b>	<b>5.85 / 10</b>	<b>5.85 / 10</b>
<b>% of Possible Score Achieved</b>	<b>49.5%</b>	<b>59.5%</b>	<b>56.5%</b>	<b>59.5%</b>	<b>58.5%</b>	<b>58.5%</b>
<b>Overall Category</b>	<b>III</b>	<b>III</b>	<b>III</b>	<b>III</b>	<b>III</b>	<b>III</b>
<b>Acreage of Assessed Aquatic Habitats within Easement</b>	<b>1.45</b>	<b>1.45</b>	<b>1.45</b>	<b>1.56</b>	<b>1.56</b>	<b>1.56</b>
<b>Functional Units (acreage x actual points)</b>	<b>7.18</b>	<b>8.63</b>	<b>8.19</b>	<b>9.28</b>	<b>9.13</b>	<b>9.13</b>

Function and Value Parameters from the 2008 MDT Montana Wetland Assessment Method	2010 Preservation	2011 Preservation	2012 Preservation	2013 Preservation	2014 Preservation	2015 Preservation
Listed/Proposed T&E Species Habitat	Low (0.0)	Low (0.1)	Low (0.1)	Low (0.1)	Low (0.0)	Low (0.0)
MTNHP Species Habitat	Mod (0.6)	Mod (0.6)	Low (0.2)	Low (0.2)	Low (0.2)	Low (0.2)
General Wildlife Habitat	High (0.9)	High (0.9)	High (0.9)	High (0.9)	High (0.9)	High (0.9)
General Fish/Aquatic Habitat	NA	NA	NA	NA	NA	NA
Flood Attenuation	Exc (1.0)	High (0.9)	Mod (0.6)	High (0.9)	High (0.9)	High (0.9)
Short and Long Term Surface Water Storage	High ( 0.8)	High ( 0.8)	High ( 0.8)	High ( 0.8)	Mod (0.6)	Mod (0.6)
Sediment/Nutrient/Toxicant Removal	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)	High (1.0)
Sediment/Shoreline Stabilization	NA	NA	NA	NA	NA	NA
Production Export/ Food Chain Support	Mod (0.7)	Exc (1.0)	Exc (1.0)	Exc (1.0)	High (0.9)	High (0.9)
Groundwater Discharge/Recharge	High (1.0)	High (1.0)	High (1.0)	High (1.0)	Mod (0.7)	Mod (0.7)
Uniqueness	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)	Mod (0.6)
Recreation/Education Potential (bonus points)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)	Low (0.05)
<b>Actual Points / Possible Points</b>	<b>6.65 / 9</b>	<b>6.95 / 9</b>	<b>6.25 / 9</b>	<b>6.55 / 9</b>	<b>5.85 / 9</b>	<b>5.85 / 9</b>
<b>% of Possible Score Achieved</b>	<b>73.9%</b>	<b>77.2%</b>	<b>69.4%</b>	<b>72.8%</b>	<b>65.0%</b>	<b>65.0%</b>
<b>Overall Category</b>	<b>II</b>	<b>II</b>	<b>II</b>	<b>II</b>	<b>III</b>	<b>II</b>
<b>Acreage of Assessed Aquatic Habitats within Easement</b>	<b>1.1</b>	<b>1.1</b>	<b>1.1</b>	<b>1.1</b>	<b>1.1</b>	<b>1.1</b>
<b>Functional Units (acreage x actual points)</b>	<b>7.32</b>	<b>7.65</b>	<b>6.88</b>	<b>7.21</b>	<b>6.44</b>	<b>6.44</b>

The 1.1 acre Preservation AA encompassed the existing forested, shrub/scrub and palustrine emergent wetlands. The existing wetland within the Preservation AA was rated as Category II with 65.0 percent of the possible points. The presence of emergent, scrub/shrub, and forested wetland types increased the structural diversity and flood attenuation ratings. Ratings were high for general wildlife habitat, flood attenuation and sediment/nutrient/toxicant removal. This AA was re-evaluated in 2014 as supporting a seasonal/intermittent water regime, a decrease from a perennial water regime recognized on previous evaluations and resulted in a decrease of actual points and functional units. The Preservation AA scored a total of 6.44 functional units in 2015.

### **3.7. Photo Documentation**

Photographs from transect end points are shown on pages C-1 to C-5 of Appendix C. Photographs taken at photo points one through seven (PP1 through PP7; Figure 2, Appendix A) during site visits in 2010, 2013, 2014, and 2015 are shown on pages C-6 to C-13 of Appendix C. Photos of the data points are included on page C-14. Photo points 4A pages C-9 and C-10 show the Shields River just outside the northwest corner of the project area from 2010, 2013, 2014, and 2015.

### **3.8. Maintenance Needs**

The irrigation diversion structure has been closed during all six site investigations. MDT is aware of the lack of water flow into the site and is working with the landowner and the irrigation district to have water diverted to the site earlier in the year. It is recommended that water be diverted into the site during the early growing season to promote increased development of hydric soils and hydrophytic vegetation within the site. Nine bird-boxes were installed at the site between 2010 and 2015. Three of the bird boxes were occupied. All fences were intact. No maintenance was required for the man-made structures.

Thirty one infestations of Canadian thistle, a Priority 2B noxious weed, were identified on site, primarily in uplands and along the site perimeter (Figure 3). The infestations ranged in area from less than 0.1 acres to between 0.1 and 1.0 acre. The cover classes ranged from a trace (<1.0 percent) to moderate (6 to 25 percent) cover. Canadian thistle was observed in communities 1, 3, 5, 13, 10, and 11. Thirteen infestations of gypsy-flower were observed on site, primarily in uplands. The size of the infestations was less than 0.1 acres with a trace (<1.0 percent) to low (1 to 5 percent) cover. MDT has an on-going weed control program and contractors sprayed the site in mid-July.

The east bank of the Shields River along the northwest corner of the Easton Ranch mitigation site remained relatively stable from project completion through the 2011 runoff event. The structural integrity of the coir-wrapped soil lifts was intact following high flows. Fine-grain deposits accumulated on the lifts as flood waters receded. The 2011 flood flows resulted in the formation of a wider base-flow channel due to a slight westward shift of the west bank, away from the site.



In early 2012, a woody debris jam was removed from the outer bend of the Shield River channel (east bank) downstream from photo point 4a, and several downed trees were removed from the cottonwood forest in the adjacent riparian zone. Removal of these stabilizing elements increased the vulnerability of the river to lateral migration. During the next high flow event (spring 2013), significant bank erosion occurred immediately upstream of photo point 4a. This erosion exposed the riprap protecting the reconstructed streambank, undermined the riprap along an approximately 85 foot long section that bank and undermined the coir wrapped soil lifts on that section, causing significant loss of soil and willow cuttings. Photos from photo point 4A on pages C-9 and C-10 document these changes.

Some re-accumulation of woody debris in the former log jam location was noted in 2014, but 2015 showed little additional accumulation and perhaps some loss of what wood had been gained the previous year. Although little additional bank erosion has been noted since the dramatic lateral cutting event of 2013, this section of bank remains exposed and vulnerable. The 2015 runoff period was fed by below-average precipitation. If some measures are not taken to provide additional stability to the outer bends of the Shields River through this reach, a future high water event may result in significant additional movement of the bank, which already threatens to capture the northwest fence corner of the project area.

### **3.9. Current Credit Summary**

Table 9 summarizes the current estimated wetland credits based on the USACE approved credit ratios (MDT 2008) and the wetland delineation completed in June 2015. Proposed mitigation included the creation of 24.95 acres of emergent and shrub/scrub wetlands, the re-establishment of a 1.56 acre flood channel, the preservation of 1.10 acres of pre-existing wetland, and the maintenance of 6.43 acres of upland buffer. Proposed wetland credits for the project site totaled 27.41 credit acres, which accounted for 0.67 acres of impacts associated with the construction of the mitigation wetland.

The 2015 delineation identified a total of 12.01 acres of wetlands within the project boundary. Approximately 9.34 acres of emergent wetland has developed to date within the constructed cells. The restored channel encompassed 1.56 acres of riverine emergent wetland. The pre-existing wetlands, which included portions of communities 3, 4 and 5, encompassed 1.1 acres. Uplands accounted for 20.64 acres of the 32.65 acre site. The current 50-foot upland buffer calculated for this site totals 11.5 acres. The expected value of 2.6 acres of upland buffer was replaced in 2015 with the GIS-calculated 50-foot upland buffer of 11.5 acres based on the existing extent of wetland development within the site. This resulted in a slight increase of credits between 2014 and 2015 although the overall extent of wetland habitat has decreased. Applying the approved USACE Mitigation ratios to each mitigation feature, a total of 12.81 acres of credit were

estimated in 2015 (Table 9), approximately 14.6 acres shy of the proposed final credit acreage.

This mitigation site has not developed wetland habitat as expected. Several of the excavated depressions that contained surface water in 2011 and 2014 were dry in 2012, 2013, and 2015, limiting the potential expansion of wetland acreage within the site (see photo sheets). The increase of wetland acreage delineated in 2014 was primarily associated with the lower topographical swales and basins and seasonal groundwater. Decreased water levels within the open-water depressions observed on site during the 2012, 2013, and 2015 field surveys were likely due to a decrease in precipitation during those years; in 2012, 2013, and 2015 precipitation was 18%, 11%, and 8% respectively, below the January to August long term average. In 2014 precipitation was 22% above the January to August long term average. It is expected that if water is not added to the site, the southern portion of the restored channel and the created wetland directly west of the channel will revert to non-wetland status. This could result in a loss of approximately 0.5 acres (0.5 estimated credits) of created and restored wetland area. There is currently a transition of hydrophytic vegetation to upland vegetation occurring in several areas of the project area which may result in a loss of even more wetland acres.

**Table 9. Summary of wetland credits at the Easton Ranch Wetland Mitigation Site from 2010 to 2015.**

Proposed Mitigation Features	Compensatory Mitigation Type	USACE Mitigation Ratios	Anticipated Final Credit Acreages	Proposed Final Wetland Credits (Acres)	2010 Wetland Acreages	2010 Estimated Credit (Acres)	2011 Wetland Acreages	2011 Estimated Credit (Acres)
Creation of palustrine emergent wetland via shallow excavation.	Creation	1:1	24.95	24.95	7.78	7.78	9.09	9.09
Re-establishment of relic flood channel.	Restoration (Re-establishment)	1:1	1.56	1.56	1.45	1.45	1.45	1.45
Preservation of existing shrub/scrub and palustrine emergent wetland.	Preservation	4:1	1.10	0.28	1.10	0.28	1.10	0.28
Establish a 50-foot wide upland buffer.	Upland Buffer	5:1	6.43	1.29	6.43*	1.29	6.43*	1.29
Project Impacts			-0.67	-0.67	-0.67	-0.67	-0.67	-0.67
<b>Total</b>				<b>27.41</b>		<b>10.12</b>		<b>11.44</b>

Proposed Mitigation Features	Compensatory Mitigation Type	2012 Wetland Acreages	2012 Estimated Credit (Acres)	2013 Wetland Acreages	2013 Credit Estimated (Acres)	2014 Wetland Acreages	2014 Credit Estimated (Acres)	2015 Wetland Acreages	2015 Credit Estimated (Acres)
Creation of palustrine emergent wetland via shallow excavation.	Creation	9.09	9.09	9.74	9.74	9.98	9.98	9.34	9.34
Re-establishment of relic flood channel.	Restoration (Re-establishment)	1.45	1.45	1.56	1.56	1.56	1.56	1.56	1.56
Preservation of existing shrub/scrub and palustrine emergent wetland.	Preservation	1.10	0.28	1.10	0.28	1.10	0.28	1.10	0.28
Establish a 50-foot wide upland buffer.	Upland Buffer	6.43*	1.29	6.43*	1.29	2.60**	0.52	11.5**	2.30
Project Impacts		-0.67	-0.67	-0.67	-0.67	-0.67	-0.67	-0.67	-0.67
<b>Total</b>			<b>11.44</b>		<b>12.19</b>		<b>11.67</b>		<b>12.81</b>

\*The upland buffer was expected to decrease as wetland areas expand within the mitigation boundary. Value presented in this table prior to 2014 (6.43ac) represented the expected extent of upland buffer once maximum wetland acreage has been achieved.

\*\*50-foot buffer calculated with GIS in 2015.

Table 10 summarizes the mitigation goals for the Easton Ranch. The Easton Ranch wetland mitigation site has shown continued progress towards achieving goals, although the targeted credit acreage has not been achieved in 2015 and will not occur without increasing hydrology throughout the footprint of the excavated areas. The site has achieved five of the six goals for this site. Although the site has developed nearly 10 acres of wetland habitat, this value falls over 50 percent short of the 25 acres originally identified as a target for wetland creation. Furthermore, expansion of the current wetland acreage is unlikely unless actions are taken to restore hydrology to the site.

Five of the mitigation goals have been achieved at this site. The constructed floodplain channel was activated during the 2011 spring runoff and resulted in development of scour holes, riffles, and point bars through natural fluvial geomorphic processes. Hydrophytic vegetation has established within the footprint of this channel. No bank erosion has been identified along the constructed channel through the course of yearly monitoring. Existing wetlands within the site have been preserved and grazing eliminated from these areas. The excavated depressions throughout the floodplain function as relic meander scars, storing surface water during periods of high flow within the Shields River. These depressional wetlands have improved the water storage capacity of the floodplain. The establishment of hydrophytic vegetation communities, preservation of existing scrub-shrub, forested, and emergent wetlands, and wildlife-friendly fencing around the site have improved wildlife habitat within the Easton Ranch wetland mitigation site.

The summary of performance standards listed in Table 11 indicates this site has not achieved the full suite of success criteria established in the mitigation plan for the Easton Ranch wetland mitigation site. All wetlands delineated within this site in 2015 met the USACE three parameter criteria for hydrology, vegetation, and soils. Groundwater has been documented filling the depressional wetlands excavated across the site. Groundwater wells established within the site during baseline evaluation had been removed during construction. Redoximorphic concentrations and other hydric characteristics have developed within the wetland soils across the site. Lower than average precipitation in 2015 and the lack of water flow onto the site has decreased hydrology of the project area. Soils disturbed during construction have developed vegetation communities and are stable with no signs of active erosion. Areas identified as wetland habitat support a prevalence of hydrophytic vegetation. Trees and shrubs planted throughout the mitigation site continue to develop and natural recruitment of aspen, willows, and cottonwoods has been documented. Approximately 165 live planted woody stems were observed in 2015. The woody plants remain small and have yet to achieve areal coverage greater than one percent site wide. The lack of woody plant growth is attributed to the lack of hydrology observed on the site.

**Table 10. Summary of mitigation goals for Easton Ranch wetland mitigation site.**

<b>Mitigation Goal for Easton Ranch</b>	<b>Goal Achieved Y/N</b>	<b>Discussion</b>
Create approximately 25 acres of new emergent, scrub/shrub and riparian wetlands by replacing existing hay fields with a variety of wetland communities that mimic habitats found in bio-reference wetland areas located north and south of the project.	N	A total of 9.34 acres of wetland habitat has been created at this site to date.
Re-establish a previously existing, relic floodplain channel and associated riparian and floodplain wetland areas totaling 1.56 acres.	Y	A 1.56-acre floodplain channel was excavated through the site. This channel was activated during peak spring runoff in 2011 with fluvial geomorphic processes resulting in scour holes, riffles, and point bars. No areas of bank erosion along this channel have developed and appears to be functioning as designed. Wetland vegetation has established within the footprint of the channel.
Preserve 1.1 acres of existing scrub/shrub, forested, and palustrine emergent communities at several locations within the project area.	Y	The 1.1 acres of existing scrub/shrub, forested, and palustrine emergent wetland communities have been preserved, livestock grazing has been eliminated, and the areas continue to exhibit wetland hydrology.
Mimic old meander scars and relic flood channels within the wetland mitigation site.	Y	Several depressional wetland areas have been constructed across the mitigation site and function as relic meander scars.
Improve water storage capacity and increase the amount of floodplain area across the site.	Y	Several depressional wetland areas have been constructed across the mitigation site and have increased the water storage capacity of the floodplain.
Increase the amount of wildlife habitat in this reach of the Shields River.	Y	Wildlife habitat has been improved and protected by excluding livestock grazing and promoting the establishment of wetland vegetation.

**Table 11. Summary of performance standards and success criteria for Easton Ranch wetland mitigation site.**

Performance Standards	Success Criteria	Criteria Achieved Y/N	Discussion
Wetland Characteristics	Meet the three parameter criteria for hydrology, vegetation, and soils as outlined in the 1987 Wetland Delineation Manual and 2010 Mountains, Valleys, Coast Region.	Y	Areas identified as wetland habitat within the mitigation site meet the three parameter criteria.
Wetland Hydrology	Soil saturation present for at least 12.5 percent of the growing season.	Y	Areas identified as wetland habitat within the mitigation site exhibit soil saturation for a minimum 12.5 percent of growing season.
	Groundwater wells will be left undisturbed within the site for the purpose of monitoring groundwater elevations during the growing season.	N	No groundwater wells remain on site. Due to construction activities, the original monitoring wells were removed from the site.
	Groundwater is filling the depressional wetlands excavated into the upland areas of the site.	Y	Indicators of groundwater filling the depressional wetlands include sparsely vegetated concave surfaces, saturation to the surface and inundation.
	Constructed stream channel is stable.	Y	The constructed floodplain channel is stable with no bank erosion identified throughout the mitigation area.
Hydric Soil	Hydric soil conditions present or appear to be forming.	Y	Hydric soil characteristics, including redoximorphic concentrations and depleted matrix, have developed throughout a majority of the constructed wetlands.
	Soil is sufficiently stable to prevent erosion.	Y	Disturbed soil is stable and does not exhibit signs of erosion.
	Soil is able to support plant cover.	Y	Plant cover has continued to develop across disturbed soils.
Hydrophytic Vegetation	Achieved when wetlands delineated as hydrophytic utilizing technical guidelines.	Y	Areas identified as wetland habitat within the mitigation site support a prevalence of hydrophytic vegetation (OBL, FACW, and FAC).
Woody Plants	Trees and shrubs will be installed and survival will be assessed.	Y	Trees and shrubs have been planted throughout the mitigation site and are assessed during each yearly monitoring visit.
	Scrub/shrub wetlands habitat will be achieved where 30 percent absolute cover by cuttings, planted and volunteer woody plants is reached within the defined monitoring period or site is showing signs of progression towards that goal at the end of the defined monitoring period.	N	Approximately 2.7 percent of the wetland areas identified within the site are dominated by woody vegetation. Planted woody species continue to survive and develop along the constructed flood channel. Natural recruitment of aspen, willows, and cottonwoods within the site continue to establish. It does not appear the site is exhibiting considerable progress toward this success criteria.
Herbaceous Plants	At least 80 percent ocular vegetation coverage by desirable hydrophytic vegetation.	Y	Desirable hydrophytic vegetation consist of greater than 80 percent of total vegetation cover within delineated wetlands.
Wetland Acreage Development	Provide 27.41 net credit acres for the project area.	N	A total of 12.81 acres of wetland credit has been generated for the mitigation site. This total includes 9.34 acres of created wetland, 1.56 acres of restored wetland, 1.10 acres of preserved wetland, establishment of a 11.5-acre upland buffer, and 0.67-acre debit from project impacts.



**Table 11 (Continued). Summary of performance standards and success criteria for Easton Ranch wetland mitigation site.**

Performance Standards	Success Criteria	Criteria Achieved Y/N	Discussion
Wetland Acreage Development	Emergent wetland habitat will be 70-75% of mitigation wetland.	N	Emergent wetland habitat comprises approximately 89% of total wetland areas delineated in 2015.
	Scrub/shrub wetland habitat will be 15-20% of wetland area.	N	Scrub/shrub wetland habitat comprises approximately 2.7% of total wetland areas delineated in 2015.
	Open water will be <5% of wetland area.	Y	Aquatic macrophytes habitat comprises approximately 8.1% of total wetland areas delineated in 2015. These inundated areas (<3-ft deep) seasonally fluctuate throughout the growing season and support a diversity of submergent and emergent vegetation. The intent of this criterion was to minimize the amount of deep open water habitat greater than 3 feet in depth.
Floodplain Channel Restoration	Considered stable when banks are vegetated with a majority of deep-rooting riparian and wetland plant species	Y	Streambanks along the constructed channel are vegetated with a diversity of deep-rooting and wetland plant species.
	Bank stability will be evaluated by reference reach comparison.	Y	Banks within the constructed floodplain channel are stable and compare to reference reach conditions with no signs of erosion or channel movement.
	Vegetation transect across the floodplain will be monitored.	Y	Vegetation transect across the floodplain has been monitored yearly and supports a prevalence of species with a root stability index greater than 6.
Bank Stabilization (Shields River)	Area visually inspected and photo documented.	Y	The results of annual inspection and photo documentation along the Shields River in the northwestern corner of the site are presented in the mitigation monitoring reports.
	Stability achieved when the banks are vegetated with a majority of deep-rooting riparian and wetland plant species.	N	The banks of the Shields River are generally dominated by upland pasture grasses. Soil lifts and the riprap installed along the bank are eroding near the NW corner of the site. Installed willow cuttings did not establish along this bank.
Upland Buffer	Noxious weeds do not exceed 10 percent cover within upland buffer area.	Y	Noxious weed cover is less than 10 percent within the upland buffer.
	Any area disturbed within creditable buffer zone must have at least 50 percent aerial cover of non-weed species by end of monitoring period.	Y	Disturbed areas have established greater than 50 percent cover by non-weed species.
Weed Control	Less than 5 percent absolute cover of state-listed noxious weed species across the site.	Y	State-listed noxious weed species across the site is less than 5 percent absolute cover.
Fencing	Install wildlife-friendly fencing along the easement boundaries.	Y	Wildlife-friendly fencing has been installed around the easement boundaries and is in good condition.
Monitoring	Monitor the site for a minimum period of five years or longer as determined by the US Army Corps.	Y	Comprehensive site monitoring has been on-going for approximately 6 years, since the completion of construction activities in 2009.

The anticipated 27.41 acres of credit development has not occurred to date; anticipated credits and 2015 calculated credits have been discussed above. To satisfy this performance standard, an additional 14.6 acres of wetland habitat would need to be created within the site. It is recommended that additional hydrology be provided to the site. Additional hydrology sources potentially available to this site include increased irrigation water diversion and a side channel of the Shields River located directly north of the site. In general, the percentages of emergent and scrub/shrub wetland habitat types fall outside the identified success criteria (Table 11). The criterion for open water to occupy less than five percent of wetland area has been achieved.

The floodplain channel is considered stable and successfully restored. The floodplain was designed to inundate during a Q2 event and was not intended to flow annually as a result of the lack of water in the system the southern portion of the channel is reverting back to non-wetland status. It is expected that this area will fluctuate in between inundations. The bank stability of the Shields River in the northwestern corner of the site has been considered marginal as the vegetation established along the banks primarily consist of upland pasture grasses lacking deep-binding roots. The streambank has significant erosion and the underlying rip-rap is now exposed at the downstream end of the bank causing an eddy pool to form. The willow cuttings installed along this reach did not establish. Results of the annual inspections and photo documentation have been reported in the annual monitoring reports. The disturbed upland buffer has developed greater than 50 percent cover by non-weed species and noxious weed cover is less than 10 percent. The percent cover of bare ground has decreased notably across the site from 2010 to 2015. Absolute cover of state-listed noxious weed species across the site is less than five percent. Weed management by MDT is ongoing. The fencing around the site was intact and in good condition and grazing has been excluded from the mitigation area. This 2015 wetland mitigation monitoring report represents the sixth year of post-construction monitoring at this site.

#### 4. REFERENCES

- Berglund, J. and R. McEldowney. 2008. *MDT Montana Wetland Assessment Method*. Prepared for Montana Department of Transportation, Helena, Montana. Post, Buckley, Schuh, & Jernigan, Helena, Montana. 42pp.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. FWS/OBS-79/31. U.S.D.I Fish and Wildlife Service. Washington D.C.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. U.S. Army Corps of Engineers. Washington, DC.
- Lichvar, R.W., M. Butterwick, N.C. Melvin, and W.N. Kirchner. 2014. *The National Wetland Plant List. 2014 Update of Wetland Ratings*. Phytoneuron 2014-41:1-42.
- Montana Department of Transportation, 2008 Easton Family Ranch Wetland Mitigation Plan, Watershed #13 – Upper Yellowstone River Basin, Park County, Montana
- Reed, P.B. 1988. *National list of plant species that occur in wetlands: North West (Region 9)*. Biological Report 88(26.9), May 1988. U.S. Fish and Wildlife Service, Washington, DC.
- U.S. Army Corps of Engineers. 2010. *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region (Version 2.0)*, ed. J. S. Wakeley, R. W. Lichvar, and C. V. Noble. ERDC/EL TR-10-3. Vicksburg, MS: U.S. Army Engineer Research and Development Center.

#### Websites:

- Montana Department of Agriculture website. Montana Noxious Weed List. Accessed in October 2015 at <http://agr.mt.gov/agr/Programs/Weeds/PDF/2015WeedList.pdf>
- Montana Natural Heritage Program website. Accessed in September 2015 at [http://mtnhp.org/nwi/PUB\\_PAB.asp](http://mtnhp.org/nwi/PUB_PAB.asp).
- USDA, Natural Resources Conservation Service Web Soil Survey. Park County, Montana. Accessed August 2015 at: <http://websoilsurvey.nrcs.usda.gov/app/>
- WRCC United States Historical Climatology Network. 2010. Accessed September 2015 at: <http://www.wrcc.dri.edu/CLIMATEDATA.html>.

## **Appendix A**

---

### Project Area Maps – Figures 2 and 3

---

MDT Wetland Mitigation Monitoring  
Easton Ranch  
Park County, Montana







Figure 2: 2015 Monitoring Activity Locations



- Legend**
- Vegetation Transect
  - Monitoring Limits
  - ⊕ Data Points
  - Photo Points
  - ▲ Bird Box

Base Photography Date:  
June 27, 2015

GRAPHICAL REPRESENTATION MAY OR MAY NOT DEPICT THE LEGAL DESCRIPTION OF ANY PARCEL HEREIN. THIS FIGURE IS A VISUAL AID ONLY; BOUNDARY RESTORATION MUST BE MADE BY A LICENSED LAND SURVEYOR. THIS FIGURE IS INTENDED TO DISPLAY INFORMATION RELEVANT TO THE REFERENCED REPORT. CONFLUENCE MAKES NO REPRESENTATION OR WARRANTY OF ANY KIND REGARDING THIS DRAWING FOR ANY USE OTHER THAN THE ORIGINAL. ANY OTHER USE IS AT THE USER'S SOLE RISK.

 CONFLUENCE consulting incorporated			REV -		
Figure 2					
DRAWN FM	CHECKED JJ	APPROVED XX	SCALE: As Shown		
			Drawn: October 11, 2015		
			PROJ MGR: J Johnson		
			2015 Monitoring Activity Locations		
			Easton Ranch Wetland Mitigation		
			Project Name		
			LOCATION: Park Co., MT		
			PROJECT NO: NH-STPP 5(39)		
			FILE: Easton/Monitor2015.mxd		







Figure 3: 2015 Mapped Site Features

Noxious Weeds  
*Cynoglossum officinale*  
*Cirsium arvense*

Infestation Size  
X = <0.1 acre  
▲ = 0.1 to 1 acre  
■ = 1 to 5 acre

Cover Class  
T = Trace (<1% cover)  
L = Low (1-5% cover)  
M = Moderate (6-25% cover)



- Vegetation Community Types**
- ① *Phleum pratense/Poa pratensis*
  - ③ *Carex spp.*
  - ④ *Salix drummondiana*
  - ⑤ *Populus balsamifera*
  - ⑦ *Aquatic Macrophytes*
  - ⑧ *Bromus spp./Trifolium spp.*
  - ⑩ *Bromus inermis/Populus tremuloides*
  - ⑪ *Juncus spp./Glyceria spp.*
  - ⑫ *Eleocharis palustris/Typha latifolia*
  - ⑬ *Bromus spp./Phleum pratense*

**Legend**

Monitoring Limits ———

Wetland Limits ———

Vegetation Communities ———

Base Photography Date:  
June 27, 2015

Acreages	
Project Area	32.65 acres
Gross Wetland	12.01 acres
Pre-existing Wetland	1.10 acres
Net Wetlands	10.91 acres
Uplands	20.64 acres

GRAPHICAL REPRESENTATION MAY OR MAY NOT DEPICT THE LEGAL DESCRIPTION OF ANY PARCEL HEREIN. THIS FIGURE IS A VISUAL AID ONLY; BOUNDARY RESTORATION MUST BE MADE BY A LICENSED LAND SURVEYOR. THIS FIGURE IS INTENDED TO DISPLAY INFORMATION RELEVANT TO THE REFERENCED REPORT. CONFLUENCE MAKES NO REPRESENTATION OR WARRANTY OF ANY KIND REGARDING THIS DRAWING FOR ANY USE OTHER THAN THE ORIGINAL. ANY OTHER USE IS AT THE USER'S SOLE RISK.

LOCATION: Park Co., MT		PROJECT NO: STPX-0034(14)		FILE: Easton/Veg2015.mxd	
Project Name EASTON RANCH		Drawing Title WETLAND MITIGATION		2015 MAPPED SITE FEATURES	
DRAWN FM	CHECKED JJ	APPROVED XX	SCALE: Noted	Drawn: October 11, 2015	PROJ MGR: F McNew
		Figure 3		REV -	



## **Appendix B**

---

2015 MDT Wetland Mitigation Site Monitoring Form  
2015 USACE Wetland Determination Data Form  
2015 MDT Montana Wetland Assessment Form

---

MDT Wetland Mitigation Monitoring  
Easton Ranch  
Park County, Montana

## MDT WETLAND MITIGATION SITE MONITORING FORM

Project Site: Easton Ranch Assessment Date/Time 6/26/2015

Person(s) conducting the assessment: R McEldowney, J Johnson, R Quire, F McNew

Weather: clear, 70F Location: Easton Ranch Mitigation Site

MDT District: Butte Milepost: NA

Legal Description: T 4N R 9E Section(s) NW 1/4 Sec 32

Initial Evaluation Date: 8/25/2010 Monitoring Year: 6 #Visits in Year: 1

Size of Evaluation Area: 34 (acres)

Land use surrounding wetland:

Agriculture (hay) to the east; undeveloped riparian corridor to the west, and herbaceous scrub/shrub wetland to the north and south. The majority of the area directly east of the project site is forested riparian associated with the Spring Creek corridor.

### HYDROLOGY

Surface Water Source: High groundwater; periodic overbank flow from Shields River.

Inundation: ☒ Average Depth: 0.2 (ft) Range of Depths: 0-2.0 (ft)

Percent of assessment area under inundation: 10 %

Depth at emergent vegetation-open water boundary: 0.5 (ft)

If assessment area is not inundated then are the soils saturated within 12 inches of surface: No

Other evidence of hydrology on the site (ex. – drift lines, erosion, stained vegetation, etc.):

Drift and sediment deposits (relic from 2011), geomorphic position, FAC-neutral, surface water, saturation, water stained leaves, dry season water table, reduced iron present, oxidized rhizospheres along living roots

### Groundwater Monitoring Wells

Record depth of water surface below ground surface, in feet.

Well ID	Water Surface Depth (ft)
---------	--------------------------

No Wells

#### Additional Activities Checklist:

- ☒ Map emergent vegetation-open water boundary on aerial photograph.
- ☒ Observe extent of surface water during each site visit and look for evidence of past surface water elevations (drift lines, erosion, vegetation staining, etc.)
- ☐ Use GPS to survey groundwater monitoring well locations, if present.

#### Hydrology Notes:

Areas of inundation within excavated depressions (created wetland AA). No signs of overbank flooding in 2015. It did not appear that irrigation water had yet to be turned on to the site in June site visit.

## VEGETATION COMMUNITIES

**Site** Easton Ranch

(Cover Class Codes **0** = < 1%, **1** = 1-5%, **2** = 6-10%, **3** = 11-20%, **4** = 21-50% , **5** = >50% )

**Community #** 1 **Community Type:** Phleum pratense / Poa pratensis **Acres** 8.25

Species	Cover class	Species	Cover class
Agrostis gigantea	0	Alopecurus pratensis	0
Alyssum alyssoides	0	Bare Ground	0
Brassica napus	0	Bromus carinatus	0
Bromus inermis	3	Carex aurea	0
Carex limosa	0	Carex nebrascensis	0
Carex parryana	0	Carum carvi	1
Chenopodium album	1	Cirsium arvense	0
Cirsium vulgare	0	Cynoglossum officinale	0
Dactylis glomerata	1	Deschampsia caespitosa	0
Elymus cinereus	0	Elymus repens	0
Elymus trachycaulus	0	Equisetum arvense	0
Equisetum hyemale	0	Festuca arundinacea	0
Juncus balticus	0	Juncus bufonius	0
Juncus effusus	0	Lotus corniculatus	1
Lycopus asper	0	Medicago lupulina	0
Melilotus officinalis	1	Phalaris arundinacea	0
Phleum pratense	5	Poa pratensis	3
Populus tremuloides	1	Potentilla anserina	0
Potentilla gracilis	0	Ranunculus sp.	0
Rumex crispus	1	Salix exigua	0
Salix lutea	0	Salix sp.	0
Schedonorus pratensis	0	Sisyrinchium idahoense	0
Solidago canadensis	0	Taraxacum officinale	1
Thlaspi arvense	0	Tragopogon dubius	0
Trifolium pratense	0	Trifolium repens	0

**Comments:**

**Community #** 3 **Community Type:** Carex spp. / **Acres** 0.44

Species	Cover class	Species	Cover class
Agrostis gigantea	0	Asclepias speciosa	0
Carex aquatilis	2	Carex nebrascensis	0
Carex utriculata	5	Cirsium arvense	0
Juncus balticus	1	Juncus effusus	1
Medicago lupulina	0	Mentha arvensis	0
Phleum pratense	0	Poa palustris	0
Populus tremuloides	0	Salix exigua	1
Scirpus microcarpus	1	Taraxacum officinale	0
Trifolium pratense	0		

**Comments:**

**Community #** 4 **Community Type:** Salix drummondiana / **Acres** 0.14

Species	Cover class	Species	Cover class
Beckmannia syzigachne	2	Carex nebrascensis	3
Carex praegracilis	1	Dactylis glomerata	2
Glyceria grandis	2	Mentha arvensis	1
Pascopyrum smithii	3	Phleum pratense	2
Poa pratensis	1	Ribes lacustre	2
Rosa woodsii	1	Salix bebbiana	1
Salix drummondiana	4	Scirpus microcarpus	2
Urtica dioica	2		

**Comments:**

**Community #** 5 **Community Type:** Populus balsamifera / **Acres** 0.63

Species	Cover class	Species	Cover class
Bromus inermis	2	Cirsium arvense	1
Glyceria striata	2	Populus angustifolia	4
Populus balsamifera	4	Salix bebbiana	2
Salix lasiandra	2	Scirpus microcarpus	2
Scutellaria lateriflora	2		

**Comments:**



**Community #** 7 **Community Type:** Aquatic macrophytes / **Acres** 0.78

Species	Cover class	Species	Cover class
Agrostis stolonifera	0	Algae, green	1
Alisma gramineum	0	Beckmannia syzigachne	1
Carex sp.	1	Eleocharis palustris	0
Glyceria grandis	0	Juncus balticus	0
Juncus ensifolius	0	Myriophyllum sp.	0
Open Water	4	Rumex crispus	0
Ruppia maritima	0		

**Comments:**

**Community #** 10 **Community Type:** Bromus inermis / Populus tremuloides **Acres** 0.22

Species	Cover class	Species	Cover class
Alopecurus pratensis	0	Bromus inermis	5
Carum carvi	1	Cirsium arvense	0
Cynoglossum officinale	0	Elymus repens	2
Equisetum arvense	0	Phleum pratense	0
Poa pratensis	2	Populus tremuloides	3
Salix sp.	0	Taraxacum officinale	2
Thlaspi arvense	0	Trifolium pratense	1

**Comments:**

**Community #** 11 **Community Type:** Juncus spp. / Glyceria spp.

**Acres**

9.9

Species	Cover class	Species	Cover class
Agrostis gigantea	0	Algae, green	0
Alisma gramineum	1	Alnus incana	0
Alopecurus pratensis	2	Bare Ground	0
Beckmannia syzigachne	0	Brassica kaber	1
Brassica napus	0	Bromus carinatus	0
Bromus inermis	0	Carex aquatilis	0
Carex nebrascensis	0	Carex praegracilis	0
Carex scoparia	0	Carex sp.	0
Carex stipata	1	Carex utriculata	0
Carum carvi	1	Cirsium arvense	0
Cirsium vulgare	0	Cynoglossum officinale	0
Deschampsia caespitosa	0	Eleocharis palustris	0
Elymus repens	0	Elymus trachycaulus	0
Epilobium ciliatum	0	Equisetum arvense	2
Equisetum hyemale	0	Festuca arundinacea	0
Geum macrophyllum	0	Glyceria grandis	1
Glyceria striata	3	Hordeum jubatum	0
Juncus balticus	2	Juncus bufonius	0
Juncus effusus	2	Juncus ensifolius	0
Juncus torreyi	1	Lotus corniculatus	0
Lycopus asper	0	Medicago lupulina	0
Melilotus officinalis	0	Mentha arvensis	1
Open Water	1	Phalaris arundinacea	0
Phleum pratense	1	Plantago major	1
Poa palustris	1	Poa pratensis	0
Populus angustifolia	0	Populus tremuloides	0
Potentilla anserina	0	Potentilla gracilis	0
Ranunculus aquatilis	0	Rumex crispus	1
Salix amygdaloides	0	Salix bebbiana	0
Salix exigua	0	Salix lutea	0
Salix sp.	0	Scirpus microcarpus	0
Scutellaria galericulata	0	Sisyrinchium idahoense	0
Sonchus arvensis	0	Taraxacum officinale	1
Thlaspi arvense	0	Trifolium pratense	0
Trifolium repens	1	Typha latifolia	1
Vicia americana	0		

**Comments:**

**Community #** 12 **Community Type:** Eleocharis palustris / Typha latifolia **Acres** 0.11

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Beckmannia syzigachne	2
Carex utriculata	2	Cirsium arvense	1
Eleocharis palustris	4	Glyceria elata	1
Mentha arvensis	1	Phalaris arundinacea	0
Plantago major	1	Rumex crispus	1
Solidago canadensis	0	Taraxacum officinale	1
Typha latifolia	3	Xanthium strumarium	0

**Comments:**

Inset of community 11, replacing one excavated depression originally characterized as community 7.

**Community #** 13 **Community Type:** Bromus spp. / Phleum pratense

**Acres** 12.17

Species	Cover class	Species	Cover class
Agrostis gigantea	1	Alopecurus pratensis	0
Alyssum alyssoides	0	Bare Ground	0
Brassica kaber	0	Brassica napus	0
Bromus arvensis	1	Bromus carinatus	0
Bromus inermis	4	Bromus tectorum	1
Carduus nutans	0	Carex aquatilis	0
Carex nebrascensis	0	Carex praegracilis	0
Carex utriculata	0	Carum carvi	1
Cirsium arvense	0	Cirsium vulgare	0
Cynoglossum officinale	0	Dactylis glomerata	0
Deschampsia caespitosa	0	Eleocharis palustris	0
Elymus cinereus	0	Elymus repens	2
Epilobium ciliatum	0	Equisetum arvense	0
Equisetum hyemale	0	Festuca arundinacea	0
Fragaria virginiana	0	Glyceria grandis	0
Glycyrrhiza lepidota	0	Hordeum jubatum	0
Juncus balticus	0	Juncus effusus	0
Lepidium perfoliatum	0	Lotus corniculatus	1
Lycopus asper	0	Medicago lupulina	1
Melilotus officinalis	0	Pascopyrum smithii	0
Persicaria lapathifolia	0	Phalaris arundinacea	0
Phleum pratense	4	Plantago major	0
Poa pratensis	3	Populus angustifolia	0
Populus balsamifera	0	Populus tremuloides	0
Potamogeton gramineus	0	Potamogeton praelongus	0
Potentilla gracilis	0	Rumex crispus	0
Salix lutea	0	Schedonorus pratensis	0
Scirpus microcarpus	1	Sinapis arvensis	0
Sisyrinchium idahoense	0	Solidago canadensis	0
Symphyotrichum sp.	0	Taraxacum officinale	0
Thlaspi arvense	0	Tragopogon dubius	0
Trifolium arvense	0	Trifolium pratense	0
Trifolium repens	0	Vicia americana	0

**Comments:**

This community replaced Community Type 8-Bromus spp./Trifolium spp. In 2015 as primary colonizing species decreased dominance and more persistent, perennial plants increased in cover. Community Type 8 replaced Community Type 2 – Chenopodium spp./Phleum pratense in 2011 as primary colonizing species decreased dominance and more persistent, perennial plants increased in cover.

**Total Vegetation Community Acreage**

**32.64**

*(Note: some area within the project bounds may be open water or other non-vegetative ground cover.)*

## VEGETATION TRANSECTS

Site: Easton Ranch Date: 6/26/2015

Transect Number: 1 Compass Direction from Start: 5

### Interval Data:

**Ending Station** 46 **Community Type:** Bromus spp. / Phleum pratense

Species	Cover class	Species	Cover class
Agrostis gigantea	2	Bromus inermis	3
Carex nebrascensis	1	Carex praegracilis	0
Carum carvi	2	Deschampsia caespitosa	1
Eleocharis palustris	0	Elymus repens	2
Juncus balticus	1	Juncus effusus	0
Phleum pratense	2	Poa pratensis	0
Taraxacum officinale	0	Trifolium pratense	0

**Ending Station** 61 **Community Type:** Juncus spp. /

Species	Cover class	Species	Cover class
Agrostis gigantea	0	Carex nebrascensis	0
Carex sp.	0	Carum carvi	1
Elymus repens	1	Equisetum arvense	0
Juncus balticus	4	Juncus effusus	4
Juncus ensifolius	2	Trifolium repens	0

**Ending Station** 101 **Community Type:** Aquatic macrophytes /

Species	Cover class	Species	Cover class
Agrostis stolonifera	1	Algae, green	1
Carex sp.	1	Eleocharis palustris	1
Glyceria grandis	1	Juncus balticus	1
Open Water	5		

**Ending Station** 134 **Community Type:** Juncus spp. /

Species	Cover class	Species	Cover class
Carex sp.	2	Cirsium vulgare	1
Deschampsia caespitosa	1	Juncus balticus	5
Juncus bufonius	0	Juncus effusus	3
Salix lutea	1		



**Ending Station** 197 **Community Type:** Phleum pratense / Poa pratensis

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Bromus inermis	3	Carex aurea	0
Carex nebrascensis	0	Cirsium arvense	2
Cirsium vulgare	1	Elymus cinereus	0
Elymus repens	3	Juncus bufonius	0
Juncus effusus	1	Lycopus asper	1
Phleum pratense	3	Poa pratensis	4
Taraxacum officinale	1	Thlaspi arvense	1
Trifolium pratense	1		

**Ending Station** 262 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Alopecurus pratensis	3	Alopecurus pratensis	1
Eleocharis palustris	2	Glyceria grandis	1
Juncus balticus	2	Juncus effusus	2
Juncus ensifolius	2	Medicago lupulina	1
Mentha arvensis	0	Open Water	4
Poa pratensis	0	Potentilla anserina	1
Salix lutea	1	Trifolium pratense	1
Typha latifolia	1		

**Ending Station** 471 **Community Type:** Bromus spp. / Phleum pratense

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Alopecurus pratensis	2	Bromus inermis	4
Carum carvi	1	Cirsium arvense	0
Cirsium vulgare	0	Dactylis glomerata	0
Deschampsia caespitosa	0	Elymus cinereus	3
Elymus cinereus	1	Elymus repens	4
Equisetum arvense	2	Medicago lupulina	1
Melilotus officinalis	0	Phleum pratense	0
Poa pratensis	3	Taraxacum officinale	2
Trifolium pratense	1		

**Ending Station** 512 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	0	Alopecurus pratensis	1
Carum carvi	1	Cirsium arvense	0
Juncus balticus	4	Juncus bufonius	1
Juncus effusus	4	Lycopus asper	0
Phleum pratense	1	Poa pratensis	1
Taraxacum officinale	0		

**Ending Station** 566 **Community Type:** Phleum pratense / Poa pratensis

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Alopecurus pratensis	4	Bromus inermis	2
Carex parryana	1	Carum carvi	1
Elymus repens	2	Equisetum arvense	0
Medicago lupulina	0	Phleum pratense	1
Poa pratensis	3	Rumex crispus	0
Taraxacum officinale	1	Thlaspi arvense	1

**Ending Station** 622 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Carum carvi	2
Cirsium arvense	0	Deschampsia caespitosa	1
Elymus repens	1	Equisetum arvense	1
Juncus balticus	1	Juncus bufonius	3
Juncus effusus	1	Lotus corniculatus	2
Melilotus officinalis	0	Mentha arvensis	1
Phleum pratense	0	Poa pratensis	1
Salix lutea	0	Taraxacum officinale	2
Trifolium pratense	2		

**Ending Station** 666 **Community Type:** Bromus spp. / Phleum pratense

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	0	Bromus inermis	2
Carum carvi	2	Cirsium arvense	2
Deschampsia caespitosa	1	Elymus repens	1
Lycopus asper	0	Medicago lupulina	1
Phalaris arundinacea	1	Phleum pratense	2
Poa pratensis	2	Potamogeton praelongus	0
Rumex crispus	1	Sisyrinchium idahoense	1
Taraxacum officinale	2	Trifolium pratense	2

**Ending Station** 722 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Alopecurus pratensis	1
Bromus inermis	1	Carex scoparia	0
Carum carvi	1	Cirsium arvense	2
Deschampsia caespitosa	1	Elymus repens	0
Juncus balticus	5	Juncus bufonius	0
Lycopus asper	0	Medicago lupulina	1
Phleum pratense	0	Poa pratensis	1
Potentilla gracilis	0	Salix exigua	0
Salix lutea	1	Sisyrinchium idahoense	0
Taraxacum officinale	1	Trifolium pratense	3

**Ending Station** 1271 **Community Type:** Bromus spp. / Phleum pratense

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Bare Ground	4	Brassica napus	1
Bromus inermis	3	Carum carvi	1
Cirsium arvense	1	Deschampsia caespitosa	0
Elymus cinereus	3	Elymus repens	4
Equisetum arvense	2	Equisetum hyemale	2
Juncus balticus	0	Lotus corniculatus	1
Medicago lupulina	1	Melilotus officinalis	2
Pascopyrum smithii	1	Phalaris arundinacea	0
Phleum pratense	3	Poa pratensis	1
Potamogeton gramineus	0	Potentilla gracilis	0
Schedonorus pratensis	0	Sisyrinchium idahoense	0
Symphyotrichum sp.	0	Taraxacum officinale	1
Thlaspi arvense	0	Tragopogon dubius	0
Trifolium pratense	4		

**Ending Station** 1324 **Community Type:** Phleum pratense / Poa pratensis

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Brassica napus	0	Bromus carinatus	1
Bromus inermis	3	Carum carvi	2
Cirsium arvense	4	Elymus cinereus	1
Elymus repens	5	Elymus trachycaulus	0
Equisetum arvense	0	Medicago lupulina	0
Phleum pratense	1	Poa pratensis	1
Taraxacum officinale	2	Thlaspi arvense	0
Trifolium pratense	2		

**Ending Station** 1376 **Community Type:** Bromus inermis / Populus tremuloides

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Alopecurus pratensis	0	Bromus inermis	4
Carum carvi	0	Cirsium arvense	1
Cynoglossum officinale	0	Elymus repens	4
Equisetum arvense	1	Phleum pratense	1
Poa pratensis	1	Populus tremuloides	3
Taraxacum officinale	1	Thlaspi arvense	1

Transect Notes:

Transect Number: 2Compass Direction from Start: 185**Interval Data:****Ending Station** 40 **Community Type:** Carex spp. /

Species	Cover class	Species	Cover class
Agrostis gigantea	0	Asclepias speciosa	0
Carex nebrascensis	5	Carex utriculata	1
Cirsium arvense	3	Juncus balticus	2
Juncus effusus	1	Mentha arvensis	0
Poa palustris	0	Scirpus microcarpus	2
Taraxacum officinale	1		

**Ending Station** 60 **Community Type:** Juncus spp. /

Species	Cover class	Species	Cover class
Agrostis gigantea	1	Bromus inermis	0
Carex nebrascensis	1	Carex scoparia	1
Carex sp.	1	Carex utriculata	1
Cirsium arvense	0	Deschampsia caespitosa	0
Eleocharis palustris	0	Epilobium ciliatum	0
Equisetum arvense	1	Glyceria grandis	1
Juncus balticus	2	Juncus bufonius	1
Juncus effusus	1	Lycopus asper	1
Mentha arvensis	0	Phleum pratense	0
Poa palustris	1	Salix lutea	0
Scutellaria galericulata	0	Sisyrinchium idahoense	0

**Ending Station** 115 **Community Type:** Bromus inermis / Phleum pratense

Species	Cover class	Species	Cover class
Bromus inermis	1	Cirsium arvense	0
Dactylis glomerata	1	Elymus cinereus	0
Elymus repens	3	Equisetum arvense	1
Juncus balticus	1	Phleum pratense	4
Poa pratensis	2	Taraxacum officinale	3

**Ending Station** 142 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	2	Carex nebrascensis	1
Carex utriculata	2	Cirsium arvense	0
Deschampsia caespitosa	1	Eleocharis palustris	0
Juncus balticus	3	Juncus effusus	2
Lotus corniculatus	1	Lycopus asper	1
Medicago lupulina	0	Mentha arvensis	1
Phleum pratense	1	Plantago major	0
Potentilla gracilis	0	Salix lutea	0
Taraxacum officinale	1	Trifolium pratense	3

**Ending Station** 315 **Community Type:** Phleum pratense / Poa pratensis

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Bromus inermis	1
Cirsium arvense	1	Elymus cinereus	0
Elymus repens	4	Equisetum arvense	1
Lotus corniculatus	0	Melilotus officinalis	1
Phalaris arundinacea	0	Phleum pratense	2
Poa pratensis	4	Potentilla anserina	0
Rumex crispus	0	Salix lutea	1
Taraxacum officinale	3	Trifolium pratense	2

**Ending Station** 351 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	2	Carex praegracilis	0
Cirsium arvense	1	Deschampsia caespitosa	1
Elymus repens	1	Juncus balticus	3
Juncus bufonius	2	Juncus effusus	1
Lotus corniculatus	0	Lycopus asper	1
Melilotus officinalis	0	Mentha arvensis	0
Phleum pratense	1	Poa pratensis	4
Potentilla anserina	1	Salix lutea	1
Scirpus microcarpus	1	Sisyrinchium idahoense	0
Trifolium pratense	1		



**Ending Station** 831 **Community Type:** Bromus spp. / Phleum pratense

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Alyssum alyssoides	0
Bromus carinatus	0	Bromus inermis	2
Carex praegracilis	0	Carum carvi	1
Cirsium arvense	1	Cirsium vulgare	0
Elymus cinereus	1	Elymus repens	4
Equisetum arvense	1	Festuca arundinacea	1
Lepidium perfoliatum	0	Lotus corniculatus	0
Medicago lupulina	0	Melilotus officinalis	2
Phalaris arundinacea	0	Phleum pratense	3
Poa pratensis	4	Rumex crispus	0
Salix lutea	0	Sisyrinchium idahoense	0
Solidago canadensis	0	Taraxacum officinale	1
Thlaspi arvense	0	Trifolium pratense	1

**Ending Station** 1239 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	2	Alnus incana	0
Alopecurus pratensis	0	Brassica napus	0
Carex nebrascensis	1	Carex praegracilis	0
Carex sp.	1	Carex utriculata	1
Cirsium arvense	1	Deschampsia caespitosa	0
Eleocharis palustris	0	Equisetum arvense	1
Festuca arundinacea	0	Glyceria grandis	0
Juncus balticus	5	Juncus bufonius	2
Lotus corniculatus	0	Lycopus asper	3
Medicago lupulina	2	Mentha arvensis	1
Poa pratensis	1	Populus angustifolia	0
Populus tremuloides	3	Potentilla anserina	1
Salix exigua	1	Salix lutea	2
Scirpus microcarpus	0	Sisyrinchium idahoense	0
Sonchus arvensis	0	Trifolium pratense	0

**Ending Station** 1326 **Community Type:** Phleum pratense / Poa pratensis

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	3	Alyssum alyssoides	0
Bromus inermis	4	Cirsium arvense	0
Elymus cinereus	0	Elymus repens	3
Equisetum arvense	1	Lotus corniculatus	0
Phleum pratense	1	Poa pratensis	1
Populus tremuloides	2	Solidago canadensis	1
Taraxacum officinale	2		

Transect Notes:

**Transect Number:** 3

**Compass Direction from Start:** 95

**Interval Data:**

**Ending Station** 136 **Community Type:** Bromus spp. / Phleum pratense

Species	Cover class	Species	Cover class
Agrostis gigantea	0	Alyssum alyssoides	1
Bare Ground	3	Bromus inermis	4
Carum carvi	3	Cirsium arvense	2
Cirsium vulgare	0	Cynoglossum officinale	0
Elymus cinereus	1	Elymus repens	2
Equisetum arvense	2	Juncus balticus	0
Lotus corniculatus	1	Melilotus officinalis	1
Phleum pratense	1	Poa pratensis	2
Populus tremuloides	0	Rumex crispus	0
Taraxacum officinale	2	Trifolium pratense	1

**Ending Station** 163 **Community Type:** Juncus spp. /

Species	Cover class	Species	Cover class
Agrostis gigantea	1	Carum carvi	1
Deschampsia caespitosa	1	Equisetum arvense	1
Juncus balticus	3	Lotus corniculatus	3
Phleum pratense	1	Poa pratensis	2
Populus tremuloides	1	Taraxacum officinale	1
Thlaspi arvense	0	Trifolium pratense	2

**Ending Station** 356 **Community Type:** Phleum pratense / Poa pratensis

Species	Cover class	Species	Cover class
Brassica napus	0	Bromus carinatus	0
Bromus inermis	2	Carum carvi	2
Cirsium arvense	0	Cynoglossum officinale	0
Deschampsia caespitosa	0	Elymus cinereus	0
Equisetum arvense	1	Lotus corniculatus	1
Poa pratensis	3	Populus tremuloides	0
Rumex crispus	0	Schedonorus pratensis	1
Solidago canadensis	1	Taraxacum officinale	3
Thlaspi arvense	1	Trifolium pratense	2
Trifolium repens	1		

**Ending Station** 646 **Community Type:** Juncus spp. /

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Carex praegracilis	0
Carex sp.	1	Carum carvi	1
Cirsium arvense	1	Deschampsia caespitosa	0
Equisetum arvense	2	Juncus balticus	5
Juncus bufonius	2	Juncus effusus	2
Juncus ensifolius	1	Lycopus asper	1
Mentha arvensis	1	Poa pratensis	0
Rumex crispus	0	Salix lutea	1
Scirpus microcarpus	0	Sisyrinchium idahoense	0
Thlaspi arvense	0	Trifolium pratense	1
Typha latifolia	0		

**Ending Station** 732 **Community Type:** Phleum pratense / Poa pratensis

<b>Species</b>	<b>Cover class</b>	<b>Species</b>	<b>Cover class</b>
Agrostis gigantea	1	Brassica napus	0
Bromus inermis	3	Carex limosa	0
Carum carvi	2	Cynoglossum officinale	0
Equisetum arvense	3	Juncus balticus	0
Juncus bufonius	0	Juncus effusus	0
Lotus corniculatus	1	Phleum pratense	3
Poa pratensis	2	Taraxacum officinale	3
Trifolium pratense	1		

Transect Notes:

## PLANTED WOODY VEGETATION SURVIVAL

Easton Ranch

Planting Type	#Planted	#Alive	Notes
Red-osier dogwood	250	12	
Sandbar willow	250	35	
Thinleaf alder	500	43	
Willow cuttings	200	75	

### Comments

A thorough sampling of the project area was employed to evaluate planted woody vegetation survival.

Easton Ranch

WILDLIFE

Birds

Were man-made nesting structures installed? Yes

If yes, type of structure: Bird Boxes

How many? 9

Are the nesting structures being used? Yes

Do the nesting structures need repairs? No

Nesting Structure Comments:

.

Species	#Observed	Behavior	Habitat
American White Pelican	1		
Bald Eagle	1		
Bullock's Oriole	1		FO,
Gray Partridge	3		
Great Blue Heron	1		
Mourning Dove	2		SS,
Northern Flicker	1		
Red-tailed Hawk	2		UP,
Red-winged Blackbird	3		
Sandhill Crane	2		
Tree Swallow	2		UP,
Vesper Sparrow	3		UP,
Western Meadowlark	1		UP,
Wilson's Snipe	2		MA,

Bird Comments

Pair of tree swallows using nest box.

BEHAVIOR CODES

**BP** = One of a breeding pair **BD** = Breeding display **F** = Foraging **FO** = Flyover **L** = Loafing **N** = Nesting

HABITAT CODES

**AB** = Aquatic bed **SS** = Scrub/Shrub **FO** = Forested **UP** = Upland buffer **I** = Island

**WM** = Wet meadow **MA** = Marsh **US** = Unconsolidated shore **MF** = Mud Flat **OW** = Open Water

## Mammals and Herptiles

Species	# Observed	Tracks	Scat	Burrows	Comments
Mountain Cottontail	1	No	No	No	
Northern Pocket Gopher		No	No	Yes	
White-tailed Deer	3	Yes	No	No	Three deer observed plus multiple tracks observed

**Wildlife Comments:**

## Easton Ranch

### PHOTOGRAPHS

Take photographs of the following permanent reference points listed in the check list below. Record the direction of the photograph using a compass. When at the site for the first time, establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3 feet above ground. Survey the location with a resource grade GPS and mark the location on the aerial photograph.

#### Photograph Checklist:

- ☒ One photograph for each of the four cardinal directions surrounding the wetland.
- ☒ At least one photograph showing upland use surrounding the wetland. If more than one upland exists then take additional photographs.
- ☒ At least one photograph showing the buffer surrounding the wetland.
- ☒ One photograph from each end of the vegetation transect, showing the transect.

Photo #	Latitude	Longitude	Bearing	Description
020011-102001	46.055286	-110.639137	340	PP7
020014,102001	46.056175	-110.64048	0	PP6
020016,102001	46.059883	-110.640404	90	PP5
020018,102001	46.060413	-110.640396	20	PP4B
020020,102002	46.060993	-110.640121	170	PP4A
020022,102002	46.061188	-110.639847	100	PP3
020024,102002	46.061028	-110.637207	200	PP2
020030,102003	46.059727	-110.637505	250	PP1
8142,8143	46.057975	-110.63992		SP01 upland
8145,8146	46.058099	-110.63998		SP02 wetland
8340	46.057281	-110.638306	5	T1 start
8341				swallow at bird box
8344	46.060627	-110.637779	185	T1 end
8345	46.06139	-110.639229	185	T2 start
8346	46.057594	-110.640343	0	T2 end
8347	46.056984	-110.640656	95	T3 start
8348	46.056114	-110.637924	265	T3 end
8349	46.0575091	-110.639425		SP03 wetland
8350	46.0575091	-110.639425		NE at SP03

#### Comments:



## ADDITIONAL ITEMS CHECKLIST

### Hydrology

- ☒ Map emergent vegetation/open water boundary on aerial photos.
- ☒ Observe extent of surface water. Look for evidence of past surface water elevations (e.g. drift lines, vegetation staining, erosion, etc).

### Photos

- ☒ One photo from the wetland toward each of the four cardinal directions
- ☒ One photo showing upland use surrounding the wetland.
- ☒ One photo showing the buffer around the wetland
- ☒ One photo from each end of each vegetation transect, toward the transect

### Vegetation

- ☒ Map vegetation community boundaries
- ☒ Complete Vegetation Transects

### Soils

- ☒ Assess soils

### Wetland Delineations

- ☒ Delineate wetlands according to applicable USACE protocol (1987 form or Supplement)
- ☒ Delineate wetland – upland boundary onto aerial photograph.

Wetland Delineation Comments

### Functional Assessments

- ☒ Complete and attach full MDT Montana Wetland Assessment Method field forms.

Functional Assessment Comments:

### Maintenance

Were man-made nesting structure installed at this site? Yes

If yes, do they need to be repaired? No

If yes, describe the problems below and indicate if any actions were taken to remedy the problems

Were man-made structures built or installed to impound water or control water flow  
into or out of the wetland? No

If yes, are the structures in need of repair?

If yes, describe the problems below.

--

# WETLAND DETERMINATION DATA FORM – Western Mountains, Valleys, and Coast Region

Project/Site: Easton City/County: Park Sampling Date: 6/26/2015  
 Applicant/Owner: MDT State: Montana Sampling Point: SP01  
 Investigator(s): R. McElowney, R. Quire Section, Township, Range: S 32 T 4N R 9E  
 Landform (hillslope, terrace, etc.): Floodplain Local relief (concave, convex, none): concave Slope (%): 0  
 Subregion (LRR): LRR E Lat: 46.057975 Long: -110.63992 Datum: SP NAD83  
 Soil Map Unit Name: Meadowcreek, rarely flooded-Nesda complex, 0 to 2% slopes NWI classification: Not Mapped

Are climatic / hydrologic conditions on the site typical for this time of year? Yes ☒ No ☐ (If no, explain in Remarks.)  
 Are Vegetation ☐, Soil ☐, or Hydrology ☐ significantly disturbed? Are "Normal Circumstances" present? Yes ☒ No ☐  
 Are Vegetation ☐, Soil ☐, or Hydrology ☐ naturally problematic? (If needed, explain any answers in Remarks.)

## SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes <input type="checkbox"/>	No <input checked="" type="checkbox"/>	Is the Sampled Area within a Wetland?	Yes <input type="checkbox"/>	No <input checked="" type="checkbox"/>
Hydric Soil Present?	Yes <input type="checkbox"/>	No <input checked="" type="checkbox"/>			
Wetland Hydrology Present?	Yes <input type="checkbox"/>	No <input checked="" type="checkbox"/>			
Remarks: Upland sample point.					

## VEGETATION - Use scientific names of plant

Tree Stratum	Plot size (30 Foot Radius)	Absolute % Cover:	Dominant Species?	Indicator Status	<b>Dominance Test worksheet</b> Number of Dominant Species that are OBL, FACW or FAC: <input type="text" value="1"/> (A) Total Number of Dominant Species Across All Strata: <input type="text" value="4"/> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <input type="text" value="25"/> % (A/B)
<u>Populus angustifolia</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>FACW</u>		

Herbaceous Stratum	Plot size ( 5 Foot Radius)	Absolute % Cover:	Dominant Species?	Indicator Status	<b>Prevalence Index worksheet</b> <table border="1"> <thead> <tr> <th>Total % Cover of:</th> <th>Multiply by:</th> </tr> </thead> <tbody> <tr> <td>OBL species 0 X 1</td> <td>0</td> </tr> <tr> <td>FACW species 16 X 2</td> <td>32</td> </tr> <tr> <td>FAC species 13 X 3</td> <td>39</td> </tr> <tr> <td>FACU species 78 X 4</td> <td>312</td> </tr> <tr> <td>UPL species 2 X 5</td> <td>10</td> </tr> <tr> <td>Column Totals</td> <td>109 (A) 393 (B)</td> </tr> </tbody> </table> <b>Prevalence Index = B/A = 3.60550</b>	Total % Cover of:	Multiply by:	OBL species 0 X 1	0	FACW species 16 X 2	32	FAC species 13 X 3	39	FACU species 78 X 4	312	UPL species 2 X 5	10	Column Totals	109 (A) 393 (B)
Total % Cover of:	Multiply by:																		
OBL species 0 X 1	0																		
FACW species 16 X 2	32																		
FAC species 13 X 3	39																		
FACU species 78 X 4	312																		
UPL species 2 X 5	10																		
Column Totals	109 (A) 393 (B)																		
<u>Agrostis stolonifera</u>	<u>10</u>	<input type="checkbox"/>	<u>FAC</u>																
<u>Carex sp.</u>	<u>1</u>	<input type="checkbox"/>	<u>NL</u>																
<u>Carum carvi</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>FACU</u>																
<u>Cirsium arvense</u>	<u>2</u>	<input type="checkbox"/>	<u>FAC</u>																
<u>Juncus bufonius</u>	<u>1</u>	<input type="checkbox"/>	<u>FACW</u>																
<u>Medicago lupulina</u>	<u>30</u>	<input checked="" type="checkbox"/>	<u>FACU</u>																
<u>Medicago sativa</u>	<u>1</u>	<input type="checkbox"/>	<u>UPL</u>																
<u>Pascopyrum smithii</u>	<u>10</u>	<input type="checkbox"/>	<u>FACU</u>																
<u>Phleum pratense</u>	<u>1</u>	<input type="checkbox"/>	<u>FAC</u>																
<u>Poa secunda</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>FACU</u>																
<u>Taraxacum officinale</u>	<u>8</u>	<input type="checkbox"/>	<u>FACU</u>																

Woody Vine Stratum	Plot size ( 30 Foot Radius)	Absolute % Cover:	Dominant Species?	Indicator Status	<b>Hydrophytic Vegetation Indicators</b> <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is <= 3.0 <input type="checkbox"/> 4 - Morphological Adaptations (Provide supporting data in remarks or on separate sheet. <input type="checkbox"/> 5 - Wetland Non-Vascular Plants <input type="checkbox"/> Problematic Hydrophytic Vegetation (Explain) Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic for #3, 4, 5.
<u>Percent Bare Ground</u>	<u>0</u>				

<b>Hydrophytic Vegetation Present?</b> Yes <input type="checkbox"/> NO <input checked="" type="checkbox"/>
---

Remarks:

US Army Corps of Engineers Western Mountains, Valleys, and Coasts - Version 2.0

## SOIL

Sampling Point: SP01

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		%	Redox Features				Type <sup>1</sup>	Loc <sup>2</sup>	Texture	Remarks
	Color (moist)			Color (moist)		%					
0-7.5	10YR	3/2	100	7.5YR	4/6	1		C	PL	Clay Loam	
7.5-15	10YR	3/1	100							Coarse Loamy Sand	Gravel throughout profile.

<sup>1</sup>Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains.<sup>2</sup>Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- |  |   |
|--|---|
| <input type="checkbox"/> Histosol (A1)                     | <input type="checkbox"/> Sandy Redox (S5)                         |
| <input type="checkbox"/> Histic Epipedon (A2)              | <input type="checkbox"/> Stripped Matrix (S6)                     |
| <input type="checkbox"/> Black Histic (A3)                 | <input type="checkbox"/> Loamy Mucky Mineral (F1) (except MLRA 1) |
| <input type="checkbox"/> Hydrogen Sulfide (A4)             | <input type="checkbox"/> Loamy Gleyed Matrix (F2)                 |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Depleted Matrix (F3)                     |
| <input type="checkbox"/> Thick Dark Surface (A12)          | <input type="checkbox"/> Redox Dark Surface (F6)                  |
| <input type="checkbox"/> Sandy Mucky Mineral (S1)          | <input type="checkbox"/> Depleted Dark Surface (F7)               |
| <input type="checkbox"/> Sandy Gleyed Matrix (S4)          | <input type="checkbox"/> Redox Depressions (F8)                   |

Indicators for Problematic Hydric Soils<sup>3</sup>:

- ☐ 2 cm Muck (A10)  
☐ Red Parent Material (TF2)  
☐ Very Shallow Dark Surface (TF12)  
☐ Other (Explain in Remarks)

<sup>3</sup>Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: \_\_\_\_\_

Depth (inches): \_\_\_\_\_

Hydric Soil Present? Yes ☐ No ☒

Remarks:

No hydric soil indicators observed during site visit.

## HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- |  |   |
|--|---|
| <input type="checkbox"/> Surface Water (A1)                        | <input type="checkbox"/> Water-Stained Leaves (B9) (except MLRA 1, 2, 4A, and 4B) |
| <input type="checkbox"/> High Water Table (A2)                     | <input type="checkbox"/> Salt Crust (B11)   |
| <input type="checkbox"/> Saturation (A3)                           | <input type="checkbox"/> Aquatic Invertebrates (B13)                              |
| <input type="checkbox"/> Water Marks (B1)                          | <input type="checkbox"/> Hydrogen Sulfide Odor (C1)                               |
| <input type="checkbox"/> Sediment Deposits (B2)                    | <input type="checkbox"/> Oxidized Rhizospheres along Living Roots (C3)            |
| <input type="checkbox"/> Drift Deposits (B3)                       | <input type="checkbox"/> Presence of Reduced Iron (C4)                            |
| <input type="checkbox"/> Algal Mat or Crust (B4)                   | <input type="checkbox"/> Recent Iron Reduction in Tilled Soils (C6)               |
| <input type="checkbox"/> Iron Deposits (B5)                        | <input type="checkbox"/> Stunted or Stressed Plants (D1) (LRR A)                  |
| <input type="checkbox"/> Surface Soil Cracks (B6)                  | <input type="checkbox"/> Other (Explain in Remarks)                               |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) |   |
| <input type="checkbox"/> Sparsely Vegetated Concave Surface (B8)   |   |

Secondary Indicators (2 or more required)

- ☐ Water-Stained Leaves (B9) (MLRA 1, 2, 4A, and 4B)  
☐ Drainage Patterns (B10)  
☐ Dry-Season Water Table (C2)  
☐ Saturation Visible on Aerial Imagery (C9)  
☐ Geomorphic Position (D2)  
☐ Shallow Aquitard (D3)  
☐ FAC-Neutral Test (D5)  
☐ Raised Ant Mounds (D6) (LRR A)  
☐ Frost-Heave Hummocks (D7)

Field Observations:

Surface Water Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_Water Table Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_Saturation Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_  
(includes capillary fringe)Wetland Hydrology Present? Yes ☐ No ☒

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

No primary or secondary hydrology indicators observed during site visit.

# WETLAND DETERMINATION DATA FORM – Western Mountains, Valleys, and Coast Region

Project/Site: Easton City/County: Park Sampling Date: 6/26/2015  
 Applicant/Owner: MDT State: Montana Sampling Point: SP02  
 Investigator(s): R. McElowney, R. Quire Section, Township, Range: S 32 T 4N R 9E  
 Landform (hillslope, terrace, etc.): Floodplain Local relief (concave, convex, none): concave Slope (%): 0  
 Subregion (LRR): LRR E Lat: 46.058099 Long: -110.63998 Datum: SP NAD83  
 Soil Map Unit Name: Meadowcreek, rarely flooded-Nesda complex, 0 to 2% slopes NWI classification: Not Mapped

Are climatic / hydrologic conditions on the site typical for this time of year? Yes ☒ No ☐ (If no, explain in Remarks.)  
 Are Vegetation ☐, Soil ☐, or Hydrology ☐ significantly disturbed? Are "Normal Circumstances" present? Yes ☒ No ☐  
 Are Vegetation ☐, Soil ☐, or Hydrology ☐ naturally problematic? (If needed, explain any answers in Remarks.)

## SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>	Is the Sampled Area within a Wetland?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>
Hydric Soil Present?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>			
Wetland Hydrology Present?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>			
Remarks: PEM Wetland sample point.					

## VEGETATION - Use scientific names of plant

Tree Stratum	Plot size (30 Foot Radius)	Absolute % Cover:	Dominant Species?	Indicator Status	<b>Dominance Test worksheet</b> Number of Dominant Species that are OBL, FACW or FAC: <input type="text" value="3"/> (A) Total Number of Dominant Species Across All Strata: <input type="text" value="3"/> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <input type="text" value="100"/> % (A/B)																																
<u>Populus angustifolia</u>		1	<input checked="" type="checkbox"/>	FACW																																	
<u>Salix lutea</u>		4	<input checked="" type="checkbox"/>	OBL																																	
<b>Prevalence Index worksheet</b> <table border="1"> <thead> <tr> <th>Total % Cover of:</th> <th>Multiply by:</th> </tr> </thead> <tbody> <tr> <td>OBL species 4 X 1</td> <td>4</td> </tr> <tr> <td>FACW species 47 X 2</td> <td>94</td> </tr> <tr> <td>FAC species 1 X 3</td> <td>3</td> </tr> <tr> <td>FACU species 9 X 4</td> <td>36</td> </tr> <tr> <td>UPL species 1 X 5</td> <td>5</td> </tr> <tr> <td>Column Totals 62 (A)</td> <td>142 (B)</td> </tr> </tbody> </table>						Total % Cover of:	Multiply by:	OBL species 4 X 1	4	FACW species 47 X 2	94	FAC species 1 X 3	3	FACU species 9 X 4	36	UPL species 1 X 5	5	Column Totals 62 (A)	142 (B)																		
Total % Cover of:	Multiply by:																																				
OBL species 4 X 1	4																																				
FACW species 47 X 2	94																																				
FAC species 1 X 3	3																																				
FACU species 9 X 4	36																																				
UPL species 1 X 5	5																																				
Column Totals 62 (A)	142 (B)																																				
<b>Prevalence Index = B/A = 2.29032</b>																																					
<b>Hydrophytic Vegetation Indicators</b> <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input checked="" type="checkbox"/> 2 - Dominance Test is >50% <input checked="" type="checkbox"/> 3 - Prevalence Index is <= 3.0 <input type="checkbox"/> 4 - Morphological Adaptations (Provide supporting data in remarks or on separate sheet.) <input type="checkbox"/> 5 - Wetland Non-Vascular Plants <input type="checkbox"/> Problematic Hydrophytic Vegetation (Explain)																																					
Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic for #3, 4, 5.																																					
<b>Hydrophytic Vegetation Present?</b> Yes <input checked="" type="checkbox"/> NO <input type="checkbox"/>																																					
<b>Herbaceous Stratum</b> Plot size ( 5 Foot Radius) <table border="1"> <tbody> <tr> <td><u>Carum carvi</u></td> <td>1</td> <td><input type="checkbox"/></td> <td>FACU</td> </tr> <tr> <td><u>Equisetum arvense</u></td> <td>1</td> <td><input type="checkbox"/></td> <td>FAC</td> </tr> <tr> <td><u>Juncus balticus</u></td> <td>40</td> <td><input checked="" type="checkbox"/></td> <td>FACW</td> </tr> <tr> <td><u>Mentha arvensis</u></td> <td>5</td> <td><input type="checkbox"/></td> <td>FACW</td> </tr> <tr> <td><u>Pascopyrum smithii</u></td> <td>1</td> <td><input type="checkbox"/></td> <td>FACU</td> </tr> <tr> <td><u>Sisyrinchium idahoense</u></td> <td>1</td> <td><input type="checkbox"/></td> <td>FACW</td> </tr> <tr> <td><u>Taraxacum officinale</u></td> <td>7</td> <td><input type="checkbox"/></td> <td>FACU</td> </tr> <tr> <td><u>Trifolium sp.</u></td> <td>1</td> <td><input type="checkbox"/></td> <td>NL</td> </tr> </tbody> </table>						<u>Carum carvi</u>	1	<input type="checkbox"/>	FACU	<u>Equisetum arvense</u>	1	<input type="checkbox"/>	FAC	<u>Juncus balticus</u>	40	<input checked="" type="checkbox"/>	FACW	<u>Mentha arvensis</u>	5	<input type="checkbox"/>	FACW	<u>Pascopyrum smithii</u>	1	<input type="checkbox"/>	FACU	<u>Sisyrinchium idahoense</u>	1	<input type="checkbox"/>	FACW	<u>Taraxacum officinale</u>	7	<input type="checkbox"/>	FACU	<u>Trifolium sp.</u>	1	<input type="checkbox"/>	NL
<u>Carum carvi</u>	1	<input type="checkbox"/>	FACU																																		
<u>Equisetum arvense</u>	1	<input type="checkbox"/>	FAC																																		
<u>Juncus balticus</u>	40	<input checked="" type="checkbox"/>	FACW																																		
<u>Mentha arvensis</u>	5	<input type="checkbox"/>	FACW																																		
<u>Pascopyrum smithii</u>	1	<input type="checkbox"/>	FACU																																		
<u>Sisyrinchium idahoense</u>	1	<input type="checkbox"/>	FACW																																		
<u>Taraxacum officinale</u>	7	<input type="checkbox"/>	FACU																																		
<u>Trifolium sp.</u>	1	<input type="checkbox"/>	NL																																		
<b>Woody Vine Stratum</b> Plot size ( 30 Foot Radius) <table border="1"> <tbody> <tr> <td><u>Percent Bare Ground</u></td> <td>40</td> </tr> </tbody> </table>						<u>Percent Bare Ground</u>	40																														
<u>Percent Bare Ground</u>	40																																				

Remarks:

## SOIL

Sampling Point: SP02

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type <sup>1</sup>	Loc <sup>2</sup>		
0-8.5	10YR	3/2	100				Clay	
8.5-16	10YR	4/2	80	2.5YR	4/8	20	C PL	Sandy Clay Loam

<sup>1</sup>Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains.<sup>2</sup>Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- |   |   |
|---|---|
| <input type="checkbox"/> Histosol (A1)                                | <input type="checkbox"/> Sandy Redox (S5)                         |
| <input type="checkbox"/> Histic Epipedon (A2)                         | <input type="checkbox"/> Stripped Matrix (S6)                     |
| <input type="checkbox"/> Black Histic (A3)                            | <input type="checkbox"/> Loamy Mucky Mineral (F1) (except MLRA 1) |
| <input type="checkbox"/> Hydrogen Sulfide (A4)                        | <input type="checkbox"/> Loamy Gleyed Matrix (F2)                 |
| <input checked="" type="checkbox"/> Depleted Below Dark Surface (A11) | <input checked="" type="checkbox"/> Depleted Matrix (F3)          |
| <input type="checkbox"/> Thick Dark Surface (A12)                     | <input type="checkbox"/> Redox Dark Surface (F6)                  |
| <input type="checkbox"/> Sandy Mucky Mineral (S1)                     | <input type="checkbox"/> Depleted Dark Surface (F7)               |
| <input type="checkbox"/> Sandy Gleyed Matrix (S4)                     | <input type="checkbox"/> Redox Depressions (F8)                   |

Indicators for Problematic Hydric Soils<sup>3</sup>:

- ☐ 2 cm Muck (A10)  
☐ Red Parent Material (TF2)  
☐ Very Shallow Dark Surface (TF12)  
☐ Other (Explain in Remarks)

<sup>3</sup>Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: \_\_\_\_\_

Depth (inches): \_\_\_\_\_

Hydric Soil Present? Yes ☒ No ☐

Remarks:

Dark colored peds scattered throughout (likely decayed plants or historic sediment deposition).

## HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- |  |   |
|--|---|
| <input type="checkbox"/> Surface Water (A1)                        | <input type="checkbox"/> Water-Stained Leaves (B9) (except MLRA 1, 2, 4A, and 4B) |
| <input type="checkbox"/> High Water Table (A2)                     | <input type="checkbox"/> Salt Crust (B11)   |
| <input type="checkbox"/> Saturation (A3)                           | <input type="checkbox"/> Aquatic Invertebrates (B13)                              |
| <input type="checkbox"/> Water Marks (B1)                          | <input type="checkbox"/> Hydrogen Sulfide Odor (C1)                               |
| <input type="checkbox"/> Sediment Deposits (B2)                    | <input checked="" type="checkbox"/> Oxidized Rhizospheres along Living Roots (C3) |
| <input type="checkbox"/> Drift Deposits (B3)                       | <input type="checkbox"/> Presence of Reduced Iron (C4)                            |
| <input type="checkbox"/> Algal Mat or Crust (B4)                   | <input type="checkbox"/> Recent Iron Reduction in Tilled Soils (C6)               |
| <input type="checkbox"/> Iron Deposits (B5)                        | <input type="checkbox"/> Stunted or Stressed Plants (D1) (LRR A)                  |
| <input type="checkbox"/> Surface Soil Cracks (B6)                  | <input type="checkbox"/> Other (Explain in Remarks)                               |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) |   |
| <input type="checkbox"/> Sparsely Vegetated Concave Surface (B8)   |   |

Secondary Indicators (2 or more required)

- ☐ Water-Stained Leaves (B9) (MLRA 1, 2, 4A, and 4B)  
☐ Drainage Patterns (B10)  
☐ Dry-Season Water Table (C2)  
☐ Saturation Visible on Aerial Imagery (C9)  
☒ Geomorphic Position (D2)  
☐ Shallow Aquitard (D3)  
☒ FAC-Neutral Test (D5)  
☐ Raised Ant Mounds (D6) (LRR A)  
☐ Frost-Heave Hummocks (D7)

Field Observations:

Surface Water Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_Water Table Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_Saturation Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_  
(includes capillary fringe)Wetland Hydrology Present? Yes ☒ No ☐

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

Project/Site: Easton City/County: Park Sampling Date: 6/26/2015  
Applicant/Owner: MDT State: Montana Sampling Point: SP03  
Investigator(s): R. McElowney, J. Johnson, F. McNew Section, Township, Range: S 32 T 4N R 9E  
Landform (hillslope, terrace, etc.): Terrace Local relief (concave, convex, none): flat Slope (%): 0  
Subregion (LRR): LRR E Lat: 46.0575091 Long: -110.639425 Datum: WGS84  
Soil Map Unit Name: Meadowcreek, rarely-flooded Nes NWI classification: Upland

**SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.**

Hydrophytic Vegetation Present?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>	<b>Is the Sampled Area within a Wetland?</b>	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>
Hydric Soil Present?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>			
Wetland Hydrology Present?	Yes <input checked="" type="checkbox"/>	No <input type="checkbox"/>			
<b>Remarks:</b> Wetland sample point. Area appears to have lost hydrology and may be transitioning to an upland community type.					

<u>Tree Stratum</u>	Plot size (30 Foot Radius)	Absolute % Cover:	Dominant Species?	Indicator Status																																												
<u>Sapling/Shrub Stratum</u>	Plot size (15 Foot Radius)																																															
<u>Herbaceous Stratum</u>	Plot size ( 5 Foot Radius)																																															
<table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td style="width: 40%;">Agrostis gigantea</td> <td style="width: 10%;">1</td> <td style="width: 5%;"><input type="checkbox"/></td> <td style="width: 45%;">FAC</td> </tr> <tr> <td>Carex nebrascensis</td> <td>1</td> <td><input type="checkbox"/></td> <td>OBL</td> </tr> <tr> <td>Carex sp.</td> <td>1</td> <td><input type="checkbox"/></td> <td>NL</td> </tr> <tr> <td>Carex utriculata</td> <td>8</td> <td><input type="checkbox"/></td> <td>OBL</td> </tr> <tr> <td>Deschampsia caespitosa</td> <td>2</td> <td><input type="checkbox"/></td> <td>FACW</td> </tr> <tr> <td>Elymus repens</td> <td>1</td> <td><input type="checkbox"/></td> <td>FAC</td> </tr> <tr> <td>Glyceria grandis</td> <td>1</td> <td><input type="checkbox"/></td> <td>OBL</td> </tr> <tr> <td>Juncus balticus</td> <td>10</td> <td><input type="checkbox"/></td> <td>FACW</td> </tr> <tr> <td>Poa pratensis</td> <td>15</td> <td><input type="checkbox"/></td> <td>FAC</td> </tr> <tr> <td>Scirpus microcarpus</td> <td>60</td> <td><input checked="" type="checkbox"/></td> <td>OBL</td> </tr> <tr> <td>Thlaspi arvense</td> <td>1</td> <td><input type="checkbox"/></td> <td>UPL</td> </tr> </tbody> </table>					Agrostis gigantea	1	<input type="checkbox"/>	FAC	Carex nebrascensis	1	<input type="checkbox"/>	OBL	Carex sp.	1	<input type="checkbox"/>	NL	Carex utriculata	8	<input type="checkbox"/>	OBL	Deschampsia caespitosa	2	<input type="checkbox"/>	FACW	Elymus repens	1	<input type="checkbox"/>	FAC	Glyceria grandis	1	<input type="checkbox"/>	OBL	Juncus balticus	10	<input type="checkbox"/>	FACW	Poa pratensis	15	<input type="checkbox"/>	FAC	Scirpus microcarpus	60	<input checked="" type="checkbox"/>	OBL	Thlaspi arvense	1	<input type="checkbox"/>	UPL
Agrostis gigantea	1	<input type="checkbox"/>	FAC																																													
Carex nebrascensis	1	<input type="checkbox"/>	OBL																																													
Carex sp.	1	<input type="checkbox"/>	NL																																													
Carex utriculata	8	<input type="checkbox"/>	OBL																																													
Deschampsia caespitosa	2	<input type="checkbox"/>	FACW																																													
Elymus repens	1	<input type="checkbox"/>	FAC																																													
Glyceria grandis	1	<input type="checkbox"/>	OBL																																													
Juncus balticus	10	<input type="checkbox"/>	FACW																																													
Poa pratensis	15	<input type="checkbox"/>	FAC																																													
Scirpus microcarpus	60	<input checked="" type="checkbox"/>	OBL																																													
Thlaspi arvense	1	<input type="checkbox"/>	UPL																																													
<u>Woody Vine Stratum</u>	Plot size ( 30 Foot Radius)																																															
<table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td style="width: 40%;">Percent Bare Ground</td> <td style="width: 10%;">1</td> <td style="width: 5%;"></td> <td style="width: 45%;"></td> </tr> </tbody> </table>					Percent Bare Ground	1																																										
Percent Bare Ground	1																																															

**Dominance Test worksheet**

Number of Dominant Species that are OBL, FACW or FAC: 1 (A)

Total Number of Dominant Species Across All Strata: 1 (B)

Percent of Dominant Species That Are OBL, FACW, or FAC: 100 % (A/B)

---

**Prevalence Index worksheet**

Total % Cover of:		Multiply by:	
OBL species	70 X 1		<span style="border: 1px solid black; padding: 2px 10px;">70</span>
FACW species	12 X 2		<span style="border: 1px solid black; padding: 2px 10px;">24</span>
FAC species	17 X 3		<span style="border: 1px solid black; padding: 2px 10px;">51</span>
FACU species	0 X 4		<span style="border: 1px solid black; padding: 2px 10px;">0</span>
UPL species	2 X 5		<span style="border: 1px solid black; padding: 2px 10px;">10</span>
Column Totals	<span style="border: 1px solid black; padding: 2px 10px;">101</span> (A)		<span style="border: 1px solid black; padding: 2px 10px;">155</span> (B)

**Prevalence Index = B/A = 1.53465**

---

**Hydrophytic Vegetation Indicators**

- ☐ 1 - Rapid Test for Hydrophytic Vegetation
- ☒ 2 - Dominance Test is >50%
- ☒ 3 - Prevalence Index is <= 3.0
- ☐ 4 - Morphological Adaptations (Provide supporting data in remarks or on separate sheet.
- ☐ 5 - Wetland Non-Vascular Plants
- ☐ Problematic Hydrophytic Vegetation (Explain)

Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic for #3, 4, 5.

---

**Hydrophytic Vegetation Present?**      Yes ☒      NO ☐

Remarks:



## SOIL

Sampling Point: SP03

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type <sup>1</sup>	Loc <sup>2</sup>		
>15							Cobbles	Soil is dry.
15	10YR	2/2					Silt Loam	No redox in upper 15 inches.

<sup>1</sup>Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains.<sup>2</sup>Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

Indicators for Problematic Hydric Soils<sup>3</sup>:

- |  |   |
|--|---|
| <input type="checkbox"/> Histosol (A1)                     | <input type="checkbox"/> Sandy Redox (S5)                         |
| <input type="checkbox"/> Histic Epipedon (A2)              | <input type="checkbox"/> Stripped Matrix (S6)                     |
| <input type="checkbox"/> Black Histic (A3)                 | <input type="checkbox"/> Loamy Mucky Mineral (F1) (except MLRA 1) |
| <input type="checkbox"/> Hydrogen Sulfide (A4)             | <input type="checkbox"/> Loamy Gleyed Matrix (F2)                 |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Depleted Matrix (F3)                     |
| <input type="checkbox"/> Thick Dark Surface (A12)          | <input type="checkbox"/> Redox Dark Surface (F6)                  |
| <input type="checkbox"/> Sandy Mucky Mineral (S1)          | <input type="checkbox"/> Depleted Dark Surface (F7)               |
| <input type="checkbox"/> Sandy Gleyed Matrix (S4)          | <input type="checkbox"/> Redox Depressions (F8)                   |

- ☐ 2 cm Muck (A10)
- ☐ Red Parent Material (TF2)
- ☐ Very Shallow Dark Surface (TF12)
- ☒ Other (Explain in Remarks)

<sup>3</sup>Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: \_\_\_\_\_

Depth (inches): \_\_\_\_\_

Hydric Soil Present? Yes ☒ No ☐

Remarks:

Site had geomorphic position in relic channel and passed the FAC-neutral test. It is unclear why site does not have hydrology.

## HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- |  |   |
|--|---|
| <input type="checkbox"/> Surface Water (A1)                        | <input type="checkbox"/> Water-Stained Leaves (B9) (except MLRA 1, 2, 4A, and 4B) |
| <input type="checkbox"/> High Water Table (A2)                     | <input type="checkbox"/> Salt Crust (B11)   |
| <input checked="" type="checkbox"/> Saturation (A3)                | <input type="checkbox"/> Aquatic Invertebrates (B13)                              |
| <input type="checkbox"/> Water Marks (B1)                          | <input type="checkbox"/> Hydrogen Sulfide Odor (C1)                               |
| <input type="checkbox"/> Sediment Deposits (B2)                    | <input type="checkbox"/> Oxidized Rhizospheres along Living Roots (C3)            |
| <input type="checkbox"/> Drift Deposits (B3)                       | <input type="checkbox"/> Presence of Reduced Iron (C4)                            |
| <input type="checkbox"/> Algal Mat or Crust (B4)                   | <input type="checkbox"/> Recent Iron Reduction in Tilled Soils (C6)               |
| <input type="checkbox"/> Iron Deposits (B5)                        | <input type="checkbox"/> Stunted or Stressed Plants (D1) (LRR A)                  |
| <input type="checkbox"/> Surface Soil Cracks (B6)                  | <input type="checkbox"/> Other (Explain in Remarks)                               |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) |   |
| <input type="checkbox"/> Sparsely Vegetated Concave Surface (B8)   |   |

Secondary Indicators (2 or more required)

- ☐ Water-Stained Leaves (B9) (MLRA 1, 2, 4A, and 4B)
- ☐ Drainage Patterns (B10)
- ☐ Dry-Season Water Table (C2)
- ☐ Saturation Visible on Aerial Imagery (C9)
- ☒ Geomorphic Position (D2)
- ☐ Shallow Aquitard (D3)
- ☒ FAC-Neutral Test (D5)
- ☐ Raised Ant Mounds (D6) (LRR A)
- ☐ Frost-Heave Hummocks (D7)

Field Observations:

Surface Water Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_Water Table Present? Yes ☐ No ☒ Depth (inches): \_\_\_\_\_Saturation Present? Yes ☒ No ☐ Depth (inches): \_\_\_\_\_  
(includes capillary fringe)Wetland Hydrology Present? Yes ☒ No ☐

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

Did not observe any primary indicators of hydrology during site visit.

# MDT Montana Wetland Assessment Form (revised March 2008)

1. Project name  2. MDT project#  Control#

3. Evaluation Date  4. Evaluators  5. Wetland/Site# (s)

6. Wetland Location(s): T  R  Sec1  T  R  Sec2

Approx Stationing or Mileposts

Watershed  Watershed/County

7. Evaluating Agency

8. Wetland size acres

Purpose of Evaluation

☐ Wetlands potentially affected by MDT project

☐ Mitigation Wetlands: pre-construction

☒ Mitigation Wetlands: post construction

☐ Other

9. Assessment area (AA) size (acres)

How assessed:

How assessed:

## 10. Classification of Wetland and Aquatic Habitats in AA

HGM Class (Brinson)	Class (Cowardin)	Modifier (Cowardin)	Water Regime	% of AA
Riverine	Emergent Wetland	Excavated	Seasonal/Intermittent	90
Depressional	Aquatic Bed	Excavated	Seasonal/Intermittent	10

11. Estimated Relative Abundance

## 12. General Condition of AA

i. Disturbance: (use matrix below to determine [circle] appropriate response – see instructions for Montana-listed noxious weed and aquatic nuisance vegetation species (ANVS) lists)

Conditions within AA	Predominant conditions adjacent to (within 500 feet of) AA		
	Managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings; and noxious weed or ANVS cover is <=15%.	Land not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to minor clearing; contains few roads or buildings; noxious weed or ANVS cover is <=30%.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >=30%.
AA occurs and is managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings; and noxious weed or ANVS cover is <=15%.	low disturbance	low disturbance	moderate disturbance
AA not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings; noxious weed or ANVS cover is <=30%.	moderate disturbance	moderate disturbance	high disturbance
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >=30%.	high disturbance	high disturbance	high disturbance

## Comments: (types of disturbance, intensity, season, etc)

Limited agriculture (hay) and a few ranch structures to the east. Undeveloped riparian corridor and herbaceous uplands to north, south, and west. Two species of noxious weeds were present within the AA. The AA is managed in a natural state, as are most of the lands within 500 feet of the AA.

## ii. Prominent noxious, aquatic nuisance, other exotic species:

Cirsium arvense and Cynoglossum officinale

## iii. Provide brief descriptive summary of AA and surrounding land use/habitat

The AA consists of four constructed wetland cells. The lowest contours of the wetland cells are seasonally inundated and have developed wetland characteristics. The higher elevations lack wetland characteristics and support upland plant communities. The cells are bordered by limited agriculture (hay) and an undeveloped riparian corridor.

**13. Structural Diversity:** (based on number of "Cowardin" **vegetated** classes present [do not include unvegetated classes], see #10 above)

Existing # of "Cowardin" Vegetated Classes in AA	Initial Rating	Is current management preventing (passive) existence of additional vegetated classes?		Modified Rating
>=3 (or 2 if 1 is forested) classes	H	NA	NA	NA
2 (or 1 if forested) classes	M	NA	NA	NA
1 class, but not a monoculture	M	<NO	YES>	L
1 class, monoculture (1 species comprises >=90% of total cover)	L	NA	NA	NA

**Comments:** The AA consists of palustrine emergent wetlands (PEM) and aquatic beds in the deeper depressions.

## SECTION PERTAINING to FUNCTIONS VALUES ASSESSMEN

### 14A. Habitat for Federally Listed or Proposed Threatened or Endangered Plants or Animals:

i. AA is Documented (D) or Suspected (S) to contain (check one based on definitions contained in instructions):

Primary or critical habitat (list species) ☐ D ☐ S

Secondary habitat (list Species) ☐ D ☐ S

Incidental habitat (list species) ☐ D ☐ S

No usable habitat ☒ S

ii. **Rating** (use the conclusions from i above and the matrix below to arrive at [check] the functional points and rating)

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	None
Functional Points and Rating	1H	.9H	.8H	.7M	.3L	.1L	0L

Sources for documented use

### 14B. Habitat for plant or animals rated S1, S2, or S3 by the Montana Natural Heritage Program: (not including species listed in 14A above)

i. AA is Documented (D) or Suspected (S) to contain (check one based on definitions contained in instructions):

Primary or critical habitat (list species) ☐ D ☐ S

Secondary habitat (list Species) ☐ D ☐ S

Incidental habitat (list species) ☒ D ☐ S Golden Eagle (S3)

No usable habitat ☐ S

ii. **Rating** (use the conclusions from i above and the matrix below to arrive at [check] the functional points and rating)

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	None
<b>S1 Species:</b> Functional Points and Rating	1H	.8H	.7M	.6M	.2L	.1L	0L
<b>S2 and S3 Species:</b> Functional Points and Rating	.9H	.7M	.6M	.5M	.2L	.1L	0L

Sources for documented use

MTNHP, 2013 field observations

**14C. General Wildlife Habitat Rating:**

i. Evidence of overall wildlife use in the AA (check substantial, moderate, or low based on supporting evidence):

Moderate

**Substantial** (based on any of the following [check]):

- ☐ observations of abundant wildlife #s or high species diversity (during any period)
- ☐ abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ presence of extremely limiting habitat features not available in the surrounding area
- ☐ interviews with local biologists with knowledge of the AA

**Minimal** (based on any of the following [check]):

- ☐ few or no wildlife observations during peak use periods
- ☐ little to no wildlife sign
- ☐ sparse adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

**Moderate** (based on any of the following [check]):

- ☐ observations of scattered wildlife groups or individuals or relatively few species during peak periods
- ☒ common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ adequate adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

ii. **Wildlife** habitat features (Working from top to bottom, check appropriate AA attributes in matrix to arrive at rating. Structural diversity is from #13. For class cover to be considered evenly distributed, the most and least prevalent **vegetated** classes must be within 20% of each other in terms of their percent composition of the AA (see #10). Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; and A = absent [see instructions for further definitions of these terms])

Structural diversity (see #13)	High								Moderate								Low			
	Even				Uneven				Even				Uneven				Even			
Class cover distribution (all vegetated classes)	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Duration of surface water in 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Low disturbance at AA (see #12)	E	E	E	H	E	E	H	H	E	H	H	M	E	H	M	M	E	H	M	M
Moderate disturbance at AA (see #12)	H	H	H	H	H	H	H	M	H	H	M	M	H	M	M	L	H	M	L	L
High disturbance at AA (see #12)	M	M	M	L	M	M	L	L	M	M	L	L	M	L	L	L	L	L	L	L

iii. **Rating** (use the conclusions from i and ii above and the matrix below to arrive at [check] the functional points and rating)

Evidence of wildlife use (i)	Wildlife habitat features rating (ii)											
	Exceptional			High			Moderate			Low		
Substantial		1E			.9H			.8H			.7M	
Moderate		.9H			.7M			.5M			.3L	
Minimal		.6M			.4M			.2L			.1L	

**Comments**

Many shorebirds and waterfowl have been documented using this site from 2010 to 2015. There is an active bald eagle (*Haliaeetus leucocephalus*) nest directly southwest of the site on the west side of the Shields River the project site is within the primary habitat zone for bald eagles (L. Urban. MDT. personal communications. 2015).

**14D. General Fish Habitat Rating:** (Assess this function if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [i.e., fish use is precluded by perched culvert or other barrier, etc.]. If the AA is not used by fish, fish use is not restorable due to habitat constraints, or is not desired from a management perspective [such as fish entrapped in a canal], then check

☒ **NA** here and proceed to 14E.)

i. **Habitat Quality and Known / Suspected Fish Species in AA** (use matrix to arrive at [check] the functional points and rating)

Duration of surface water in AA	Permanent / Perennial						Seasonal / Intermittent						Temporary / Ephemeral					
	Optimal		Adequate		Poor		Optimal		Adequate		Poor		Optimal		Adequate		Poor	
Aquatic hiding / resting / escape cover	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S
Thermal cover optimal / suboptimal	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S
FWP Tier I fish species	1E	.9H	.8H	.7M	.6M	.5M	.9H	.8H	.7M	.6M	.5M	.4M	.7M	.6M	.5M	.4M	.3L	.3L
FWP Tier II or Native Game fish species	.9H	.8H	.7M	.6M	.5M	.5M	.8H	.7M	.6M	.5M	.4M	.4M	.6M	.5M	.4M	.3L	.2L	.2L
FWP Tier III or Introduced Game fish	.8H	.7M	.6M	.5M	.5M	.4M	.7M	.6M	.5M	.4M	.4M	.3L	.5M	.4M	.3L	.2L	.2L	.1L
FWP Non-Game Tier IV or No fish species	.5M	.5M	.5M	.4M	.4M	.3L	.4M	.4M	.4M	.3L	.3L	.2L	.2L	.2L	.2L	.1L	.1L	.1L



Sources used for identifying fish sp. potentially found in AA:

ii. **Modified Rating** (NOTE: Modified score cannot exceed 1 or be less than 0.1)

a) Is fish use of the AA significantly reduced by a culvert, dike, or other man-made structure or activity or is the waterbody included on the current final MDEQ list of waterbodies in need of TMDL development with listed "Probable Impaired Uses" including cold or warm water fishery or aquatic life support, or do aquatic nuisance plant or animal species (see Appendix E) occur in fish habitat? Y ☐ N ☒ If yes, reduce score in i above by 0.1: **Modified Rating**

b) Does the AA contain a documented spawning area or other critical habitat feature (i.e., sanctuary pool, upwelling area, etc.- specify in comments) for native fish or introduced game fish? ☐ Y ☒ N If yes, add 0.1 to the adjusted score in i or iia above:

**Modified Rating**

iii. **Final Score and Rating:**

**Comments:** Wetland cells are isolated from Shields River with no fish habitat present.

**14E. Flood Attenuation:** (Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA are not flooded from in-channel or overbank flow, click ☐ NA here and proceed to 14F.)

i. **Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

Estimated or Calculated Entrenchment (Rosgen 1994, 1996)	Slightly entrenched - C, D, E stream types			Moderately entrenched - B stream type			Entrenched-A, F, G stream types		
% of flooded wetland classified as forested and/or scrub/shrub	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains <b>no outlet or restricted outlet</b>	1H	.9H	.6M	.8H	.7M	.5M	.4M	.3L	.2L
AA contains <b>unrestricted outlet</b>	.9H	.8H	.5M	.7M	.6M	.4M	.3L	.2L	.1L

Slightly Entrenched ER = >2.2			Moderately Entrenched ER = 1.41 - 2.2		Entrenched ER = 1.0 - 1.4	
C stream type	D stream type	E stream type	B stream type		A stream type	F stream type



**Floodprone width**  / **Bankfull width**  = **Entrenchment ratio**

ii. Are 10 acres of wetland in the AA subject to flooding AND are man-made features which may be significantly damaged by floods located within 0.5 mile downstream of the AA (check)? Y ☐ N ☒

**Comments:**

AA receives overbank flow from Shields River during high flow events.

**14F. Short and Long Term Surface Water Storage:** (Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow. If no wetlands in the AA are subject to flooding or ponding, click ☐ NA here and proceed to 14G.)

i. **Rating** (Working from top to bottom, use the matrix below to arrive at [check] the functional points and rating. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral [see instructions for further definitions of these terms].)

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding	>5 acre feet			1.1 to 5 acre feet			≤1 acre foot		
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond <b>≥ 5 out of 10 years</b>	1H	.9H	.8H	.8H	.6M	.5M	.4M	.3L	.2L
Wetlands in AA flood or pond <b>&lt; 5 out of 10 years</b>	.9H	.8H	.7M	.7M	.5M	.4M	.3L	.2L	.1L

**Comments:** (9.34 acre wetland) \* (1 ft. max depth at highwater) = 9.34 acre feet.

**14G. Sediment/Nutrient/Toxicant Retention and Removal:** (Applies to wetlands with potential to receive sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, click ☐ **NA** here and proceed to 14H.)

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating [H = high, M = moderate, or L = low])

Sediment, nutrient, and toxicant input levels within AA	AA receives or surrounding land use with potential to deliver levels of sediments, nutrients, or compounds at levels such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use with potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% cover of wetland vegetation in AA	≥ 70%		< 70%		≥ 70%		< 70%	
Evidence of flooding / ponding in AA	Yes	No	Yes	No	Yes	No	Yes	No
AA contains <b>no or restricted</b> outlet	1H	.8H	.7M	.5M	.5M	.4M	.3L	.2L
AA contains <b>unrestricted</b> outlet	.9H	.7M	.6M	.4M	.4M	.3L	.2L	.1L

**Comments:** There was evidence of ponding and flooding in 2011, 2012, 2014, and 2015. There was no evidence of ponding or flooding in 2013.

**14H Sediment/Shoreline Stabilization:** (Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action. If 14H does not apply, click ☐ **NA** here and proceed to 14I.)

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

% Cover of <b>wetland</b> streambank or shoreline by species with stability ratings of 6 (see <b>Appendix F</b> ).	Duration of surface water adjacent to rooted vegetation					
	Permanent / Perennial		Seasonal / Intermittent		Temporary / Ephemeral	
≥ 65%	1H		.9H		.7M	
35-64%	.7M		.6M		.5M	
< 35%	.3L		.2L		.1L	

Deep-rooted species observed in 2015 include bulrush, spikerush, sedges, and rushes.

**Comments:**

#### 14I. Production Export/Food Chain Support:

**i. Level of Biological Activity** (synthesis of wildlife and fish habitat ratings [check])

General Fish Habitat Rating (14D.iii.)	General Wildlife Habitat Rating (14C.iii.)					
	E/H		M		L	
E/H	H		H		M	
M	H		M		M	
L	M		M		L	
N/A	H		M		L	

**ii. Rating** (Working from top to bottom, use the matrix below to arrive at [check] the functional points and rating. Factor A = acreage of vegetated wetland component in the AA; Factor B = level of biological activity rating from above (14I.i.); Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to duration of surface water in the AA, where P/P, S/I, and T/E are as previously defined, and A = "absent" [see instructions for further definitions of these terms].)

A	Vegetated component >5 acres						Vegetated component 1-5 acres						Vegetated component <1 acre					
	High		Moderate		Low		High		Moderate		Low		High		Moderate		Low	
B	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
C	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
P/P	1E	.7H	.8H	.5M	.6M	.4M	.9H	.6M	.7H	.4M	.5M	.3L	.8H	.6M	.6M	.4M	.3L	.2L
S/I	.9H	.6M	.7H	.4M	.5M	.3L	.8H	.5M	.6M	.3L	.4M	.2L	.7H	.5M	.5M	.3L	.3L	.2L
T/E/A	.8H	.5M	.6M	.3L	.4M	.2L	.7H	.4M	.5M	.2L	.3L	.1L	.6M	.4M	.4M	.2L	.2L	.1L

**iii. Modified Rating** (NOTE: Modified score cannot exceed 1 or be less than 0.1.) **Vegetated Upland Buffer (VUB):** Area with 30% plant cover, 15% noxious weed or ANVS cover, and that is not subjected to periodic mechanical mowing or clearing (unless for weed control).

a) Is there an average 50 foot-wide vegetated upland buffer around 75% of the AA circumference? Y ☐ N ☒ If yes, add 0.1 to the score in ii above and adjust rating accordingly: **Modified Rating** .8H

**Comments:** Vegetated area greater than 5 acres with moderate level of biological activity and seasonal hydrology.

**14J. Groundwater Discharge/Recharge:** (check the appropriate indicators in i & ii below)

**i. Discharge Indicators**

- ☐ The AA is a slope wetland
- ☒ Springs or seeps are known or observed
- ☐ Vegetation growing during dormant season/drought
- ☐ Wetland occurs at the toe of a natural slope
- ☐ Seeps are present at the wetland edge
- ☐ AA permanently flooded during drought periods
- ☐ Wetland contains an outlet, but no inlet
- ☒ Shallow water table and the site is saturated to the surface
- ☐ Other:

**ii. Recharge Indicators**

- ☒ Permeable substrate present without underlying impeding layer
- ☐ Wetland contains inlet but no outlet
- ☐ Stream is a known 'losing' stream; discharge volume decreases
- ☐ Other:

**iii. Rating** (use the information from i and ii above and the table below to arrive at [check] the functional points and rating)

Criteria	Duration of saturation at AA Wetlands FROM GROUNDWATER DISCHARGE OR WITH WATER THAT IS RECHARGING THE GROUNDWATER SYSTEM			
	P/P	S/I	T	None
Groundwater Discharge or Recharge	1H	.7M	.4M	.1L
Insufficient Data/Information	NA			

**Comments:** Ponding was observed on site in 2014 and 2015.

**14K. Uniqueness:**

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

Replacement potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate		
	rare	common	abundant	rare	common	abundant	rare	common	abundant
Estimated relative abundance (#11)									
Low disturbance at AA (#12i)	1H	.9H	.8H	.8H	.6M	.5M	.5M	.4M	.3L
Moderate disturbance at AA (#12i)	.9H	.8H	.7M	.7M	.5M	.4M	.4M	.3L	.2L
High disturbance at AA (#12i)	.8H	.7H	.6M	.6M	.4M	.3L	.3L	.2L	.1L

**Comments:** Vegetation is becoming well established with tree populations colonizing the northern mitigation boundary.

**14L. Recreation/Education Potential:** (affords "bonus" points if AA provides recreation or education opportunity)

**i. Is the AA a known or potential rec./ed. site:** (check) ☒ Y ☐ N (if 'Yes' continue with the evaluation; if 'No' then click ☐ NA here and proceed to the overall summary and rating page)

**ii. Check categories that apply to the AA:** ☒ Educational/scientific study; ☒ Consumptive rec.; ☐ Non-consumptive rec.; ☐ Other

**iii. Rating** (use the matrix below to arrive at [check] the functional points and rating)

Known or Potential Recreation or Education Area	Known	Potential
Public ownership or public easement with general public access (no permission required)	.2H	.15H
Private ownership with general public access (no permission required)	.15H	.1M
Private or public ownership without general public access, or requiring permission for public access	.1M	.05L

**Comments:**

Permission is required for access to this site.

**General Site Notes**

FUNCTION & VALUE SUMMARY & OVERALL RATING FOR WETLAND/SITE #(S): Creation

Function & Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units: (Actual Points x Estimated AA Acreage)	Indicate the four most prominent functions with an asterisk (*)
A. Listed/Proposed T&E Species Habitat	L	0	1	0	<input type="checkbox"/>
B. MT Natural Heritage Program Species Habitat	L	.2	1	1.868	<input type="checkbox"/>
C. General Wildlife Habitat	M	.7	1	6.538	<input checked="" type="checkbox"/>
D. General Fish Habitat	NA	0	0	0	<input type="checkbox"/>
E. Flood Attenuation	M	.5	1	4.67	<input type="checkbox"/>
F. Short and Long Term Surface Water Storage	H	.8	1	7.472	<input checked="" type="checkbox"/>
G. Sediment/Nutrient/Toxicant Removal	H	.9	1	8.406	<input checked="" type="checkbox"/>
H. Sediment/Shoreline Stabilization	M	.6	1	5.604	<input type="checkbox"/>
I. Production Export/Food Chain Support	H	.8	1	7.472	<input checked="" type="checkbox"/>
J. Groundwater Discharge/Recharge	M	.7	1	6.538	<input type="checkbox"/>
K. Uniqueness	M	.4	1	3.736	<input type="checkbox"/>
L. Recreation/Education Potential (bonus points)	L	.05	NA	0.467	<input type="checkbox"/>
Totals:		5.65	10	52.771	
Percent of Possible Score			56.5 %		

**Category I Wetland:** (must satisfy **one** of the following criteria; otherwise go to Category II)

- ☐ Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**
- ☐ Score of 1 functional point for Uniqueness; **or**
- ☐ Score of 1 functional point for Flood Attenuation **and** answer to Question 14E.ii is "yes"; **or**
- ☐ Percent of possible score > 80% (round to nearest whole #).

**Category II Wetland:** (Criteria for Category I not satisfied **and** meets any **one** of the following criteria; otherwise go to Category IV)

- ☐ Score of 1 functional point for MT Natural Heritage Program Species Habitat; **or**
- ☐ Score of .9 or 1 functional point for General Wildlife Habitat; **or**
- ☐ Score of .9 or 1 functional point for General Fish Habitat; **or**
- ☐ "High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish/Aquatic Habitat; **or**
- ☐ Score of .9 functional point for Uniqueness; **or**
- ☐ Percent of possible score > 65% (round to nearest whole #).

**Category III Wetland:** (Criteria for Categories I, II, or IV not satisfied)



**Category IV Wetland:** (Criteria for Categories I or II are not satisfied and all of the following criteria are met; otherwise go to Category III)

- ☐ "Low" rating for Uniqueness; **and**
- ☐ Vegetated wetland component < 1 acre (do not include upland vegetated buffer); **and**
- ☐ Percent of possible score < 35% (round to nearest whole #).

**OVERALL ANALYSIS AREA RATING:**

(check appropriate category based on the criteria outlined above)

I	II	III	IV
---	----	-----	----



## MDT Montana Wetland Assessment Form (revised March 2008)

1. Project name	Easton Ranch	2. MDT project#	ST(X-34(14))	Control#	4866
3. Evaluation Date	6/26/2015	4. Evaluators	R. McEldowney, F. McNew		
5. Wetland/Site# (s)	Preservation				
6. Wetland Location(s): T	4N	R	9E	Sec1	32
Approx Stationing or Mileposts	NA				
Watershed	10070003	Watershed/County	Upper Yellowstone Watershed/Park County		
7. Evaluating Agency	Confluence for MDT				
Purpose of Evaluation			8. Wetland size acres		
<input type="checkbox"/> Wetlands potentially affected by MDT project <input type="checkbox"/> Mitigation Wetlands: pre-construction <input type="checkbox"/> Mitigation Wetlands: post construction <input checked="" type="checkbox"/> Other Preserved PSS/PFO/PEM Habitat			How assessed: Measured e.g. by GPS  9. Assessment area (AA) size (acres) 1.1 How assessed: Measured e.g. by GPS		

### 10. Classification of Wetland and Aquatic Habitats in AA

HGM Class (Brinson)	Class (Cowardin)	Modifier (Cowardin)	Water Regime	% of AA
Riverine	Scrub-Shrub Wetland		Seasonal/Intermittent	10
Riverine	Forested Wetland		Seasonal/Intermittent	20
Riverine	Emergent Wetland		Seasonal/Intermittent	70

11. Estimated Relative Abundance Common

### 12. General Condition of AA

i. Disturbance: (use matrix below to determine [circle] appropriate response – see instructions for Montana-listed noxious weed and aquatic nuisance vegetation species (ANVS) lists)

Conditions within AA	Predominant conditions adjacent to (within 500 feet of) AA		
	Managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings; and noxious weed or ANVS cover is <=15%.	Land not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to minor clearing; contains few roads or buildings; noxious weed or ANVS cover is <=30%.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >=30%.
AA occurs and is managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings; and noxious weed or ANVS cover is <=15%.	low disturbance	low disturbance	moderate disturbance
AA not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings; noxious weed or ANVS cover is <=30%.	moderate disturbance	moderate disturbance	high disturbance
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >=30%.	high disturbance	high disturbance	high disturbance

### Comments: (types of disturbance, intensity, season, etc)

AA consists of existing riverine PFO/PSS/PEM wetlands located adjacent to the created depressional wetlands and flood channel. AA and adjacent areas are managed in a natural state, disturbance is low.

### ii. Prominent noxious, aquatic nuisance, other exotic species:

Cirsium arvense and Cynoglossum officinale

### iii. Provide brief descriptive summary of AA and surrounding land use/habitat

AA contains small areas of existing PFO/PSS/PEM wetlands located at the northwest (Shields River) and southcentral ends of the mitigation area. The existing PFO/PEM habitat located at the southern end of the AA receives direct hydrologic inputs from the created flood channel. Both wetland features are bordered by created wetlands and the Shields River riparian corridor.

**13. Structural Diversity:** (based on number of "Cowardin" **vegetated** classes present [do not include unvegetated classes], see #10 above)

Existing # of "Cowardin" Vegetated Classes in AA	Initial Rating	Is current management preventing (passive) existence of additional vegetated classes?		Modified Rating
>=3 (or 2 if 1 is forested) classes	H	NA	NA	NA
2 (or 1 if forested) classes	M	NA	NA	NA
1 class, but not a monoculture	M	<NO	YES>	L
1 class, monoculture (1 species comprises >=90% of total cover)	L	NA	NA	NA

**Comments:** PEM, PFO, and PSS vegetated communities are present on site.

## SECTION PERTAINING to FUNCTIONS VALUES ASSESSMEN

### 14A. Habitat for Federally Listed or Proposed Threatened or Endangered Plants or Animals:

i. AA is Documented (D) or Suspected (S) to contain (check one based on definitions contained in instructions):

Primary or critical habitat (list species) ☐ D ☐ S

Secondary habitat (list Species) ☐ D ☐ S

Incidental habitat (list species) ☐ D ☐ S

No usable habitat ☒ S

ii. **Rating** (use the conclusions from i above and the matrix below to arrive at [check] the functional points and rating)

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	None
Functional Points and Rating	1H	.9H	.8H	.7M	.3L	.1L	0L

Sources for documented use

### 14B. Habitat for plant or animals rated S1, S2, or S3 by the Montana Natural Heritage Program: (not including species listed in 14A above)

i. AA is Documented (D) or Suspected (S) to contain (check one based on definitions contained in instructions):

Primary or critical habitat (list species) ☐ D ☐ S

Secondary habitat (list Species) ☐ D ☐ S

Incidental habitat (list species) ☒ D ☐ S Golden Eagle (S3)

No usable habitat ☐ S

ii. **Rating** (use the conclusions from i above and the matrix below to arrive at [check] the functional points and rating)

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	None
<b>S1 Species:</b> Functional Points and Rating	1H	.8H	.7M	.6M	.2L	.1L	0L
<b>S2 and S3 Species:</b> Functional Points and Rating	.9H	.7M	.6M	.5M	.2L	.1L	0L

Sources for documented use

MTNHP, 2013 field observations.

#### 14C. General Wildlife Habitat Rating:

- i. Evidence of overall wildlife use in the AA (check substantial, moderate, or low based on supporting evidence):

Moderate

**Substantial** (based on any of the following [check]):

- ☐ observations of abundant wildlife #s or high species diversity (during any period)
- ☐ abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ presence of extremely limiting habitat features not available in the surrounding area
- ☐ interviews with local biologists with knowledge of the AA

**Minimal** (based on any of the following [check]):

- ☐ few or no wildlife observations during peak use periods
- ☐ little to no wildlife sign
- ☐ sparse adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

**Moderate** (based on any of the following [check]):

- ☐ observations of scattered wildlife groups or individuals or relatively few species during peak periods
- ☒ common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ adequate adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

- ii. **Wildlife** habitat features (Working from top to bottom, check appropriate AA attributes in matrix to arrive at rating. Structural diversity is from #13. For class cover to be considered evenly distributed, the most and least prevalent **vegetated** classes must be within 20% of each other in terms of their percent composition of the AA (see #10). Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; and A = absent [see instructions for further definitions of these terms])

Structural diversity (see #13)	High								Moderate								Low			
	Even				Uneven				Even				Uneven				Even			
Class cover distribution (all vegetated classes)	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Duration of surface water in 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Low disturbance at AA (see #12)	E	E	E	H	E	E	H	H	E	H	H	M	E	H	M	M	E	H	M	M
Moderate disturbance at AA (see #12)	H	H	H	H	H	H	H	M	H	H	M	M	H	M	M	L	H	M	L	L
High disturbance at AA (see #12)	M	M	M	L	M	M	L	L	M	M	L	L	M	L	L	L	L	L	L	L

- iii. **Rating** (use the conclusions from i and ii above and the matrix below to arrive at [check] the functional points and rating)

Evidence of wildlife use (i)	Wildlife habitat features rating (ii)											
	Exceptional			High			Moderate			Low		
Substantial		1E			.9H			.8H			.7M	
Moderate		.9H			.7M			.5M			.3L	
Minimal		.6M			.4M			.2L			.1L	

#### Comments

Moderate use of site by moose, deer, golden eagle, and other avian species. There is an active bald eagle (*Haliaeetus leucocephalus*) nest directly southwest of the site on the west side of the Shields River the project site is within the primary habitat zone for bald eagles (L. Urban. MDT. personal communications. 2015).

- 14D. **General Fish Habitat Rating:** (Assess this function if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [i.e., fish use is precluded by perched culvert or other barrier, etc.]. If the AA is not used by fish, fish use is not restorable due to habitat constraints, or is not desired from a management perspective [such as fish entrapped in a canal], then check

☒ NA here and proceed to 14E.)

- i. **Habitat Quality and Known / Suspected Fish Species in AA** (use matrix to arrive at [check the functional points and rating])

Duration of surface water in AA	Permanent / Perennial						Seasonal / Intermittent						Temporary / Ephemeral					
	Optimal		Adequate		Poor		Optimal		Adequate		Poor		Optimal		Adequate		Poor	
Aquatic hiding / resting / escape cover	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S
Thermal cover optimal / suboptimal	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S
FWP Tier I fish species	1E	.9H	.8H	.7M	.6M	.5M	.9H	.8H	.7M	.6M	.5M	.4M	.7M	.6M	.5M	.4M	.3L	.3L
FWP Tier II or Native Game fish species	.9H	.8H	.7M	.6M	.5M	.5M	.8H	.7M	.6M	.5M	.4M	.4M	.6M	.5M	.4M	.3L	.2L	.2L
FWP Tier III or Introduced Game fish	.8H	.7M	.6M	.5M	.5M	.4M	.7M	.6M	.5M	.4M	.4M	.3L	.5M	.4M	.3L	.2L	.2L	.1L
FWP Non-Game Tier IV or No fish species	.5M	.5M	.5M	.4M	.4M	.3L	.4M	.4M	.4M	.3L	.3L	.2L	.2L	.2L	.2L	.1L	.1L	.1L

Sources used for identifying fish sp. potentially found in AA:

ii. **Modified Rating** (NOTE: Modified score cannot exceed 1 or be less than 0.1)

a) Is fish use of the AA significantly reduced by a culvert, dike, or other man-made structure or activity or is the waterbody included on the current final MDEQ list of waterbodies in need of TMDL development with listed "Probable Impaired Uses" including cold or warm water fishery or aquatic life support, or do aquatic nuisance plant or animal species (see Appendix E) occur in fish habitat? Y ☐ N ☒ If yes, reduce score in i above by 0.1: **Modified Rating**

b) Does the AA contain a documented spawning area or other critical habitat feature (i.e., sanctuary pool, upwelling area, etc.- specify in comments) for native fish or introduced game fish? ☐ Y ☒ N If yes, add 0.1 to the adjusted score in i or iia above:

**Modified Rating**

iii. **Final Score and Rating:**


**Comments:** No fish habitat on site.

**14E. Flood Attenuation:** (Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA are not flooded from in-channel or overbank flow, click ☐ NA here and proceed to 14F.)

i. **Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

Estimated or Calculated Entrenchment (Rosgen 1994, 1996)	Slightly entrenched - C, D, E stream types			Moderately entrenched - B stream type			Entrenched-A, F, G stream types		
% of flooded wetland classified as forested and/or scrub/shrub	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains <b>no outlet or restricted outlet</b>	1H	.9H	.6M	.8H	.7M	.5M	.4M	.3L	.2L
AA contains <b>unrestricted outlet</b>	.9H	.8H	.5M	.7M	.6M	.4M	.3L	.2L	.1L

Slightly Entrenched ER = >2.2			Moderately Entrenched ER = 1.41 - 2.2		Entrenched ER = 1.0 - 1.4	
C stream type	D stream type	E stream type	B stream type		A stream type	F stream type



**Floodprone width**  / **Bankfull width**  = **Entrenchment ratio**

ii. Are 10 acres of wetland in the AA subject to flooding AND are man-made features which may be significantly damaged by floods located within 0.5 mile downstream of the AA (check)? Y ☐ N ☒

**Comments:**

Approximately 30% of the preservation AA contains forested and/or scrub/shrub wetland with surface water outlet to the south into relic isolated channel. The Shields River is slightly entrenched at this location.

**14F. Short and Long Term Surface Water Storage:** (Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow. If no wetlands in the AA are subject to flooding or ponding, click ☐ NA here and proceed to 14G.)

i. **Rating** (Working from top to bottom, use the matrix below to arrive at [check] the functional points and rating. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral [see instructions for further definitions of these terms].)

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding	>5 acre feet			1.1 to 5 acre feet			≤1 acre foot		
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond <b>≥ 5 out of 10 years</b>	1H	.9H	.8H	.8H	.6M	.5M	.4M	.3L	.2L
Wetlands in AA flood or pond <b>&lt; 5 out of 10 years</b>	.9H	.8H	.7M	.7M	.5M	.4M	.3L	.2L	.1L

**Comments:** (1.10 acre of preserved wetland) x (approximate average of 1.0 ft. of inundation during high water) = 1.10 acre feet



**14G. Sediment/Nutrient/Toxicant Retention and Removal:** (Applies to wetlands with potential to receive sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, click ☐ **NA** here and proceed to 14H.)

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating [H = high, M = moderate, or L = low])

Sediment, nutrient, and toxicant input levels within AA	AA receives or surrounding land use with potential to deliver levels of sediments, nutrients, or compounds at levels such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use with potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% cover of wetland vegetation in AA	≥ 70%		< 70%		≥ 70%		< 70%	
Evidence of flooding / ponding in AA	Yes	No	Yes	No	Yes	No	Yes	No
AA contains <b>no or restricted</b> outlet	1H	.8H	.7M	.5M	.5M	.4M	.3L	.2L
AA contains <b>unrestricted</b> outlet	.9H	.7M	.6M	.4M	.4M	.3L	.2L	.1L

**Comments:** Wetland vegetation cover exceeds 70%. AA contains restricted outlet.

**14H Sediment/Shoreline Stabilization:** (Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action. If 14H does not apply, click ☒ **NA** here and proceed to 14I.)

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

% Cover of <b>wetland</b> streambank or shoreline by species with stability ratings of 6 (see <b>Appendix F</b> ).	Duration of surface water adjacent to rooted vegetation					
	Permanent / Perennial		Seasonal / Intermittent		Temporary / Ephemeral	
≥ 65%	1H		.9H		.7M	
35-64%	.7M		.6M		.5M	
< 35%	.3L		.2L		.1L	

No shoreline in the project area.

**Comments:**

#### 14I. Production Export/Food Chain Support:

**i. Level of Biological Activity** (synthesis of wildlife and fish habitat ratings [check])

General Fish Habitat Rating (14D.iii.)	General Wildlife Habitat Rating (14C.iii.)					
	E/H		M		L	
E/H	H		H		M	
M	H		M		M	
L	M		M		L	
N/A	H		M		L	

**ii. Rating** (Working from top to bottom, use the matrix below to arrive at [check] the functional points and rating. Factor A = acreage of vegetated wetland component in the AA; Factor B = level of biological activity rating from above (14I.i.); Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to duration of surface water in the AA, where P/P, S/I, and T/E are as previously defined, and A = "absent" [see instructions for further definitions of these terms].)

A	Vegetated component >5 acres						Vegetated component 1-5 acres						Vegetated component <1 acre					
	High		Moderate		Low		High		Moderate		Low		High		Moderate		Low	
	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
P/P	1E	.7H	.8H	.5M	.6M	.4M	.9H	.6M	.7H	.4M	.5M	.3L	.8H	.6M	.6M	.4M	.3L	.2L
S/I	.9H	.6M	.7H	.4M	.5M	.3L	.8H	.5M	.6M	.3L	.4M	.2L	.7H	.5M	.5M	.3L	.3L	.2L
T/E/A	.8H	.5M	.6M	.3L	.4M	.2L	.7H	.4M	.5M	.2L	.3L	.1L	.6M	.4M	.4M	.2L	.2L	.1L

**iii. Modified Rating** (NOTE: Modified score cannot exceed 1 or be less than 0.1.) **Vegetated Upland Buffer (VUB):** Area with 30% plant cover, 15% noxious weed or ANVS cover, and that is not subjected to periodic mechanical mowing or clearing (unless for weed control).

a) Is there an average 50 foot-wide vegetated upland buffer around 75% of the AA circumference? Y ☒ N ☐ If yes, add 0.1 to the score in ii above and adjust rating accordingly: **Modified Rating** .9H

**Comments:** There is a restricted surface water outlet to the south.

**14J. Groundwater Discharge/Recharge:** (check the appropriate indicators in i & ii below)

**i. Discharge Indicators**

- ☐ The AA is a slope wetland
- ☐ Springs or seeps are known or observed
- ☐ Vegetation growing during dormant season/drought
- ☐ Wetland occurs at the toe of a natural slope
- ☐ Seeps are present at the wetland edge
- ☐ AA permanently flooded during drought periods
- ☐ Wetland contains an outlet, but no inlet
- ☒ Shallow water table and the site is saturated to the surface
- ☐ Other:

**ii. Recharge Indicators**

- ☒ Permeable substrate present without underlying impeding layer
- ☐ Wetland contains inlet but no outlet
- ☐ Stream is a known 'losing' stream; discharge volume decreases
- ☐ Other:

**iii. Rating** (use the information from i and ii above and the table below to arrive at [check] the functional points and rating)

Criteria	Duration of saturation at AA Wetlands FROM GROUNDWATER DISCHARGE OR WITH WATER THAT IS RECHARGING THE GROUNDWATER SYSTEM			
	P/P	S/I	T	None
Groundwater Discharge or Recharge	1H	.7M	.4M	.1L
Insufficient Data/Information	NA			

Comments:

**14K. Uniqueness:**

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

Replacement potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate		
	rare	common	abundant	rare	common	abundant	rare	common	abundant
Low disturbance at AA (#12i)	1H	.9H	.8H	.8H	.6M	.5M	.5M	.4M	.3L
Moderate disturbance at AA (#12i)	.9H	.8H	.7M	.7M	.5M	.4M	.4M	.3L	.2L
High disturbance at AA (#12i)	.8H	.7H	.6M	.6M	.4M	.3L	.3L	.2L	.1L

Comments:  Site disturbance is low and structural diversity is high.

**14L. Recreation/Education Potential:** (affords "bonus" points if AA provides recreation or education opportunity)

**i. Is the AA a known or potential rec./ed. site:** (check) Y ☐ N ☒ (if 'Yes' continue with the evaluation; if 'No' then click ☒ NA here and proceed to the overall summary and rating page)

**ii. Check categories that apply to the AA:** ☐ Educational/scientific study; ☒ Consumptive rec.; ☒ Non-consumptive rec.; ☐ Other

**iii. Rating** (use the matrix below to arrive at [check] the functional points and rating)

Known or Potential Recreation or Education Area	Known	Potential
Public ownership or public easement with general public access (no permission required)	.2H	.15H
Private ownership with general public access (no permission required)	.15H	.1M
Private or public ownership without general public access, or requiring permission for public access	.1M	.05L

Comments:

Permission is required for site access.

**General Site Notes**

FUNCTION & VALUE SUMMARY & OVERALL RATING FOR WETLAND/SITE #(S): Preservation

Function & Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units: (Actual Points x Estimated AA Acreage)	Indicate the four most prominent functions with an asterisk (*)
A. Listed/Proposed T&E Species Habitat	L	0	1	0	<input type="checkbox"/>
B. MT Natural Heritage Program Species Habitat	L	.2	1	0.22	<input type="checkbox"/>
C. General Wildlife Habitat	H	.9	1	0.99	<input checked="" type="checkbox"/>
D. General Fish Habitat	NA	0	0	0	<input type="checkbox"/>
E. Flood Attenuation	H	.9	1	0.99	<input type="checkbox"/>
F. Short and Long Term Surface Water Storage	M	.6	1	0.66	<input type="checkbox"/>
G. Sediment/Nutrient/Toxicant Removal	H	1	1	1.1	<input checked="" type="checkbox"/>
H. Sediment/Shoreline Stabilization	NA	0	0	0	<input type="checkbox"/>
I. Production Export/Food Chain Support	H	.9	1	0.99	<input checked="" type="checkbox"/>
J. Groundwater Discharge/Recharge	M	.7	1	0.77	<input checked="" type="checkbox"/>
K. Uniqueness	M	.6	1	0.66	<input type="checkbox"/>
L. Recreation/Education Potential (bonus points)	L	.05	NA	0.055	<input type="checkbox"/>
Totals:		5.85	9	6.435	
Percent of Possible Score			65 %		

**Category I Wetland:** (must satisfy **one** of the following criteria; otherwise go to Category II)

- ☐ Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**
- ☐ Score of 1 functional point for Uniqueness; **or**
- ☐ Score of 1 functional point for Flood Attenuation **and** answer to Question 14E.ii is "yes"; **or**
- ☐ Percent of possible score > 80% (round to nearest whole #).

**Category II Wetland:** (Criteria for Category I not satisfied **and** meets any **one** of the following criteria; otherwise go to Category IV)

- ☐ Score of 1 functional point for MT Natural Heritage Program Species Habitat; **or**
- ☒ Score of .9 or 1 functional point for General Wildlife Habitat; **or**
- ☐ Score of .9 or 1 functional point for General Fish Habitat; **or**
- ☐ "High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish/Aquatic Habitat; **or**
- ☐ Score of .9 functional point for Uniqueness; **or**
- ☒ Percent of possible score > 65% (round to nearest whole #).

**Category III Wetland:** (Criteria for Categories I, II, or IV not satisfied)

☐

**Category IV Wetland:** (Criteria for Categories I or II are not satisfied and all of the following criteria are met; otherwise go to Category III)

- ☐ "Low" rating for Uniqueness; **and**
- ☐ Vegetated wetland component < 1 acre (do not include upland vegetated buffer); **and**
- ☐ Percent of possible score < 35% (round to nearest whole #).

**OVERALL ANALYSIS AREA RATING:**

(check appropriate category based on the criteria outlined above)

I	II	III	IV
---	----	-----	----

# MDT Montana Wetland Assessment Form (revised March 2008)

1. Project name  2. MDT project#  Control#

3. Evaluation Date  4. Evaluators  5. Wetland/Site# (s)

6. Wetland Location(s): T  R  Sec1  T  R  Sec2

Approx Stationing or Mileposts

Watershed  Watershed/County

7. Evaluating Agency

8. Wetland size acres

Purpose of Evaluation

☐ Wetlands potentially affected by MDT project

☐ Mitigation Wetlands: pre-construction

☐ Mitigation Wetlands: post construction

☒ Other

9. Assessment area (AA) size (acres)

How assessed:

## 10. Classification of Wetland and Aquatic Habitats in AA

HGM Class (Brinson)	Class (Cowardin)	Modifier (Cowardin)	Water Regime	% of AA
Riverine	Emergent Wetland	Excavated	Seasonal/Intermittent	100
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

11. Estimated Relative Abundance

## 12. General Condition of AA

i. Disturbance: (use matrix below to determine [circle] appropriate response – see instructions for Montana-listed noxious weed and aquatic nuisance vegetation species (ANVS) lists)

Conditions within AA	Predominant conditions adjacent to (within 500 feet of) AA		
	Managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings; and noxious weed or ANVS cover is <=15%.	Land not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to minor clearing; contains few roads or buildings; noxious weed or ANVS cover is <=30%.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >=30%.
AA occurs and is managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings; and noxious weed or ANVS cover is <=15%.	<input type="text" value="low disturbance"/>	<input type="text" value="low disturbance"/>	<input type="text" value="moderate disturbance"/>
AA not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings; noxious weed or ANVS cover is <=30%.	<input type="text" value="moderate disturbance"/>	<input type="text" value="moderate disturbance"/>	<input type="text" value="high disturbance"/>
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >=30%.	<input type="text" value="high disturbance"/>	<input type="text" value="high disturbance"/>	<input type="text" value="high disturbance"/>

## Comments: (types of disturbance, intensity, season, etc)

Limited agriculture (hay) and ranch structures to the east. Undeveloped riparian corridor and herbaceous uplands to north, south, and west. Two species of noxious weeds were present within the AA. The AA is managed in a natural state, as are most of the lands within 500 feet of the AA.

## ii. Prominent noxious, aquatic nuisance, other exotic species:

## iii. Provide brief descriptive summary of AA and surrounding land use/habitat

The AA consists of one constructed secondary stream channel which bisects the mitigation area. The channel is active during high flow events, is seasonally inundated by shallow ground water early in the growing season and has developed wetland characteristics.

**13. Structural Diversity:** (based on number of "Cowardin" **vegetated** classes present [do not include unvegetated classes], see #10 above)

Existing # of "Cowardin" Vegetated Classes in AA	Initial Rating	Is current management preventing (passive) existence of additional vegetated classes?		Modified Rating
>=3 (or 2 if 1 is forested) classes	H	NA	NA	NA
2 (or 1 if forested) classes	M	NA	NA	NA
1 class, but not a monoculture	M	<NO	YES>	L
1 class, monoculture (1 species comprises >=90% of total cover)	L	NA	NA	NA

**Comments:** Planted shrubs along channel are surviving but have yet to develop a shrub community, emergent vegetation present.

## SECTION PERTAINING to FUNCTIONS VALUES ASSESSMEN

### 14A. Habitat for Federally Listed or Proposed Threatened or Endangered Plants or Animals:

i. AA is Documented (D) or Suspected (S) to contain (check one based on definitions contained in instructions):

Primary or critical habitat (list species) ☐ D ☐ S

Secondary habitat (list Species) ☐ D ☐ S

Incidental habitat (list species) ☐ D ☐ S

No usable habitat ☒ S

ii. **Rating** (use the conclusions from i above and the matrix below to arrive at [check] the functional points and rating)

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	None
Functional Points and Rating	1H	.9H	.8H	.7M	.3L	.1L	0L

Sources for documented use

### 14B. Habitat for plant or animals rated S1, S2, or S3 by the Montana Natural Heritage Program: (not including species listed in 14A above)

i. AA is Documented (D) or Suspected (S) to contain (check one based on definitions contained in instructions):

Primary or critical habitat (list species) ☐ D ☐ S

Secondary habitat (list Species) ☐ D ☐ S

Incidental habitat (list species) ☒ D ☐ S Golden Eagle (S3)

No usable habitat ☐ S

ii. **Rating** (use the conclusions from i above and the matrix below to arrive at [check] the functional points and rating)

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	None
<b>S1 Species:</b> Functional Points and Rating	1H	.8H	.7M	.6M	.2L	.1L	0L
<b>S2 and S3 Species:</b> Functional Points and Rating	.9H	.7M	.6M	.5M	.2L	.1L	0L

Sources for documented use

MTNHP;observed on site in 2013



**14C. General Wildlife Habitat Rating:**

i. Evidence of overall wildlife use in the AA (check substantial, moderate, or low based on supporting evidence):

Moderate

**Substantial** (based on any of the following [check]):

- ☐ observations of abundant wildlife #s or high species diversity (during any period)
- ☐ abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ presence of extremely limiting habitat features not available in the surrounding area
- ☐ interviews with local biologists with knowledge of the AA

**Minimal** (based on any of the following [check]):

- ☐ few or no wildlife observations during peak use periods
- ☐ little to no wildlife sign
- ☐ sparse adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

**Moderate** (based on any of the following [check]):

- ☐ observations of scattered wildlife groups or individuals or relatively few species during peak periods
- ☒ common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- ☐ adequate adjacent upland food sources
- ☐ interviews with local biologists with knowledge of the AA

ii. **Wildlife** habitat features (Working from top to bottom, check appropriate AA attributes in matrix to arrive at rating. Structural diversity is from #13. For class cover to be considered evenly distributed, the most and least prevalent **vegetated** classes must be within 20% of each other in terms of their percent composition of the AA (see #10). Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; and A = absent [see instructions for further definitions of these terms])

Structural diversity (see #13)	High								Moderate								Low			
	Even				Uneven				Even				Uneven				Even			
Class cover distribution (all vegetated classes)	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Duration of surface water in 10% of AA																				
Low disturbance at AA (see #12)	E	E	E	H	E	E	H	H	E	H	H	M	E	H	M	M	E	H	M	M
Moderate disturbance at AA (see #12)	H	H	H	H	H	H	H	M	H	H	M	M	H	M	M	L	H	M	L	L
High disturbance at AA (see #12)	M	M	M	L	M	M	L	L	M	M	L	L	M	L	L	L	L	L	L	L

iii. **Rating** (use the conclusions from i and ii above and the matrix below to arrive at [check] the functional points and rating)

Evidence of wildlife use (i)	Wildlife habitat features rating (ii)											
	Exceptional			High			Moderate			Low		
Substantial		1E			.9H			.8H			.7M	
Moderate		.9H			.7M			.5M			.3L	
Minimal		.6M			.4M			.2L			.1L	

**Comments**

AA has frequent deer and moose sightings. There is an active bald eagle (*Haliaeetus leucocephalus*) nest directly southwest of the site on the west side of the Shields River the project site is within the primary habitat zone for bald eagles (L. Urban. MDT. personal communications. 2015).

**14D. General Fish Habitat Rating:** (Assess this function if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [i.e., fish use is precluded by perched culvert or other barrier, etc.]. If the AA is not used by fish, fish use is not restorable due to habitat constraints, or is not desired from a management perspective [such as fish entrapped in a canal], then check

☒ **NA** here and proceed to 14E.)

i. **Habitat Quality and Known / Suspected Fish Species in AA** (use matrix to arrive at [check the functional points and rating])

Duration of surface water in AA	Permanent / Perennial						Seasonal / Intermittent						Temporary / Ephemeral					
	Optimal		Adequate		Poor		Optimal		Adequate		Poor		Optimal		Adequate		Poor	
Aquatic hiding / resting / escape cover																		
Thermal cover optimal / suboptimal	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S	O	S
FWP Tier I fish species	1E	.9H	.8H	.7M	.6M	.5M	.9H	.8H	.7M	.6M	.5M	.4M	.7M	.6M	.5M	.4M	.3L	.3L
FWP Tier II or Native Game fish species	.9H	.8H	.7M	.6M	.5M	.5M	.8H	.7M	.6M	.5M	.4M	.4M	.6M	.5M	.4M	.3L	.2L	.2L
FWP Tier III or Introduced Game fish	.8H	.7M	.6M	.5M	.5M	.4M	.7M	.6M	.5M	.4M	.4M	.3L	.5M	.4M	.3L	.2L	.2L	.1L
FWP Non-Game Tier IV or No fish species	.5M	.5M	.5M	.4M	.4M	.3L	.4M	.4M	.4M	.3L	.3L	.2L	.2L	.2L	.2L	.1L	.1L	.1L

Sources used for identifying fish sp. potentially found in AA:

ii. **Modified Rating** (NOTE: Modified score cannot exceed 1 or be less than 0.1)

a) Is fish use of the AA significantly reduced by a culvert, dike, or other man-made structure or activity or is the waterbody included on the current final MDEQ list of waterbodies in need of TMDL development with listed "Probable Impaired Uses" including cold or warm water fishery or aquatic life support, or do aquatic nuisance plant or animal species (see Appendix E) occur in fish habitat? Y ☐ N ☒ If yes, reduce score in i above by 0.1: **Modified Rating**

b) Does the AA contain a documented spawning area or other critical habitat feature (i.e., sanctuary pool, upwelling area, etc.- specify in comments) for native fish or introduced game fish? ☐ Y ☒ N If yes, add 0.1 to the adjusted score in i or iia above:

**Modified Rating**

iii. **Final Score and Rating:**


**Comments:** Although activated during high-flow events within the Shields River, no permanent fish habitat is present within AA.

**14E. Flood Attenuation:** (Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA are not flooded from in-channel or overbank flow, click ☐ NA here and proceed to 14F.)

i. **Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

Estimated or Calculated Entrenchment (Rosgen 1994, 1996)	Slightly entrenched - C, D, E stream types			Moderately entrenched - B stream type			Entrenched-A, F, G stream types		
% of flooded wetland classified as forested and/or scrub/shrub	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains <b>no outlet or restricted outlet</b>	1H	.9H	.6M	.8H	.7M	.5M	.4M	.3L	.2L
AA contains <b>unrestricted outlet</b>	.9H	.8H	.5M	.7M	.6M	.4M	.3L	.2L	.1L

Slightly Entrenched ER = >2.2			Moderately Entrenched ER = 1.41 - 2.2		Entrenched ER = 1.0 - 1.4	
C stream type	D stream type	E stream type	B stream type		A stream type	F stream type



**Floodprone width**  / **Bankfull width**  = **Entrenchment ratio**

ii. Are 10 acres of wetland in the AA subject to flooding AND are man-made features which may be significantly damaged by floods located within 0.5 mile downstream of the AA (check)? Y ☐ N ☒

**Comments:**

Outlet is restricted. AA subject to overflow from Shields River and empties into old meanders of the Shields River at the south end of AA.

**14F. Short and Long Term Surface Water Storage:** (Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow. If no wetlands in the AA are subject to flooding or ponding, click ☐ NA here and proceed to 14G.)

i. **Rating** (Working from top to bottom, use the matrix below to arrive at [check] the functional points and rating. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral [see instructions for further definitions of these terms].)

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding	>5 acre feet			1.1 to 5 acre feet			≤1 acre foot		
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond <b>≥ 5 out of 10 years</b>	1H	.9H	.8H	.8H	.6M	.5M	.4M	.3L	.2L
Wetlands in AA flood or pond <b>&lt; 5 out of 10 years</b>	.9H	.8H	.7M	.7M	.5M	.4M	.3L	.2L	.1L

**Comments:** (1.56 acre of restoration) x (average 1 ft. ponding/flow at high water) = 1.56 acre feet

**14G. Sediment/Nutrient/Toxicant Retention and Removal:** (Applies to wetlands with potential to receive sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, click ☐ **NA** here and proceed to 14H.)

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating [H = high, M = moderate, or L = low])

Sediment, nutrient, and toxicant input levels within AA	AA receives or surrounding land use with potential to deliver levels of sediments, nutrients, or compounds at levels such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use with potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% cover of wetland vegetation in AA	≥ 70%		< 70%		≥ 70%		< 70%	
Evidence of flooding / ponding in AA	Yes	No	Yes	No	Yes	No	Yes	No
AA contains <b>no or restricted</b> outlet	1H	.8H	.7M	.5M	.5M	.4M	.3L	.2L
AA contains <b>unrestricted</b> outlet	.9H	.7M	.6M	.4M	.4M	.3L	.2L	.1L

**Comments:** Cover in AA is greater than 70% and outlet is topographically restricted.

**14H Sediment/Shoreline Stabilization:** (Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action. If 14H does not apply, click ☐ **NA** here and proceed to 14I.)

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

% Cover of <b>wetland</b> streambank or shoreline by species with stability ratings of 6 (see <b>Appendix F</b> ).	Duration of surface water adjacent to rooted vegetation					
	Permanent / Perennial		Seasonal / Intermittent		Temporary / Ephemeral	
≥ 65%	1H		.9H		.7M	
35-64%	.7M		.6M		.5M	
< 35%	.3L		.2L		.1L	

**Comments:**

Increased vegetation development from 2013 to 2015 of species with high stability ratings including Salix, Carex and Juncus.

#### 14I. Production Export/Food Chain Support:

**i. Level of Biological Activity** (synthesis of wildlife and fish habitat ratings [check])

General Fish Habitat Rating (14D.iii.)	General Wildlife Habitat Rating (14C.iii.)					
	E/H		M		L	
E/H	H		H		M	
M	H		M		M	
L	M		M		L	
N/A	H		M		L	

**ii. Rating** (Working from top to bottom, use the matrix below to arrive at [check] the functional points and rating. Factor A = acreage of vegetated wetland component in the AA; Factor B = level of biological activity rating from above (14I.i.); Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to duration of surface water in the AA, where P/P, S/I, and T/E are as previously defined, and A = "absent" [see instructions for further definitions of these terms].)

A	Vegetated component >5 acres						Vegetated component 1-5 acres						Vegetated component <1 acre					
	High		Moderate		Low		High		Moderate		Low		High		Moderate		Low	
	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
P/P	1E	.7H	.8H	.5M	.6M	.4M	.9H	.6M	.7H	.4M	.5M	.3L	.8H	.6M	.6M	.4M	.3L	.2L
S/I	.9H	.6M	.7H	.4M	.5M	.3L	.8H	.5M	.6M	.3L	.4M	.2L	.7H	.5M	.5M	.3L	.3L	.2L
T/E/A	.8H	.5M	.6M	.3L	.4M	.2L	.7H	.4M	.5M	.2L	.3L	.1L	.6M	.4M	.4M	.2L	.2L	.1L

**iii. Modified Rating** (NOTE: Modified score cannot exceed 1 or be less than 0.1.) **Vegetated Upland Buffer (VUB):** Area with 30% plant cover, 15% noxious weed or ANVS cover, and that is not subjected to periodic mechanical mowing or clearing (unless for weed control).

a) Is there an average 50 foot-wide vegetated upland buffer around 75% of the AA circumference? Y ☒ N ☐ If yes, add 0.1 to the score in ii above and adjust rating accordingly: **Modified Rating** .7M

**Comments:**

Channel is seasonally inundated and has a restricted outlet at the southern end of the mitigation site.

**14J. Groundwater Discharge/Recharge:** (check the appropriate indicators in i & ii below)

**i. Discharge Indicators**

- ☐ The AA is a slope wetland
- ☒ Springs or seeps are known or observed
- ☒ Vegetation growing during dormant season/drought
- ☐ Wetland occurs at the toe of a natural slope
- ☐ Seeps are present at the wetland edge
- ☐ AA permanently flooded during drought periods
- ☐ Wetland contains an outlet, but no inlet
- ☒ Shallow water table and the site is saturated to the surface
- ☐ Other:

**ii. Recharge Indicators**

- ☒ Permeable substrate present without underlying impeding layer
- ☐ Wetland contains inlet but no outlet
- ☐ Stream is a known 'losing' stream; discharge volume decreases
- ☐ Other:

**iii. Rating** (use the information from i and ii above and the table below to arrive at [check] the functional points and rating)

Criteria	Duration of saturation at AA Wetlands FROM GROUNDWATER DISCHARGE OR WITH WATER THAT IS RECHARGING THE GROUNDWATER SYSTEM			
	P/P	S/I	T	None
Groundwater Discharge or Recharge	1H	.7M	.4M	.1L
Insufficient Data/Information	NA			

**Comments:** Channel is intermittently inundated by shallow groundwater and high flows from the Shields River.

**14K. Uniqueness:**

**i. Rating** (working from top to bottom, use the matrix below to arrive at [check] the functional points and rating)

Replacement potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate		
	rare	common	abundant	rare	common	abundant	rare	common	abundant
Low disturbance at AA (#12i)	1H	.9H	.8H	.8H	.6M	.5M	.5M	.4M	.3L
Moderate disturbance at AA (#12i)	.9H	.8H	.7M	.7M	.5M	.4M	.4M	.3L	.2L
High disturbance at AA (#12i)	.8H	.7H	.6M	.6M	.4M	.3L	.3L	.2L	.1L

**Comments:** Emergent wetland within seasonal flood channel. Common wetland type within basin with 10-50% of area wetlands similar to

**14L. Recreation/Education Potential:** (affords "bonus" points if AA provides recreation or education opportunity)

**i. Is the AA a known or potential rec./ed. site:** (check) ☒ Y ☐ N (if 'Yes' continue with the evaluation; if 'No' then click ☐ NA here and proceed to the overall summary and rating page)

**ii. Check categories that apply to the AA:** ☒ Educational/scientific study; ☐ Consumptive rec.; ☐ Non-consumptive rec.; ☐ Other

**iii. Rating** (use the matrix below to arrive at [check] the functional points and rating)

Known or Potential Recreation or Education Area	Known	Potential
Public ownership or public easement with general public access (no permission required)	.2H	.15H
Private ownership with general public access (no permission required)	.15H	.1M
Private or public ownership without general public access, or requiring permission for public access	.1M	.05L

**Comments:**

Permission is required for site access.

**General Site Notes**

FUNCTION & VALUE SUMMARY & OVERALL RATING FOR WETLAND/SITE #(S): Restoration

Function & Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units: (Actual Points x Estimated AA Acreage)	Indicate the four most prominent functions with an asterisk (*)
A. Listed/Proposed T&E Species Habitat	L	0	1	0	<input type="checkbox"/>
B. MT Natural Heritage Program Species Habitat	L	.2	1	0.312	<input type="checkbox"/>
C. General Wildlife Habitat	M	.7	1	1.092	<input type="checkbox"/>
D. General Fish Habitat	NA	0	0	0	<input type="checkbox"/>
E. Flood Attenuation	M	.6	1	0.936	<input type="checkbox"/>
F. Short and Long Term Surface Water Storage	M	.6	1	0.936	<input type="checkbox"/>
G. Sediment/Nutrient/Toxicant Removal	H	1	1	1.56	<input checked="" type="checkbox"/>
H. Sediment/Shoreline Stabilization	H	.9	1	1.404	<input checked="" type="checkbox"/>
I. Production Export/Food Chain Support	M	.7	1	1.092	<input checked="" type="checkbox"/>
J. Groundwater Discharge/Recharge	M	.7	1	1.092	<input checked="" type="checkbox"/>
K. Uniqueness	M	.4	1	0.624	<input type="checkbox"/>
L. Recreation/Education Potential (bonus points)	L	.05	NA	0.078	<input type="checkbox"/>
Totals:		5.85	10	9.126	
Percent of Possible Score			58.5 %		

**Category I Wetland:** (must satisfy **one** of the following criteria; otherwise go to Category II)

- ☐ Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**
- ☐ Score of 1 functional point for Uniqueness; **or**
- ☐ Score of 1 functional point for Flood Attenuation **and** answer to Question 14E.ii is "yes"; **or**
- ☐ Percent of possible score > 80% (round to nearest whole #).

**Category II Wetland:** (Criteria for Category I not satisfied **and** meets any **one** of the following criteria; otherwise go to Category IV)

- ☐ Score of 1 functional point for MT Natural Heritage Program Species Habitat; **or**
- ☐ Score of .9 or 1 functional point for General Wildlife Habitat; **or**
- ☐ Score of .9 or 1 functional point for General Fish Habitat; **or**
- ☐ "High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish/Aquatic Habitat; **or**
- ☐ Score of .9 functional point for Uniqueness; **or**
- ☐ Percent of possible score > 65% (round to nearest whole #).

**Category III Wetland:** (Criteria for Categories I, II, or IV not satisfied)



**Category IV Wetland:** (Criteria for Categories I or II are not satisfied and all of the following criteria are met; otherwise go to Category III)

- ☐ "Low" rating for Uniqueness; **and**
- ☐ Vegetated wetland component < 1 acre (do not include upland vegetated buffer); **and**
- ☐ Percent of possible score < 35% (round to nearest whole #).

**OVERALL ANALYSIS AREA RATING:**  
(check appropriate category based on the criteria outlined above)

I	II	III	IV
---	----	-----	----



## **Appendix C**

---

### Project Area Photographs

---

MDT Wetland Mitigation Monitoring  
Easton Ranch  
Park County, Montana



Veg Tran 1 – Start Location: Veg Com 8 foreground

Taken in 2010



Veg Tran 1 – Start Location: Veg Com 8 foreground

Taken in 2013



Veg Tran 1 – Start Location: Veg Com 8 foreground

Taken in 2014



Veg Tran 1 – Start Location: Veg Com 13 foreground

Taken in 2015





Veg Tran 1 – End Location: Veg Com 8 foreground

Taken in 2010



Veg Tran 1 – End Location: Veg Com 8 foreground

Taken in 2013



Veg Tran 1 – End Location: Veg Com 8 foreground

Taken in 2014



Veg Tran 1 – End Location: Veg Com 13 foreground

Taken in 2015





Veg Tran 2 – Start Location: Veg Com 1 foreground

Taken in 2010



Veg Tran 2 – Start Location: Veg Com 1 foreground

Taken in 2013



Veg Tran 2 – Start Location: Veg Com 1 foreground

Taken in 2014



Veg Tran 2 – Start Location: Veg Com 1 foreground

Taken in 2015





Veg Tran 2 – End Location: Veg Com 1 foreground

Taken in 2010



Veg Tran 2 – End Location: Veg Com 1 foreground

Taken in 2013



Veg Tran 2 – End Location: Veg Com 1 foreground

Taken in 2014



Veg Tran 2 – End Location: Veg Com 1 foreground

Taken in 2015





Veg Tran 3 – Start Location: Veg Com 1 foreground

Taken in 2010



Veg Tran 3 – Start Location: Veg Com 1 foreground

Taken in 2013



Veg Tran 3 – Start Location: Veg Com 1 foreground

Taken in 2014



Veg Tran 3 – Start Location: Veg Com 1 foreground

Taken in 2015





Photo Point 1 – Photo 1

Location: East Boundary

Date: 2010



Photo Point 1 – Photo 1

Location: East Boundary

Date: 2013



Photo Point 1 – Photo 1

Location: East Boundary

Date: 2014



Photo Point 1 – Photo 1

Location: East Boundary

Date: 2015



Photo Point 2 – Photo 1

Location: Northeast Corner of Site      Date: 2010



Photo Point 2 – Photo 1

Location: Northeast Corner of Site      Date: 2013



Photo Point 2 – Photo 1

Location: Northeast Corner of Site      Date: 2014



Photo Point 2 – Photo 1

Location: Northeast Corner of Site      Date: 2015





Photo Point 3 – Photo 1

Location: Northwest Corner of Site

Date: 2010



Photo Point 3 – Photo 1

Location: Northwest Corner of Site

Date: 2013



Photo Point 3 – Photo 1

Location: Northwest Corner of Site

Date: 2014



Photo Point 3 – Photo 1

Location: Northwest Corner of Site

Date: 2015





Photo Point 4a – Photo 1

Location: Shields Bank-DS

Date: 2010



Photo Point 4a – Photo 1

Location: Shields Bank-DS

Date: 2013



Photo Point 4a – Photo 1

Location: Shields Bank-DS

Date: 2014



Photo Point 4a – Photo 1

Location: Shields Bank-DS

Date: 2015





Photo Point 4a – Photo 1

Location: Shields Bank-US

Date: 2010



Photo Point 4a – Photo 1

Location: Shields Bank-US

Date: 2013



Photo Point 4a – Photo 1

Location: Shields Bank-US

Date: 2014



Photo Point 4a – Photo 1

Location: Shields Bank-US

Date: 2015





Photo Point 5 – Photo 1

Location: West Boundary

Date: 2010



Photo Point 5 – Photo 1

Location: West Boundary

Date: 2013



Photo Point 5 – Photo 1

Location: West Boundary

Date: 2014



Photo Point 5 – Photo 1

Location: West Boundary

Date: 2015



Photo Point 6 – Photo 1

Location: Southwest Corner of Site

Date: 2010



Photo Point 6 – Photo 1

Location: Southwest Corner of Site

Date: 2013



Photo Point 6 – Photo 1

Location: Southwest Corner of Site

Date: 2014



Photo Point 6 – Photo 1

Location: Southwest Corner of Site

Date: 2015





Photo Point 7 – Photo 1

Location: Southeast Corner of Site

Date: 2010



Photo Point 7 – Photo 1

Location: Southeast Corner of Site

Date: 2013



Photo Point 7 – Photo 1

Location: Southeast Corner of Site

Date: 2014



Photo Point 7 – Photo 1

Location: Southeast Corner of Site

Date: 2015



Wetland delineation sample point SP03. Facing NE toward the sample point.



Wetland delineation sample point SP03.



Bird box at T1 start.



## **Appendix D**

---

### Project Plan Sheets

---

MDT Wetland Mitigation Monitoring  
Easton Ranch  
Park County, Montana

