
**MONTANA DEPARTMENT OF TRANSPORTATION
WETLAND MITIGATION MONITORING REPORT: YEAR 2004**

*Wigeon Reservoir
Alzada, Montana*



Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Ave
Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING
~ A DIVISION OF **PBS&J**
P.O. Box 239
Helena, MT 59624

June 2005

Project No: B43054.00 - 0416



MONTANA DEPARTMENT OF TRANSPORTATION

WETLAND MITIGATION MONITORING REPORT:

YEAR 2004

*Wigeon Reservoir
Alzada, Montana*

Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Ave
Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING
A DIVISION OF PBS&J
P.O. Box 239
Helena, MT 59624

June 2005

Project No: B43054.00 - 0416



TABLE OF CONTENTS

| | |
|---|-----------|
| 1.0 INTRODUCTION..... | 1 |
| 2.0 METHODS..... | 1 |
| 2.1 Monitoring Dates and Activities..... | 1 |
| 2.2 Hydrology | 1 |
| 2.3 Vegetation..... | 3 |
| 2.4 Soils..... | 3 |
| 2.5 Wetland Delineation | 3 |
| 2.6 Mammals, Reptiles, and Amphibians | 3 |
| 2.7 Birds..... | 4 |
| 2.8 Macroinvertebrates | 4 |
| 2.9 Functional Assessment..... | 4 |
| 2.10 Photographs..... | 4 |
| 2.11 GPS Data..... | 4 |
| 2.12 Maintenance Needs..... | 4 |
| 3.0 RESULTS | 5 |
| 3.1 Hydrology | 5 |
| 3.2 Vegetation..... | 5 |
| 3.3 Soils..... | 7 |
| 3.4 Wetland Delineation | 8 |
| 3.5 Wildlife | 8 |
| 3.6 Macroinvertebrates | 9 |
| 3.7 Functional Assessment..... | 9 |
| 3.8 Photographs..... | 10 |
| 3.9 Maintenance Needs/Recommendations | 10 |
| 3.10 Current Credit Summary..... | 10 |
| 4.0 REFERENCES..... | 11 |

TABLES

| | |
|---------|---|
| Table 1 | <i>2001-2004 Wigeon Reservoir vegetation species list.</i> |
| Table 2 | <i>2001-2004 transect data summary.</i> |
| Table 3 | <i>Fish and wildlife species observed on the Wigeon Reservoir Mitigation Site from 2001-2004.</i> |
| Table 4 | <i>Summary of 2001-2004 wetlands function/value ratings and functional points at the Wigeon Reservoir Mitigation Project.</i> |

FIGURES

| | |
|----------|---|
| Figure 1 | <i>Project Site Location Map</i> |
| Figure 2 | <i>Monitoring Activity Locations 2004</i> |
| Figure 3 | <i>Mapped Site Features 2004</i> |

CHARTS

| | |
|---------|---|
| Chart 1 | <i>Length of vegetation communities along Transect 1.</i> |
| Chart 2 | <i>Transect maps showing vegetation types from the start of transect (0 feet) to the end of transect.</i> |
| Chart 3 | <i>Bioassessment scores for 2001-2004.</i> |

APPENDICES

| | |
|------------|--|
| Appendix A | <i>Figures 2 and 3</i> |
| Appendix B | <i>2004 Wetland Mitigation Site Monitoring Form</i> <i>2004 Bird Survey Forms</i> <i>2004 Wetland Delineation Forms</i> <i>2004 Functional Assessment Forms</i> |
| Appendix C | <i>Representative Photographs</i> <i>2004 Aerial Photograph</i> |
| Appendix D | <i>Bird Survey Protocol</i> <i>GPS Protocol</i> |
| Appendix E | <i>2004 Macroinvertebrate Sampling Protocol and Data</i> |

Cover Photo: Southwest edge of reservoir; view is east.

1.0 INTRODUCTION

The Wigeon wetland was created to provide mitigation credits for wetland impacts associated with Montana Department of Transportation (MDT) roadway projects that have been constructed in Watershed #16 of District Four (Glendive District). The site is located in Carter County, Montana, approximately 22 miles directly north of Alzada (**Figure 1**) in Sections 23 and 26, Township 5 South, Range 59 East. Elevations range from approximately 3,169 to 3,175 feet above sea level.

Construction was completed on this site in October of 1997 with the goal of creating a reservoir to provide nesting and brood rearing habitat for waterfowl and other wildlife species. An impoundment was constructed to collect surface water runoff from an intermittent tributary of Prairie Dog Creek. The site boundary is illustrated on **Figure 2, Appendix A**.

This wetland was designed by the BLM in association with the MDT to provide specific wetland functions including: nesting and brood rearing habitat for waterfowl; water for wildlife habitat; increased habitat diversity; water storage and retention; and creating open water and emergent wetland types.

2.0 METHODS

2.1 Monitoring Dates and Activities






The site was visited once on July 26, 2004. All information within the Wetland Mitigation Site Monitoring Form (**Appendix B**) and macroinvertebrate samples were collected at this time. Activities and information conducted/collected included: wetland delineation; wetland/open water boundary mapping; vegetation community mapping; vegetation transects; soils data; hydrology data; bird and general wildlife use; photograph points; macroinvertebrate sampling; functional assessment; and assessment of the maintenance needs at inflow area and outflow structure.

2.2 Hydrology

Wetland hydrology indicators were recorded using procedures outlined in the US Army Corps (COE) 1987 Wetland Delineation Manual (Environmental Laboratory 1987). Hydrology data were recorded on the Routine Wetland Delineation Data Form (**Appendix B**) at each wetland determination point.

All additional hydrologic data were recorded on the mitigation site monitoring form (**Appendix B**). Where possible, the boundary between wetlands and open water (no rooted vegetation) aquatic habitats was mapped on the aerial photograph and an estimate of the average water depth at this boundary was recorded (**Figure 3, Appendix A**). There are no groundwater monitoring wells at the site.

FIGURE 1. PROJECT LOCATION
Wigeon Reservoir
Mitigation Site



1: 24,000

PROJECT #: 130091.028
DATE: APRIL 2001
LOCATION:
PROJECT MANAGER: B. DUTTON
DRAWN BY: B. NOECKER



1120 CEDAR PO BOX 8254 MISSOULA, MT 59807

2.3 Vegetation

General vegetation types were delineated on an aerial photograph during the July site visit (**Figure 3, Appendix A**). Coverage of the dominant species in each community type is listed on the monitoring form (**Appendix B**). A comprehensive plant species list for the entire site was compiled in 2001 and has been updated with the new species encountered during the current season. Observations from past years will be compared with new data to document vegetation changes over time. Wigeon Reservoir is not fenced, and cattle have unrestricted access to the site. Woody species were not planted on this site.

One transect was established during the 2001 monitoring event to represent the range of vegetation conditions over time, especially the establishment and increase of hydrophytic vegetation. The transect was again sampled in 2004. The location of this transect is shown on **Figure 2, Appendix A**. Percent cover for each species was recorded on the vegetation transect form (**Appendix B**). Transect ends were marked with metal fence posts and their locations were recorded with the GPS unit during 2001. Photographs of the transect were taken from both ends during the 2004 site visit.

2.4 Soils

Soils were evaluated during the site visit according to the procedure outlined in the COE 1987 Wetland Delineation Manual. Soil data were recorded for each wetland determination point on the COE Routine Wetland Delineation Data Form (**Appendix B**).

2.5 Wetland Delineation

A wetland delineation was conducted within the area immediately adjacent to and including the reservoir according to the 1987 COE Wetland Delineation Manual. Wetland and upland areas within the monitoring area were investigated for the presence of wetland hydrology, hydrophytic vegetation and hydric soils. The indicator status of vegetation was derived from the National List of Plant Species that Occur in Wetlands: North Plains Region 4 (Reed 1988). The information was recorded on the COE Routine Wetland Delineation Forms (**Appendix B**). The wetland/upland and open water boundaries were used to calculate the wetland area developed at the reservoir.

2.6 Mammals, Reptiles, and Amphibians

Mammal, reptile, and amphibian species observations were recorded on the wetland monitoring form during the site visit (**Appendix B**). Indirect use indicators were also recorded including tracks, scat and burrows. A comprehensive wildlife species list for the entire site was compiled and will be updated as new species are encountered. Observations from past years will be compared with new data to determine if wildlife use is changing over time.

2.7 Birds

Bird observations were recorded during the site visit according to the established bird survey protocol (**Appendix D**). A general, qualitative bird list has been compiled using these observations. Observations will be compared between years in future studies.

2.8 Macroinvertebrates

One macroinvertebrate sample was collected during the 2004 site visit following the sampling protocol (**Appendix E**). Samples were preserved as outlined in the sampling procedure and sent to Rhithron Associates for analysis. The approximate sampling location is indicated on **Figure 2, Appendix A**. Results are included in **Appendix F**.

2.9 Functional Assessment

A functional assessment form was completed in 2004 for the Wigeon reservoir using the 1999 MDT Montana Wetland Assessment Method. Field data necessary for this assessment were collected on a condensed data sheet. The remainder of the assessment was completed in the office (**Appendix B**).

2.10 Photographs

Photographs were taken showing the current land use surrounding the site, the wetland buffer, the monitoring area, and the vegetation transect. A description and compass direction for each photograph were recorded on the wetland monitoring form. The approximate locations of the photos are shown on **Figure 2, Appendix A**. All photographs were taken using a digital camera; representative photos are included in **Appendix C**.

2.11 GPS Data

During the 2001 monitoring season, survey points were collected using a resource grade Trimble, Geoexplorer III hand-held GPS unit (**Appendix D**). Points collected included: the vegetation transect beginning and ending locations; photograph locations; and the jurisdictional wetland boundary. The wetland boundary was updated in 2004 by adjusting the boundary by hand-drawing on an aerial photograph.

2.12 Maintenance Needs

There are no inflow or outflow structures or nest boxes at this site. The only hydrologic control structure at the Wigeon wetland is the dike; no pipes or other outflow structures were installed to convey water through the dike or out of the reservoir. The dike structure was examined for obvious maintenance needs during the July visit.

3.0 RESULTS

3.1 Hydrology

Open water / aquatic bed represented 49% of the area within the wetland boundary. The reservoir was not at full pool as a result of the drought. Negligible emergent vegetation was observed within the open water or within community type 3 at the time of investigation because of the low water level. Water depths were estimated to range between 1 and 4 feet deep throughout the reservoir. The open water boundary is depicted on **Figure 3, Appendix A**. The primary source of hydrology is an intermittent tributary of Prairie Dog Creek and the secondary source is likely groundwater. No problems with the dike were noted.

According to the Western Regional Climate Center (WRCC 2005), the Ridgeway 1S station annual mean (1952 – 2004) precipitation was 13.27 inches; the average precipitation through the month of July for this period was 8.02 inches. For 2004, precipitation through July was 4.9 inches or 61% of the mean. Since the year 2000, with the exception of 2003, yearly precipitation has been below average.

3.2 Vegetation

Vegetation species identified on the site are presented in **Table 1** and in the monitoring form (**Appendix B**). **Table 2** and **Charts 1** and **2** illustrate transect data trends. Three (3) major vegetation communities were mapped on the mitigation area map (**Figure 3, Appendix A**). The communities include: Type 1, *Artemisia tridentata*/*Bouteloua gracilis*; Type 2, *Hordeum jubatum*; and Type 3, *Chenopodium glauca*/*Eleocharis* spp. (in 2003, dominant species in this community type changed from *Typha latifolia* / *Eleocharis palustris*). Dominant species within each community are listed on the monitoring form (**Appendix B**). Community Type 3 was primarily represented by *Chenopodium* and exposed substrate (mud) during the 2004 season as a result of very low water levels. No wetland vegetation of any growth stage (i.e. single vegetative blades) was observed in community type 3 as was the case in 2003. No new species were observed in 2004.

The transect length was increased in 2003 to include the exposed substrate that has resulted from the lower water level caused by drought. These transect data (**Table 2** and **Chart 1**) indicate that the site has completely lost the preferred hydrophytic vegetation community, which included *Scirpus*, *Carex*, *Juncus* and *Eleocharis* species, and has subsequently been replaced by *Chenopodium*, a non-preferred FACW weedy species.

Table 1: 2001-2004 Wigeon Reservoir vegetation species list.

| Scientific Name ¹ | Region 4 (North Plains) Wetland Indicator Status ² |
|---|---|
| <i>Achillea millefolium</i> | FACU |
| <i>Agropyron cristatum</i> | -(UPL) |
| <i>Agropyron dasystachyum</i> | FAC |
| <i>Agropyron smithii</i> | FACU |
| <i>Artemesia cana</i> | FACU |
| <i>Artemesia tridentate</i> | (UPL) |
| <i>Bouteloua gracilis</i> | (UPL) |
| <i>Bromus japonicus</i> | (UPL) |
| <i>Carex spp.</i> | (unknown, FAC-OBL) |
| <i>Carex utriculata</i> | OBL |
| <i>Chenopodium glaucum</i> | FACW |
| <i>Eleocharis acicularis</i> | OBL |
| <i>Eleocharis palustris</i> | OBL |
| <i>Festuca idahoensis</i> | (UPL) |
| <i>Grindelia gracilifolia</i> | FACW |
| <i>Hordeum jubatum</i> | FACW |
| <i>Juncus spp.</i> | (unknown, FAC-OBL) |
| <i>Myosotis scorpioides</i> | OBL |
| <i>Najas flexilis</i> | OBL |
| <i>Opuntia spp.</i> | (UPL) |
| <i>Phleum pretense</i> | FACU |
| <i>Puccinelliana nuttalliana</i> | OBL |
| <i>Sagittaria spp.</i> | OBL |
| <i>Scirpus spp.</i> | OBL |
| <i>Thlaspi arvense</i> | (FACU) |
| <i>Typha latifolia</i> | OBL |
| <i>Xanthium strumarium</i> | FAC |

¹ **Bolded** species indicate those documented within the analysis area for the first time in 2004.

² Species either not included or classified as “non-indicator” for the National List of Plant Species that Occur in Wetlands: North Plains (Region 4); status in parentheses are probable and based on biologist’s experience.

Table 2: 2001-2004 transect data summary.

| Monitoring Year | 2001 | 2002 | 2003 | 2004 |
|--|------|------|-----------------|------|
| Transect Length (feet) | 39 | 39 | 54 ¹ | 54 |
| # Vegetation Community Transitions along Transect | 2 | 2 | 2 | 2 |
| # Vegetation Communities along Transect | 3 | 1 | 3 | 3 |
| # Hydrophytic Vegetation Communities along Transect | 1 | 1 | 2 | 2 |
| Total Vegetative Species | 11 | 7 | 6 | 4 |
| Total Hydrophytic Species | 4 | 3 | 4 | 2 |
| Total Upland Species | 7 | 4 | 2 | 2 |
| Estimated % Total Vegetative Cover | 76 | 78 | 88 | 91 |
| % Transect Length Comprised of Hydrophytic Vegetation Communities | 21 | 38 | 56 | 56 |
| % Transect Length Comprised of Upland Vegetation Communities | 79 | 61 | 44 | 44 |
| % Transect Length Comprised of Unvegetated Open Water | 0 | 0 | 0 | 0 |
| % Transect Length Comprised of Bare Substrate | 0 | 0 | 0 | 0 |

¹ Lengthened in 2003 as a result of exposed substrate area.

Chart 1: Length of vegetation communities along Transect 1.





Chart 2: Transect maps showing vegetation types from the start of transect (0 feet) to the end of transect. Transect 1 was lengthened from 39 feet (2001-2002) to 54 feet (2003-2004) as a result of exposed substrate area. Vegetation species within community types are not static across years.



3.3 Soils

The site was mapped as part of the Carter County Soil Survey. The dominant soil on the site is the Moyerson-Orinoco complex (277D) a silty clay loam, and the Gerdrum-Absher (165C) complex (Typic Natriboralfs). The taxonomic classification of the 277D series components are Ustic Torriorthent and Ardic Ustorthent, respectively.

The Myerson-Orinoco (277D) is typical of sedimentary plains and hills and the Gerdrum-Absher complex (165C) occurs in alluvial fans and stream terraces. Neither of these soil series are hydric or have hydric inclusions. Both soils types are poor for wetland plant establishment and have a high saline content.

Soil pit (SP) 1 was excavated within the wetland vegetation community. At 10 inches the soil was black (10YR 2/1) clay and saturated at a depth of 6 inches. SP-2 was excavated in the upland community; at 10 inches the soil was a very dark gray (2.5Y 4/1) with yellowish brown (10 YR 5/8) mottles.

3.4 Wetland Delineation

The delineated wetland boundary is depicted on **Figure 3, Appendix A**. The COE data forms are included in **Appendix B**.

The 5.5 acres of gross wetland area encompasses 2.81 acres of primarily goosefoot (*Chenopodium glaucum*, FAC), foxtail (*Hordeum jubatum*), and mud. No other wetland species were observed. The first year of monitoring, 2001, the gross wetland area totaled 8.2 acres and included 3.0 acres of emergent wetland. The drought has caused a 33% decline in the gross wetland area.

3.5 Wildlife

Wildlife species are listed in **Table 3**. Activities and densities associated with these observations are included on the monitoring form in **Appendix B**. Leopard frogs, a “species of special concern” (S3) by the Montana Natural Heritage Program (MNHP), were first observed in 2001.

Table 3. Fish and wildlife species observed on the Wigeon Reservoir Mitigation Site from 2001-2004.

| | |
|---|--|
| AMPHIBIANS AND REPTILES | |
| Plains garter snake (<i>Thamnophis radix</i>) | |
| Painted turtle (<i>Chrysemys picta</i>) | |
| Leopard frogs (<i>Rana pipiens</i>) | |
| BIRDS | |
| American Avocet (<i>Recurvirostra americana</i>) ¹ | Northern Pintail (<i>Anas acuta</i>) ¹ |
| American Wigeon (<i>Anas americanus</i>) | Red-winged Blackbird (<i>Agelaius phoeniceus</i>) |
| Blue-winged teal (<i>Anas discors</i>) | Redhead (<i>Aythya Americana</i>) |
| Bobolink (<i>Dolichonyx oryzivorus</i>) | Ring-necked Duck (<i>Aythya collaris</i>) ¹ |
| Canada Goose (<i>Branta canadensis</i>) ¹ | Ruddy Duck (<i>Oxyura jamaicensis</i>) |
| Eared grebes (<i>Podiceps nigricollis</i>) | Greater Sage Grouse (<i>Centrocercus urophasianus</i>) |
| Horned Lark (<i>Eremophila alpestris</i>) | Savanannah Sparrow (<i>Passerculus sandwichensis</i>) |
| Gadwall (<i>Anas strepera</i>) | Spotted sandpiper (<i>Actitis macularia</i>) |
| Grasshopper Sparrow (<i>Ammodramus savannarum</i>) | Swallow (<i>Hirundo</i> spp.) |
| Green-winged Teal (<i>Anas crecca</i>) ¹ | Upland Sandpiper (<i>Bartramia longicauda</i>) |
| Killdeer (<i>Charadrius vociferous</i>) | Wilson’s Phalarope (<i>Phalaropus tricolor</i>) ¹ |
| Mallards (<i>Anas platyrhynchos</i>) | Willet (<i>Catoptrophorus semipalmatus</i>) ¹ |
| Meadow Lark (<i>Sturnella neglecta</i>) | |

¹ June 2003 MDT sightings.

Table 3. Fish and Wildlife Species Observed on the Wigeon Reservoir Mitigation Site from 2001-2004.


| |
|----------------------------------|
| MAMMALS |
| Deer (<i>Odocoileus</i> spp.) |
| Raccoon (<i>Procyon lotor</i>) |

[†] June 2003 MDT sightings.

3.6 Macroinvertebrates

The optimal conditions noted in 2001 have yet to be restored at this site on Wigeon Reservoir, but scores indicate a stabilization of conditions since 2002 (**Bollman, Appendix E**). In that year, biotic condition was rated poor while scores for both 2003 and 2004 indicate sub-optimal conditions (**Chart 3**). Taxa richness increased significantly, buoyed by the addition of several chironomid taxa which were not reported from the site in 2003. The biotic index value was near the median value for all sites over all years of study. Habitat complexity may have improved between the 2 years with improved availability of substrate habitats apparent.

Chart 3: Bioassessment scores for 2001-2004.



3.7 Functional Assessment

Completed functional assessment forms are included in **Appendix B** and summarized below in **Table 4**. Functional units have decreased 28% since 2001 because of a 2.7-acre decrease in gross wetland area caused by drought. The wetland continues to rank as a Category II wetland as it provides primary habitat for an MNHP species of special concern, the leopard frog. The diversity of wildlife that use the reservoir is high as evidenced by the diversity of waterfowl, amphibians and reptiles. Disturbance by cattle or observations of heavily cattle-tracked areas appeared to be less in 2004 at the time of the investigation. Thus, the disturbance rating was revised from high to moderate, which increased some of the values resulting in an increase in actual functional points.

3.8 Photographs

Representative photos taken from photo points and transect ends are included in **Appendix C**. A 2004 aerial photograph is provided in **Appendix C**.

3.9 Maintenance Needs/Recommendations

No observable problems were noted concerning the dike structure. Fencing the outer limits of the wetland boundary prior to the end of the drought is recommended to protect reestablishing hydrophytic wetland vegetation. Several watering-access points can be incorporated into the fence perimeter to allow cattle access.

Table 4: Summary of 2001-2004 wetland function/value ratings and functional points at the Wigeon Reservoir mitigation project.

| Function and Value Parameters From the 1999 MDT Montana Wetland Assessment Method | 2001 | 2002 | 2003 | 2004 |
|---|--------------|--------------|--------------|--------------|
| Listed/Proposed T&E Species Habitat | Low (0) | Low (0) | Low (0) | Low (0) |
| MNHP Species Habitat | High (1) | High (1) | High (1) | High (1) |
| General Wildlife Habitat | Mod (.5) | Mod (.7) | Mod (.7) | High (.9) |
| General Fish/Aquatic Habitat | Mod (.6) | Mod (.6) | Mod (.6) | Mod (.5) |
| Flood Attenuation | Mod (.5) | Mod (.5) | Mod (.5) | Mod (.5) |
| Short and Long Term Surface Water Storage | High (1) | High (1) | High (1) | High (1) |
| Sediment, Nutrient, Toxicant Removal | Mod (.7) | Mod (.7) | Mod (.7) | Mod (.7) |
| Sediment/Shoreline Stabilization | Mod (.7) | Mod (.7) | Low (.3) | Low (.3) |
| Production Export/Food Chain Support | Mod (.6) | Mod (.6) | Mod (.6) | High (.8) |
| Groundwater Discharge/Recharge | High (1) | High (1) | High (1) | High (1) |
| Uniqueness | Low (.3) | Low (.2) | Low (.2) | Low (.3) |
| Recreation/Education Potential | Low (.1) | Low (.2) | Low (.2) | Low (.5) |
| Actual Points/Possible Points | 7/12 | 7.2/12 | 6.8/12 | 7.5/12 |
| % of Possible Score Achieved | 58% | 60% | 56% | 63% |
| Overall Category | II | II | II | II |
| Total Acreage of Assessed Wetlands within Easement | 8.20 | 8.09 | 8.09 | 5.50 |
| Functional Units (acreage x actual points) | 57.40 | 58.24 | 55.00 | 41.25 |
| Net Acreage Gain | 8.20 | 8.09 | 8.09 | 5.50 |
| Net Functional Unit Gain | 57.40 | 58.24 | 55.00 | 41.25 |

3.10 Current Credit Summary

The 5.5 acres of gross wetland area encompasses 2.81 acres of goosefoot, foxtail, and mud. No other wetland species were observed. In 2001, the gross wetland area totaled 8.2 acres and included 3.0 acres of emergent wetland. The drought has caused a 33% decline in the gross wetland area and nearly 100% loss of desirable wetland vegetation species. It is likely, however, that wetland area and species will be regained with normal precipitation. Functional units have decreased 28% since 2001 as a result of a 2.7 acre decrease in gross wetland area caused by the drought. However, the wetland continues to rank as a Category II wetland.

Fencing the outer limits of the wetland boundary prior to the end of the drought is recommended to protect reestablishing hydrophytic wetland vegetation. Several watering-access points can be incorporated into the fence perimeter to allow cattle access.

4.0 REFERENCES


- Berglund, J. 1999. *MDT Montana Wetland Assessment Method*. May. Montana Department of Transportation, Helena, Montana.
- Bollman, W. 2004. MDT Wetland Mitigation Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001, 2002, 2003, and 2004. Rhithron Associates Inc. Missoula, Montana.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. U.S. Army Corps of Engineers. Washington, DC.
- Montana Department of Transportation (MDT). 1996. *MDT Biological Resources Report: Alzada South*. Montana Department of Transportation, Helena, Montana.
- Reed, P. 1988. *National list of plant species that occur in wetlands: North Plains (Region 4)*. Biological Report 88(26.4), May. U.S. Fish and Wildlife Service, Washington, D.C.
- U.S. Army Corps of Engineers (COE). 1987. *Corps of Engineers Wetlands Delineation Manual*. U.S. Army Corps. Washington, DC.
- USDA Natural Resource Conservation Service (NRCS). 2003. *Soil Survey of Carter County Area, Montana*.
- Western Regional Climate Center (WRCC). 2005. Ridgeway 1S Station:
<http://www.wrcc.dri.edu/cgi-bin/cliMAIN.pl?mtridg>.

Appendix A

FIGURES 2 - 3

*MDT Wetland Mitigation Monitoring
Wigeon Reservoir
Alzada, Montana*

Figure 2 Monitoring Activity Locations 2004




Legend

- Monitoring Area Limits
- Vegetation Transect
- Photograph Point
- Aerial Reference Point
- Soil Sample Point
- Macro-invertebrate Sample
- Base Photograph July 25, 2004

| | | |
|---|--|--|
| PROJ NO: 330054.416 FILE NAME: BASE2004.dwg SCALE: 1"= 150 ft LOCATION: Wigeon Reservoir | DRAWN: RA CHECKED: APPVD: BD PROJ MGR: BD | PROJECT NAME MDT Wigeon Reservoir Wetland Mitigation DRAWING TITLE Monitoring Activity Locations 2004 |
| SHEET NUMBER 2 OF | | REV 02 DATE: 06/23/05 |



Figure 3 Mapped Site Features 2004



Legend

- Monitoring Area Limits
- Wetland Boundary
- Vegetation Community Boundary
- Open Water Boundary
- Base Photograph July 25, 2004

- Vegetation CommunityType:**
- ① Artemisia tridentata/Bouteloua gracilis
 - ② Hordeum jubatum
 - ③ Chenopodium glaucum/Eleocharis spp.

Wetland Area 2004

| | |
|------------|------------|
| Gross Area | 5.50 Acres |
| Open Water | 2.69 Acres |
| Net Area | 2.81 Acres |



Appendix B

2004 WETLAND MITIGATION SITE MONITORING FORM

2004 BIRD SURVEY FORMS

2004 WETLAND DELINEATION FORMS

2004 FUNCTIONAL ASSESSMENT FORMS

MDT Wetland Mitigation Monitoring

Wigeon Reservoir

Alzada, Montana

VEGETATION COMMUNITIES

Community No.: 1 Community Title (main species): Artemesia spp./Bouteloua gracilis

| Dominant Species | % Cover | Dominant Species | % Cover |
|-----------------------------|---------|-------------------------------|---------|
| <i>Artemesia cana</i> | 10 | <i>Agropyron cristatum</i> | 10 |
| <i>Opuntia spp.</i> | 10 | <i>Grindelia gracifolia</i> | 10 |
| <i>Achillea millefolium</i> | 10 | <i>Agropyron dasystachyum</i> | 5 |
| <i>Bouteloua gracilis</i> | 25 | <i>Artemesia tridentata</i> | 5 |
| <i>Festuca idahoensis</i> | 15 | <i>Melolotis officinale</i> | (*) |

COMMENTS/PROBLEMS: yellow clover has increased in some areas

Community No.: 2 Community Title (main species): Hordeum jubatum

| Dominant Species | % Cover | Dominant Species | % Cover |
|-----------------------------|---------|----------------------------------|---------|
| <i>Hordeum jubatum</i> | 60 | <i>Melilotis officinalis</i> | 2 |
| <i>Phleum pratense</i> | 1 | <i>Puccinelliana nuttalliana</i> | <1 |
| <i>Grindelia gracifolia</i> | 1 | | |
| <i>Xanthium strumarium</i> | 1 | | |
| <i>Chenopodium glaucum</i> | 40 | | |

COMMENTS/PROBLEMS: _____

Community No.: 3 Community Title (main species): Chenopodium glaucum/Eleocharis spp.

| Dominant Species | % Cover | Dominant Species | % Cover |
|------------------------------|---------|------------------------------------|---------|
| <i>Typha latifolia</i> | 0 | <i>Chenopodium glaucum</i> | 45 |
| <i>Eleocharis palustris</i> | 0 | <i>Puccinellia nuttalliana</i> | 0 |
| <i>Scirpus spp.</i> | 0 | exposed pond substrate (dried mud) | 55 |
| <i>Sagittaria spp.</i> | 0 | <i>Hordeum jubatum</i> | <1 |
| <i>Eleocharis acicularis</i> | 0 | | |

COMMENTS/PROBLEMS: **2004 CT 3 composition has changed to primarily mud and *Chenopodium*.**
Will continue to identify it as CT 3.

Additional Activities Checklist:

Record and map vegetative communities on air photo

PHOTOGRAPHS

Using a camera with a 50 mm lenses and color film take photographs of the following permanent reference points listed in the checklist below. Record the direction of the photograph using a compass. (The first time at each site establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3' above ground, survey the location with a resource grade GPS and mark the location on the air photo.)

Checklist:

- One photo for each of the 4 cardinal directions surrounding wetland
- At least one photo showing upland use surrounding wetland – if more than one upland use exists, take additional photos
- At least one photo showing buffer surrounding wetland
- One photo from each end of vegetation transect showing transect

| Location | Photograph Description | Compass Reading |
|----------|--|-----------------|
| A | wetland view | 194 |
| B | Edge of open water between photo points D and E. | 180 |
| C | wetland buffer | 280 |
| D | wetland view | 90 |
| E | wetland view | 0 |
| F | wetland view | 330 |
| G | wetland transect end | 10 |
| H | UPL transect beginning | 190 |

COMMENTS/PROBLEMS: _____

GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points with the GPS unit set at 5 second recording rate. Record file numbers fore site in designated GPS field notebook

Checklist:

- Jurisdictional wetland boundary
- 4-6 landmarks recognizable on the air photo
- Start and end points of vegetation transect(s)
- Photo reference points
- Groundwater monitoring well locations

COMMENTS/PROBLEMS: _____ wetland boundary hand-drawn

WETLAND DELINEATION

(Attach Corps of Engineers delineation forms)

At each site conduct the items on the checklist below:

- Delineate wetlands according to the 1987 Army Corps manual.
- Delineate wetland-upland boundary on the air photo
- * Survey wetland-upland boundary with a resource grade GPS survey

COMMENTS/PROBLEMS: *hand-drawn 2004

FUNCTIONAL ASSESSMENT

(Complete and attach full MDT Montana Wetland Assessment Method field forms; also attach abbreviated field forms, if used)

COMMENTS/PROBLEMS: _____

MAINTENANCE

Were man-made nesting structures installed at this site? YES___ NO

If yes, do they need to be repaired? YES___ NO___

If yes, describe problems below and indicate if any actions were taken to remedy the problems.

Were man-made structures build or installed to impound water or control water flow into or out of the wetland?

YES NO___

If yes, are the structures working properly and in good working order? YES NO___

If no, describe the problems below.

COMMENTS/PROBLEMS: Water levels continue to decrease; likely the result of drought in this area. Cattle tracks noted, however no wetland vegetation remains to be grazed.

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: Wigeon Date: 7/26/04 Examiner: LB/LWC Transect # 1

Approx. transect length: 54 feet Compass Direction from Start (Upland): 10 deg

| Vegetation type A: | | CT 1 |
|----------------------------------|----|--------|
| Length of transect in this type: | 24 | feet |
| Species: | | Cover: |
| AGRSMI | | 10 |
| HORJUB | | 40 |
| MELOFF | | 40 |
| CHENGLA | | 10 |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| Total Vegetative Cover: | | 100% |

| Vegetation type B: | | CT 2 |
|----------------------------------|----|--------|
| Length of transect in this type: | 15 | feet |
| Species: | | Cover: |
| HORJUB | | <1 |
| CHEGLA. | | 99 |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| Total Vegetative Cover: | | 100% |

| Vegetation type C: | | CT 3 |
|----------------------------------|----|--------|
| Length of transect in this type: | 15 | feet |
| Species: | | Cover: |
| CHEGLA | | 55 |
| mud | | 35 |
| HORJUB | | 10 |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| Total Vegetative Cover: | | 65% |

| Vegetation type D: | | CT 4 |
|----------------------------------|--|--------|
| Length of transect in this type: | | feet |
| Species: | | Cover: |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| Total Vegetative Cover: | | |

SOILS

| Map Unit Name | | Moyerson-Orinoco (277D)-non-hydric | | Drainage Class: <u>mod. well</u> | |
|---|---------|---|-------------------------------|--|---------------------------------------|
| (Series and Phase): | | | | Field Observations | |
| Taxonomy (Subgroup): | | <u>NA</u> | | Confirm Mapped Type? <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No | |
| Profile Description: | | | | | |
| Depth inches | Horizon | Matrix Color (Munsell Moist) | Mottle Colors (Munsell Moist) | Mottle Abundance/Contrast | Texture, Concretions, Structure, etc. |
| 10" | A | 10 YR 2/1 | | | clay |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| Hydric Soil Indicators: | | | | | |
| <input type="checkbox"/> Histosol | | <input type="checkbox"/> Concretions | | | |
| <input type="checkbox"/> Histic Epipedon | | <input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils | | | |
| <input type="checkbox"/> Sulfidic Odor | | <input type="checkbox"/> Organic Streaking in Sandy Soils | | | |
| <input type="checkbox"/> Aquic Moisture Regime | | <input type="checkbox"/> Listed on Local Hydric Soils List | | | |
| <input type="checkbox"/> Reducing Conditions | | <input type="checkbox"/> Listed on National Hydric Soils List | | | |
| <input checked="" type="checkbox"/> Gleyed or Low-Chroma Colors | | <input type="checkbox"/> Other (Explain in Remarks) | | | |

WETLAND DETERMINATION

| | | | |
|---|---|-----------------------------|--|
| Hydrophytic Vegetation Present? | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No | Is this Sampling Point Within a Wetland? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No |
| Wetland Hydrology Present? | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No | |
| Hydric Soils Present? | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No | |
| Remarks: | | | |
| Entire CT 3 does qualify as WL as a result of >30% <i>Chenopodium glaucum</i> , however, half of CT 3- the portion adjacent to the water edge- is technically just mud and not vegetated. | | | |

Approved by HQUSACE 2/92

SOILS

| Map Unit Name | | Moyerson-Orinoco (277D)-non-hydric | | Drainage Class: | mod. well |
|---|---------|---|-------------------------------|---------------------------|---|
| (Series and Phase): | | | | Field Observations | |
| Taxonomy (Subgroup): | | NA | | Confirm Mapped Type? | Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> |
| Profile Description: | | | | | |
| Depth inches | Horizon | Matrix Color (Munsell Moist) | Mottle Colors (Munsell Moist) | Mottle Abundance/Contrast | Texture, Concretions, Structure, etc. |
| 10 | A | 2.5Y 4/1 | 10 YR 5/8 | | silt |
| | | | | | |
| | | | | | |
| | | | | | |
| Hydric Soil Indicators: | | | | | |
| <input type="checkbox"/> Histosol | | <input type="checkbox"/> Concretions | | | |
| <input type="checkbox"/> Histic Epipedon | | <input type="checkbox"/> High Organic Content in surface Layer in Sandy Soils | | | |
| <input type="checkbox"/> Sulfidic Odor | | <input type="checkbox"/> Organic Streaking in Sandy Soils | | | |
| <input type="checkbox"/> Aquic Moisture Regime | | <input type="checkbox"/> Listed on Local Hydric Soils List | | | |
| <input type="checkbox"/> Reducing Conditions | | <input type="checkbox"/> Listed on National Hydric Soils List | | | |
| <input type="checkbox"/> Gleyed or Low-Chroma Colors | | <input type="checkbox"/> Other (Explain in Remarks) | | | |
| Area seems to have a history of inundation based on soil colors; constructed in 1997; 4 years prior to first monitoring event by LWC. | | | | | |

WETLAND DETERMINATION

| | | | |
|---|---|--|--|
| Hydrophytic Vegetation Present? | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No | Is this Sampling Point Within a Wetland? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No |
| Wetland Hydrology Present? | <input type="checkbox"/> Yes | <input checked="" type="checkbox"/> No | |
| Hydric Soils Present? | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No | |
| Remarks: | | | |
| This area may convert to WL if water level increases. | | | |

Approved by HQUSACE 2/92

14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS

i. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S
- Secondary habitat (list species) D S
- Incidental habitat (list species) D S
- No usable habitat D S

ii. **Rating** (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

| | | | | | | | |
|-----------------------------|-------------|-------------|---------------|---------------|----------------|----------------|------|
| Highest Habitat Level | doc/primary | sus/primary | doc/secondary | sus/secondary | doc/incidental | sus/incidental | none |
| Functional Point and Rating | --- | --- | --- | --- | --- | --- | --- |

If documented, list the source (e.g., observations, records, etc.): _____

14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.

Do not include species listed in 14A(i).

i. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S Rana pipiens
- Secondary habitat (list species) D S _____
- Incidental habitat (list species) D S _____
- No usable habitat D S _____

iii. **Rating** (Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

| | | | | | | | |
|-----------------------------|-------------|-------------|---------------|---------------|----------------|----------------|------|
| Highest Habitat Level: | doc/primary | sus/primary | doc/secondary | sus/secondary | doc/incidental | sus/incidental | none |
| Functional Point and Rating | 1 (H) | --- | --- | --- | --- | --- | --- |

If documented, list the source (e.g., observations, records, etc.): _____

14C. General Wildlife Habitat Rating

i. **Evidence of overall wildlife use in the AA:** (Check either substantial, moderate, or low)

Substantial (based on any of the following)

- observations of abundant wildlife #s or high species diversity (during any period)
- abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- presence of extremely limiting habitat features not available in the surrounding area
- interviews with local biologists with knowledge of the AA

Low (based on any of the following)

- few or no wildlife observations during peak use periods
- little to no wildlife sign
- sparse adjacent upland food sources
- interviews with local biologists with knowledge of AA

Moderate (based on any of the following)

- observations of scattered wildlife groups or individuals or relatively few species during peak periods
- common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- adequate adjacent upland food sources
- interviews with local biologists with knowledge of the AA

ii. **Wildlife Habitat Features** (Working from top to bottom, select appropriate AA attributes to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from #13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see #10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A = absent.

| Structural Diversity (from #13) | <input type="checkbox"/> High | | | | | | | | <input checked="" type="checkbox"/> Moderate | | | | | | | | <input type="checkbox"/> Low | | | |
|--|-------------------------------|-----|-----|----|---------------------------------|-----|-----|----|--|-----|-----|----|---------------------------------|-----|-----|----|-------------------------------|-----|-----|----|
| | <input type="checkbox"/> Even | | | | <input type="checkbox"/> Uneven | | | | <input checked="" type="checkbox"/> Even | | | | <input type="checkbox"/> Uneven | | | | <input type="checkbox"/> Even | | | |
| Class Cover Distribution (all vegetated classes) | | | | | | | | | | | | | | | | | | | | |
| Duration of Surface Water in ≥ 10% of AA | P/P | S/I | T/E | A | P/P | S/I | T/E | A | P/P | S/I | T/E | A | P/P | S/I | T/E | A | P/P | S/I | T/E | A |
| Low disturbance at AA (see #12) | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Moderate disturbance at AA (see #12) | -- | -- | -- | -- | -- | -- | -- | -- | H | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| High disturbance at AA (see #12) | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |

iii. **Rating** (Using 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.)

| Evidence of Wildlife Use from 14C(i) | Wildlife Habitat Features Rating from 14C(ii) | | | |
|--------------------------------------|--|--|-----------------------------------|------------------------------|
| | <input type="checkbox"/> Exceptional | <input checked="" type="checkbox"/> High | <input type="checkbox"/> Moderate | <input type="checkbox"/> Low |
| Substantial | -- | .9 (H) | -- | -- |
| Moderate | -- | -- | -- | -- |
| Low | -- | -- | -- | -- |

Comments: Avian species composition likely high but are not observed. As we spend more time there it is apparent recording diversity is limited by observation periods.

14D. GENERAL FISH/AQUATIC HABITAT RATING NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat, excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

i. **Habitat Quality** (Pick the appropriate AA attributes in matrix to pick the exceptional (E), high (H), moderate (M), or low (L) quality rating.)

| Duration of Surface Water in AA | <input checked="" type="checkbox"/> Permanent/Perennial | | | <input type="checkbox"/> Seasonal / Intermittent | | | <input type="checkbox"/> Temporary / Ephemeral | | |
|--|---|--------|------|--|--------|------|--|--------|------|
| Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation) | >25% | 10-25% | <10% | >25% | 10-25% | <10% | >25% | 10-25% | <10% |
| Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities. | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities. | -- | -- | M | -- | -- | -- | -- | -- | -- |

ii. **Modified Habitat Quality:** Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support?

Y N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating: E H M L

iii. **Rating** (Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to pick the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).)

| Types of Fish Known or Suspected Within AA | Modified Habitat Quality from 14D(ii) | | | |
|--|---------------------------------------|-------------------------------|--|------------------------------|
| | <input type="checkbox"/> Exceptional | <input type="checkbox"/> High | <input checked="" type="checkbox"/> Moderate | <input type="checkbox"/> Low |
| Native game fish | -- | -- | -- | -- |
| Introduced game fish | -- | -- | -- | -- |
| Non-game fish | -- | -- | .5 (M) | -- |
| No fish | -- | -- | -- | -- |

Comments: Fish fry observed by MDT June 2003.

14E. FLOOD ATTENUATION NA (proceed to 14G)

Applies only to wetlands subject to flooding via in-channel or overbank flow.

If wetlands in AA do not flooded from in-channel or overbank flow, check NA above.

i. **Rating** (Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

| Estimated wetland area in AA subject to periodic flooding | <input type="checkbox"/> ≥ 10 acres | | | <input checked="" type="checkbox"/> <10, >2 acres | | | <input type="checkbox"/> ≤2 acres | | |
|---|-------------------------------------|--------|------|---|--------|--------|-----------------------------------|--------|------|
| % of flooded wetland classified as forested, scrub/shrub, or both | 75% | 25-75% | <25% | 75% | 25-75% | <25% | 75% | 25-75% | <25% |
| AA contains no outlet or restricted outlet | -- | -- | -- | -- | -- | .5 (M) | -- | -- | -- |
| AA contains unrestricted outlet | -- | -- | -- | -- | -- | -- | -- | -- | -- |

ii. **Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA?** (check) Y N Comments: _____

14F. SHORT AND LONG TERM SURFACE WATER STORAGE NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow.

If no wetlands in the AA are subject to flooding or ponding, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Abbreviations: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

| Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding. | <input checked="" type="checkbox"/> >5 acre feet | | | <input type="checkbox"/> <5, >1 acre feet | | | <input type="checkbox"/> ≤1 acre foot | | |
|--|--|-----|-----|---|-----|-----|---------------------------------------|-----|-----|
| Duration of surface water at wetlands within the AA | P/P | S/I | T/E | P/P | S/I | T/E | P/P | S/I | T/E |
| Wetlands in AA flood or pond ≥ 5 out of 10 years | 1 (H) | -- | -- | -- | -- | -- | -- | -- | -- |
| Wetlands in AA flood or pond < 5 out of 10 years | -- | -- | -- | -- | -- | -- | -- | -- | -- |

Comments: _____

14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL NA (proceed to 14H)

Applies to wetlands with potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input.

If no wetlands in the AA are subject to such input, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

| Sediment, Nutrient, and Toxicant Input Levels Within AA | AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present. | | | | Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present. | | | |
|---|---|-----------------------------|---|-----------------------------|--|-----------------------------|--------------------------------|-----------------------------|
| | <input type="checkbox"/> ≥ 70% | | <input checked="" type="checkbox"/> < 70% | | <input type="checkbox"/> ≥ 70% | | <input type="checkbox"/> < 70% | |
| Evidence of flooding or ponding in AA | <input type="checkbox"/> Yes | <input type="checkbox"/> No | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No | <input type="checkbox"/> Yes | <input type="checkbox"/> No | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| AA contains no or restricted outlet | -- | -- | .7 (M) | -- | -- | -- | -- | -- |
| AA contains unrestricted outlet | -- | -- | -- | -- | -- | -- | -- | -- |

Comments: _____

14H. SEDIMENT/Shoreline Stabilization NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

| % Cover of wetland streambank or shoreline by species with deep, binding rootmasses. | Duration of Surface Water Adjacent to Rooted Vegetation | | |
|--|---|--|--|
| | <input checked="" type="checkbox"/> Permanent / Perennial | <input type="checkbox"/> Seasonal / Intermittent | <input type="checkbox"/> Temporary / Ephemeral |
| ≥ 65 % | -- | -- | -- |
| 35-64 % | -- | -- | -- |
| < 35 % | .3 (L) | -- | -- |

Comments:

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function. A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet; P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

| A | <input type="checkbox"/> Vegetated component >5 acres | | | | | | <input checked="" type="checkbox"/> Vegetated component 1-5 acres | | | | | | <input type="checkbox"/> Vegetated component <1 acre | | | | | |
|-------|---|----------------------------|-----------------------------------|----------------------------|------------------------------|----------------------------|---|----------------------------|--|----------------------------|------------------------------|----------------------------|--|----------------------------|-----------------------------------|----------------------------|------------------------------|----------------------------|
| B | <input type="checkbox"/> High | | <input type="checkbox"/> Moderate | | <input type="checkbox"/> Low | | <input type="checkbox"/> High | | <input checked="" type="checkbox"/> Moderate | | <input type="checkbox"/> Low | | <input type="checkbox"/> High | | <input type="checkbox"/> Moderate | | <input type="checkbox"/> Low | |
| C | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input checked="" type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N | <input type="checkbox"/> Y | <input type="checkbox"/> N |
| P/P | -- | -- | -- | -- | -- | -- | -- | -- | .8H | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| S/I | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| T/E/A | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- |

Comments:

14J. GROUNDWATER DISCHARGE/RECHARGE (D/R) (Check the indicators in i & ii below that apply to the AA)

i. **Discharge Indicators**

- Springs are known or observed.
- Vegetation growing during dormant season/drought.
- Wetland occurs at the toe of a natural slopes.
- Seeps are present at the wetland edge.
- AA permanently flooded during drought periods.
- Wetland contains an outlet, but no inlet.
- Other

ii. **Recharge Indicators**

- Permeable substrate presents without underlying impeding layer.
- Wetland contains inlet but not outlet.
- Other

iii. **Rating:** Use the information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

| Criteria | Functional Point and Rating |
|---|-----------------------------|
| AA has known Discharge/Recharge area or one or more indicators of D/R present | 1 (H) |
| No Discharge/Recharge indicators present | -- |
| Available Discharge/Recharge information inadequate to rate AA D/R potential | -- |

Comments: intermittent stream flowing into reservoir likely provides groundwater inflows; no surface outlet

14K. UNIQUENESS

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

| Replacement Potential | AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP. | | | AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP. | | | AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate. | | |
|---------------------------------------|--|---------------------------------|-----------------------------------|---|---------------------------------|-----------------------------------|---|--|-----------------------------------|
| Estimated Relative Abundance from #11 | <input type="checkbox"/> rare | <input type="checkbox"/> common | <input type="checkbox"/> abundant | <input type="checkbox"/> rare | <input type="checkbox"/> common | <input type="checkbox"/> abundant | <input type="checkbox"/> rare | <input checked="" type="checkbox"/> common | <input type="checkbox"/> abundant |
| Low disturbance at AA (#12i) | -- | -- | -- | -- | -- | -- | -- | -- | -- |
| Moderate disturbance at AA (#12i) | -- | -- | -- | -- | -- | -- | -- | .3L | -- |
| High disturbance at AA (#12i) | -- | -- | -- | -- | -- | -- | -- | -- | -- |

Comments:

14L. RECREATION / EDUCATION POTENTIAL

i. Is the AA a known recreational or educational site? Yes (Rate High (1.0), then proceed to 14L(ii) only] No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA: Educational / scientific study Consumptive rec. Non-consumptive rec. Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

- Yes [Proceed to 14L (ii) and then 14L(iv).]
- No [Rate as low in 14L(iv)]

iv. **Rating** (Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

| Ownership | Disturbance at AA from #12(i) | |
|-------------------|-------------------------------|--|
| | <input type="checkbox"/> Low | <input checked="" type="checkbox"/> Moderate |
| Public ownership | -- | .5(M) |
| Private ownership | -- | -- |

Comments: _____

FUNCTION, VALUE SUMMARY, AND OVERALL RATING

| Function and Value Variables | Rating | Actual Functional Points | Possible Functional Points | Functional Units (Actual Points x Estimated AA Acreage) |
|--|--------|--------------------------|--|--|
| A. Listed/Proposed T&E Species Habitat | L | 0.00 | 1 | |
| B. MT Natural Heritage Program Species Habitat | H | 1.00 | 1 | |
| C. General Wildlife Habitat | H | 0.90 | 1 | |
| D. General Fish/Aquatic Habitat | M | 0.50 | 1 | |
| E. Flood Attenuation | M | 0.50 | 1 | |
| F. Short and Long Term Surface Water Storage | H | 1.00 | 1 | |
| G. Sediment/Nutrient/Toxicant Removal | M | 0.70 | 1 | |
| H. Sediment/Shoreline Stabilization | L | 0.30 | 1 | |
| I. Production Export/Food Chain Support | H | 0.80 | 1 | |
| J. Groundwater Discharge/Recharge | H | 1.00 | 1 | |
| K. Uniqueness | L | 0.30 | 1 | |
| L. Recreation/Education Potential | M | 0.50 | 1 | |
| Totals: | | 7.50 | 12.00 | 21 |
| Percent of Total Possible Points: | | | 63% (Actual / Possible) x 100 [rd to nearest whole #] | |

| |
|--|
| <p>Category I Wetland: (Must satisfy one of the following criteria. If not proceed to Category II.)</p> <p><input type="checkbox"/> Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; or</p> <p><input type="checkbox"/> Score of 1 functional point for Uniqueness; or</p> <p><input type="checkbox"/> Score of 1 functional point for Flood Attenuation and answer to Question 14E(ii) is "yes"; or</p> <p><input type="checkbox"/> Percent of total Possible Points is > 80%.</p> |
| <p>Category II Wetland: (Criteria for Category I not satisfied and meets any one of the following Category II criteria. If not satisfied, proceed to Category IV.)</p> <p><input checked="" type="checkbox"/> Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; or</p> <p><input checked="" type="checkbox"/> Score of .9 or 1 functional point for General Wildlife Habitat; or</p> <p><input type="checkbox"/> Score of .9 or 1 functional point for General Fish/Aquatic Habitat; or</p> <p><input type="checkbox"/> "High" to "Exceptional" ratings for both General Wildlife Habitat and General Fish / Aquatic Habitat; or</p> <p><input type="checkbox"/> Score of .9 functional point for Uniqueness; or</p> <p><input type="checkbox"/> Percent of total possible points is > 65%.</p> |
| <p><input type="checkbox"/> Category III Wetland: (Criteria for Categories I, II, or IV not satisfied.)</p> |
| <p>Category IV Wetland: (Criteria for Categories I or II are not satisfied and <u>all</u> of the following criteria are met; If not satisfied, proceed to Category III.)</p> <p><input type="checkbox"/> "Low" rating for Uniqueness; and</p> <p><input type="checkbox"/> "Low" rating for Production Export / Food Chain Support; and</p> <p><input type="checkbox"/> Percent of total possible points is < 30%.</p> |

OVERALL ANALYSIS AREA (AA) RATING: (Check appropriate category based on the criteria outlined above.)

I II III IV

Appendix C

REPRESENTATIVE PHOTOGRAPHS 2004 AERIAL PHOTOGRAPH

*MDT Wetland Mitigation Monitoring
Wigeon Reservoir
Alzada, Montana*

2004 WIGEON RESERVOIR



Location: A **Description: Wetland view** **Compass Reading: 194°**



Location: B **Description: Wetland buffer** **Compass Reading: 22°**



Location: C **Description: Wetland buffer** **Compass Reading: 280°**



Location: D vicinity **Description: Wetland view**
Compass Reading: 46°



Location: E **Description: Wetland view** **Compass Reading: 0°**



Location: F **Description: Wetland view**
Compass Reading: 330°

2004 WIGEON RESERVOIR



Location: G **Description:** Upland transect end
Compass Reading: 10°



Location: H **Description:** Wetland transect end
Compass Reading: 190°



Wigeon Reservoir 2004

Appendix D

BIRD SURVEY PROTOCOL GPS PROTOCOL

*MDT Wetland Mitigation Monitoring
Wigeon Reservoir
Alzada, Montana*

BIRD SURVEY PROTOCOL

The following is an outline of the MDT Wetland Mitigation Site Monitoring Bird Survey Protocol. Though each site is vastly different, the bird survey data collection methods must be standardized to a certain degree to increase repeatability. An Area Search within a restricted time frame will be used to collect the following data: a bird species list, density, behavior, and habitat-type use. There will be some decisions that team members must make to fit the protocol to their particular site. Each of the following sections and the desired result describes the protocol established to reflect bird species use over time.

Species Use within the Mitigation Wetland: Survey Method

Result: To conduct a bird survey of the wetland mitigation site within a restricted period of time and the budget allotment.

Sites that can be circumambulated or walked throughout.

These types of sites will include ponds, enhanced historic river channels, wet meadows, and any area that can be surveyed from the entirety of its perimeter or walked throughout. If the wetland is not uncomfortably inundated, conduct several “meandering” transects through the site in an orderly fashion (record the number and approximate location/direction of the transects in the field notebook; they do not have to be formalized or staked). If a very small portion of the site cannot be crossed due to inundation, this method will also apply. Though the sizes of the site vary, each site will require surveying to the fullest extent possible within a set time limit. The optimum times to conduct the survey are in the morning hours. Conduct the survey from sunrise to no later than 11:00 AM. (Note: some sites may have to be surveyed in the late afternoon or evening due to time constraints or weather; if this is the case, record the time of day and include this information in your report discussion.) If the survey is completed before 11:00 AM and no additions are being made to the list, then the task is complete. The overall limiting factor regarding the number of hours that are spent conducting this survey is the number of budgeted hours; this determination must be made by site by each individual.

In many cases, binoculars will be the only instrument that is needed to identify and count the birds using the wetland. If the wetland includes deep water habitat that can not be assessed with binoculars, then a scope and tripod are necessary. If this is the case, establish as many lookout posts as necessary from key vantage points to collect the data. Depending on the size of the open water, more time may be spent viewing the mitigation area from these vantage points than is spent walking the peripheries of more shallow-water wetlands.

Sites that cannot be circumambulated.

These types of sites will include large-bodied waters, such as reservoirs, particularly those with deep water habitat (>6 ft) close to the shore and no wetland development in that area of the shoreline. If one area of the reservoir was graded in such a way to create or enhance the development of a wetland, then that will be the area in which the ambulatory bird survey is conducted. The team member must then determine the length of the shoreline that will be surveyed during each visit.

As stated above in the ambulatory site section, these large sites most likely will have to be surveyed from established vantage points.

Species Use within the Mitigation Wetland: Data Recording

Result: A complete list of bird species using the site, an estimate of bird densities and associated behaviors, and identification of habitat use.

1. Bird Species List

Record the bird species on the Bird Survey - Field Data Sheet using the appropriate 4-letter code of the common name. The coding uses the first two letters of the first two words of the birds' common name or if one name, the first four (4) letters. For example, mourning dove is coded MODO and mallard is MALL. If an unknown individual is observed, use the following protocol and define your abbreviation at the bottom of the field data sheet: unknown shorebird: UNSB; unknown brown bird (UNBR); unknown warbler (UNWA); unknown waterfowl (UNWF). For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parentheses; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded: UNBB / FO (25). You may also note on the data sheet if that particular individual is using a constructed nest box.

2. Bird Density

In the office, sum the Bird Survey – Field Data Sheet data by species and by behavior. Record this data in the Bird Summary Table.

3. Bird Behavior

Bird behavior must be identified by what is known. When a species is simply observed, the behavior that it is immediately exhibiting is what is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair individual (BP); foraging (F); flyover (FO); loafing (L; e.g. sleeping, roosting, floating with head tucked under wing are loafing behaviors); and, nesting (N). If more behaviors are observed that do have a specific descriptive word, use them and we will add it to the protocol; descriptive words or phrases such as “migrating” or “living on site” are unknown behaviors.

4. Bird Species Habitat Use

We are interested in what bird species are using which particular habitat within the mitigation wetlands. This data is easily collected by simply recording what habitat the species was initially observed. Use the following broad category habitat classifications: aquatic bed (AB - rooted floating, floating-leaved, or submergent vegetation); forested (FO); marsh (MA – cattail, bulrush, emergent vegetation, etc. with surface water); open water (OW – primarily unvegetated); scrub-shrub (SS); and upland buffer (UP); wet meadow (WM – sedges, rushes, grasses with little to no surface water). If other categories are observed onsite that are not suggested here, we will make a new category next year.

GPS Mapping and Aerial Photo Referencing Procedure

The wetland boundaries, photograph location points and sampling locations were field located with mapping grade Trimble Geo III GPS units. The data was collected with a minimum of three positions per feature using Course/Acquisition code. The collected data was then transferred to a PC and differentially corrected to the nearest operating Community Base Station. The corrected data was then exported to ACAD drawings in Montana State Plain Coordinates NAD 83 international feet.

The GPS positions collected and processed had a 68% accuracy of 7 feet except in isolated areas of Tasks .008 and .011, where it went to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

Aerial reference points were used to position the aerial photographs. This positioning did not remove the distortion inherent in all photos; this imagery is to be used as a visual aide only. The located wetland boundaries were given a final review by the wetland biologist and adjustments were made if necessary.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

Appendix E

MACROINVERTEBRATE SAMPLING PROTOCOL AND DATA

*MDT Wetland Mitigation Monitoring
Wigeon Reservoir
Alzada, Montana*

AQUATIC INVERTEBRATE SAMPLING PROTOCOL

Equipment List

- D-frame sampling net with 1 mm mesh. Wildco is a good source of these.
- Spare net.
- 1-liter plastic sample jars, wide-mouth. VWR has these: catalog #36319-707.
- 95% ethanol: Northwest Scientific in Billings carries this.

All these other things are generally available at hardware or sporting goods stores. Make the labels on an ink jet printer preferably.

- hip waders.
- pre-printed sample labels (printed on Rite-in-the-Rain or other coated paper, two labels per sample).
- pencil.
- plastic pail (3 or 5 gallon).
- large tea strainer or framed screen.
- towel.
- tape for affixing label to jar.
- cooler with ice for sample storage.

Site Selection

Select the sampling site with these considerations in mind:

- Select a site accessible with hip waders. If substrates are too soft, lay a wide board down to walk on.
- Determine a location that is representative of the overall condition of the wetland.

Sampling

Wetland invertebrates inhabit the substrate, the water column, the stems and leaves of aquatic vegetation, and the water surface. Your goal is to sweep the collecting net through each of these habitat types, and then to combine the resulting samples into the 1-liter sample jar.

Dip out about a gallon of water into the pail. Pour about a cup of ethanol into the sample jar. Fill out the top half of the sample labels, using pencil, since ink will dissolve in the ethanol.

Ideally, you can sample a swath of water column from near-shore outward to a depth of approximately 3 feet with a long sweep of the net, keeping the net at about half the depth of the water throughout the sweep. Sweep the water surface as well. Pull the net through a vegetated area, beneath the water surface, for at least a meter of distance.

Sample the substrate by pulling the net along the bottom, bumping it against the substrate several times as you pull.



This step is optional, but it gives you a chance to see that you've collected some invertebrates. Rinse the net into the bucket, and look for insects, crustaceans, etc. If necessary, repeat the sampling process in a nearby location, and add the net contents to the bucket. Remember to sample all four environments.

Sieve the contents of the bucket through the straining device and pour or carefully scrape the contents of the strainer into the sample jar.

If you skip the bucket-and-sieve steps, simply lift handfuls of material out of the sampling net into the jars. In either case, please include some muck or mud and some vegetation in the jar. Often, you will have collected a large amount of vegetable material. If this is the case, lift out handfuls of material from the sieve into the jar, until the jar is about half full. Please limit material you include in the sample, so that there is only a single jar for each sample.

Top off the sample jar with enough ethanol to cover all the material in the jar. Leave as little headroom as possible.

It is not necessary to sample habitats in any specified order. Keep in mind that disturbing the habitats prior to sampling will chase off the animals you are trying to capture.

Complete the sample labels. Place one label inside the sample jar and tape the other label securely to the outside of the jar. Dry the jar before attaching the outer label if necessary. In some situations, it may be necessary to collect more than one sample at a site. If you take multiple samples from the same site, clearly indicate this by using individual sample numbers, along with the total number of samples collected at the site (e.g. Sample #3 of 5 total samples).

Photograph the sampled site.

Sample Handling/Shipping

- In the field, keep collected samples cool by storing them in a cooler. Only a small amount of ice is necessary.
- Inventory all samples, preparing a list of all sites and enumerating all samples, before shipping or delivering to the laboratory.
- Deliver samples to Rhithron.

MDT Wetland Mitigation Monitoring Project
Aquatic Invertebrate Monitoring
Summary 2001 - 2004

METHODS

Among other monitoring activities, aquatic invertebrate assemblages were collected at a number of mitigation wetlands throughout Montana. This report summarizes data generated from four years of collection.

The method employed to assess these wetlands is based on constructing an index using a battery of 12 bioassessment metrics or attributes (Table 1) tested and recommended by Stribling et al. (1995) in a report to the Montana Department of Health and Environmental Science. In that study, it was determined that some of the metrics were of limited use in some geographic regions, and for some wetland types. Despite that finding, all 12 metrics are used in this evaluation of mitigated wetlands, since detailed geographic information and wetland classifications were unavailable.

Scoring criteria for metrics were developed by generally following the tactic used by Stribling et al. Boxplots were generated using a statistical software package, and distributions, median values, ranges, and quartiles for each metric were examined. All sites in all years of sampling were used. Camp Creek, which was sampled in 2002, 2003, and 2004, was assessed using the tested metric battery developed for montane streams of Western Montana (Bollman 1998). The fauna at the Camp Creek site was different from that of the other sites, and suggested montane stream conditions rather than wetland conditions. For the wetlands, "optimal" scores were generally those that fell above the 75th percentile (for those metrics that decrease in value in response to stress) or below the 25th percentile (for metrics that respond to stress by an increase in value) of all scores. Additional scoring ranges were established by bisecting the range below the 75th percentile for decreasing scores (or above the 25th percentile for increasing scores) into "sub-optimal" and "poor" assessment categories. A score of 5, 3, or 1 was assigned to optimal, sub-optimal, and poor metric performance, respectively. In this way, metric values were translated into normalized metric scores, and scores for all metrics were summed to produce a total bioassessment score. Total bioassessment scores were classified according to a similar process, using the ranges and distributions of total scores for all sites studied in all years.

The purpose of constructing an index from biological attributes or metrics is to provide a means of integrating information to facilitate the determination of whether management action is needed. The nature of the action needed is not determined solely by the index score, however, but by consideration of an analysis of the component metrics, the taxonomic composition of the assemblages, and other issues. The diagnostic functions of the metrics and taxonomic data need more study; our understanding of the interrelationships of natural environmental factors and anthropogenic disturbances are tentative. Thus, the further interpretive remarks accompanying the raw taxonomic and metric data are offered cautiously.

Sample processing

Aquatic invertebrate samples were collected at mitigation wetland sites in the summer months of 2001, 2002, 2003, and 2004 by personnel of Land and Water Consulting, Inc. Sampling procedures utilized were based on the protocols developed by the Montana Department of Environmental Quality (MT DEQ). Sampling consisted of D-frame net sweeps through emergent vegetation (when present), the water column, over the water surface, and included disturbing and scraping substrates at each sampled sites. Samples were preserved in ethanol at each wetland site and subsequently delivered to Rhithron Associates, Inc. for processing, taxonomic determinations, and data analysis.

At Rhithron's laboratory, Caton subsamplers and stereomicroscopes with 10X magnification were used to randomly select a minimum of 100 organisms, when possible, from each sample. In some cases, the entire sample contained fewer than 100 organisms; in these cases, all organisms from the sample were taken. Taxa were identified in general accordance with the taxonomic resolution standards set out in the MT DEQ Standard Operating Procedures for Sampling and Sample Analysis (Bukantis 1998). All samples were re-identified by a second taxonomist for quality assurance purposes. The identified samples have been archived at Rhithron's laboratory. Taxonomic data and organism counts were entered into an Excel 2000 spreadsheet, and metrics were calculated and scored using spreadsheet formulae.

Bioassessment metrics

An index based on the performance of 12 metrics was constructed, as described above. Table 1 lists those metrics, describes their calculation and the expected response of each to increased degradation or impairment of the wetland.

In addition to the summed scores of each metric and the associated impairment classification described above, each individual metric informs the bioassessment to some degree. The four richness metrics (Total taxa, POET, Chironomidae taxa, and Crustacea taxa + Mollusca taxa) can be interpreted to express habitat complexity as well as water quality. Complex, diverse habitats consist of variable substrates, emergent vegetation, variable water depths and other factors, and are potential features of long-established stable wetlands with minimal human disturbance. In the study conducted by Stribling et al. (1995), all four richness metrics were found to be significantly associated with water quality parameters including conductance, salinity, and total dissolved solids.

Four composition metrics (%Chironomidae, %Orthocladinae of Chironomidae, %Crustacea + %Mollusca, and %Amphipoda) measure the relative contributions of certain taxonomic groups that may have significant responses to habitat and/or water quality impacts. For example, amphipods have been demonstrated to increase in abundance in alkaline conditions. Short-lived, relatively mobile taxa such as chironomids dominate ephemeral environments; many are hemoglobin-bearers capable of tolerating de-oxygenated conditions.

Two tolerance metrics (the Hilsenhoff Biotic Index and %Dominant taxon) were included in the bioassessment battery. The HBI indicates the overall invertebrate assemblage tolerance to nutrient enrichment, warm water, and/or low dissolved oxygen conditions. The percent abundance of the dominant taxon has been demonstrated to be strongly associated with pH, conductance, salinity, total organic carbon, and total dissolved solids.

Two trophic measures (%Collector-gatherers and %Filterers) may be helpful in expressing functional integrity of the invertebrate assemblage, which can be impacted by poor water quality or habitat degradation. High proportions of filtering organisms suggest nutrient and/or organic enrichment, while abundant collectors suggest more positive functional conditions and well-developed wetland morphology. These organisms graze periphyton growing on stable surfaces such as macrophytes.

RESULTS

In 2001, 29 sites were sampled statewide. Nineteen of these sites were revisited in 2002, and 13 new sites were sampled. In 2003, 17 sites that had been visited in both 2001 and 2002 were re-sampled, and 11 sites sampled for the first time in 2001 were re-visited. In addition, 2 new sites were sampled. In 2004, 25 sites were re-visited, and 6 new sites were sampled. Thus, the 2004 database contains data for 122 sampling events at 50 unique sites. Table 2 summarizes sites and sampling years.

Metric scoring criteria were re-developed each year as new data was added. For 2004, all 122 records were utilized. Ranges of individual metrics, as well as median metric values remained remarkably consistent in each of the 4 years; minimal changes resulted from the addition of new data in 2004. The summary metric values and scores for the 2004 samples are given in Tables 3a-3d.

Literature cited

Bollman, W. 1998. Montana Valleys and Foothill Prairies Ecoregion. Master's Thesis. (M.S.) University of Montana. Missoula, Montana.

Bukantis, R. 1998. Rapid bioassessment macroinvertebrate protocols: Sampling and sample analysis SOP's. Working draft. Montana Department of Environmental Quality. Planning Prevention and Assistance Division. Helena, Montana.

Stribling, J.B., J. Lathrop-Davis, M.T. Barbour, J.S. White, and E.W. Leppo. 1995. Evaluation of environmental indicators for the wetlands of Montana: the multimetric approach using benthic macroinvertebrates. Report to the Montana Department of Health and Environmental Science. Helena, Montana.

Table 1. Aquatic invertebrate metrics employed in the MTDT mitigation wetland monitoring study, 2001- 2004.

| Metric | Metric Calculation | Expected Response to Degradation or Impairment |
|--------------------------------|---|---|
| Total taxa | Count of unique taxa identified to lowest recommended taxonomic level | Decrease |
| POET | Count unique Plecoptera, Trichoptera, Ephemeroptera, and Odonata taxa identified to lowest recommended taxonomic level | Decrease |
| Chironomidae taxa | Count unique midge taxa identified to lowest recommended taxonomic level | Decrease |
| Crustacea taxa + Mollusca taxa | Count unique Crustacea taxa and Mollusca taxa identified to lowest recommended taxonomic level | Decrease |
| % Chironomidae | Percent abundance of midges in the subsample | Increase |
| Orthocladiinae/Chironomidae | Number of individual midges in the sub-family Orthocladiinae / total number of midges in the subsample. | Decrease |
| %Amphipoda | Percent abundance of amphipods in the subsample | Increase |
| %Crustacea + %Mollusca | Percent abundance of crustaceans in the subsample plus percent abundance of molluscs in the subsample | Increase |
| HBI | Relative abundance of each taxon multiplied times that taxon's modified Hilsenhoff Biotic Index value. These numbers are summed over all taxa in the subsample. | Increase |
| %Dominant taxon | Percent abundance of the most abundant taxon in the subsample | Increase |
| %Collector-Gatherers | Percent abundance of organisms in the collector-gatherer functional group | Decrease |
| %Filterers | Percent abundance of organisms in the filterer functional group | Increase |

Table 2. Montana Department of Transportation Mitigated Wetlands Monitoring Project sites. 2001 – 2004.

| 2001 | 2002 | 2003 | 2004 |
|-------------------------|-------------------------|-------------------------|-------------------------|
| Beaverhead 1 | Beaverhead 1 | Beaverhead 1 | Beaverhead 1 |
| Beaverhead 2 | Beaverhead 2 | | |
| Beaverhead 3 | Beaverhead 3 | | Beaverhead 3 |
| Beaverhead 4 | Beaverhead 4 | Beaverhead 4 | |
| Beaverhead 5 | Beaverhead 5 | Beaverhead 5 | Beaverhead 5 |
| Beaverhead 6 | Beaverhead 6 | Beaverhead 6 | Beaverhead 6 |
| Big Sandy 1 | | | |
| Big Sandy 2 | | | |
| Big Sandy 3 | | | |
| Big Sandy 4 | | | |
| Johnson-Valier | | | |
| VIDA | | | |
| Cow Coulee | Cow Coulee | Cow Coulee | |
| Fourchette - Puffin | Fourchette - Puffin | Fourchette - Puffin | Fourchette - Puffin |
| Fourchette - Flashlight | Fourchette - Flashlight | Fourchette - Flashlight | Fourchette - Flashlight |
| Fourchette - Penguin | Fourchette - Penguin | Fourchette - Penguin | Fourchette - Penguin |
| Fourchette - Albatross | Fourchette - Albatross | Fourchette - Albatross | Fourchette - Albatross |
| Big Spring | Big Spring | Big Spring | Big Spring |
| Vince Ames | | | |
| Ryegate | | | |
| Lavinia | | | |
| Stillwater | Stillwater | Stillwater | Stillwater |
| Roundup | Roundup | Roundup | Roundup |
| Wigeon | Wigeon | Wigeon | Wigeon |
| Ridgeway | Ridgeway | Ridgeway | Ridgeway |
| Musgrave - Rest. 1 |
| Musgrave - Rest. 2 |
| Musgrave - Enh. 1 |
| Musgrave - Enh. 2 | | | |
| | Hoskins Landing | Hoskins Landing | Hoskins Landing |
| | Peterson - 1 | Peterson - 1 | Peterson - 1 |
| | Peterson - 2 | | Peterson - 2 |
| | Peterson - 4 | Peterson - 4 | Peterson - 4 |
| | Peterson - 5 | Peterson - 5 | Peterson - 5 |
| | Jack Johnson - main | Jack Johnson - main | |
| | Jack Johnson - SW | Jack Johnson - SW | |
| | Creston | Creston | Creston |
| | Lawrence Park | | |
| | Perry Ranch | | |
| | SF Smith River | SF Smith River | SF Smith River |
| | Camp Creek | Camp Creek | Camp Creek |
| | Kleinschmidt | Kleinschmidt - pond | Kleinschmidt - pond |
| | | Kleinschmidt - stream | Kleinschmidt - stream |
| | | Ringling - Galt | |
| | | | Circle |
| | | | Cloud Ranch Pond |
| | | | Cloud Ranch Stream |
| | | | Colloid |
| | | | Jack Creek |
| | | | Norem |

Table 3a.

| | BEAVER HEAD #1 | BEAVER HEAD #3 | BEAVER HEAD #5 | BEAVER HEAD #6 | BIG SPRING CREEK | CIRCLE | CLOUD RANCH POND | CLOUD RANCH STREAM | COLLOID | CRESTON |
|------------------------|-------------------|-------------------|-------------------|-------------------|------------------------|-------------|------------------------|--------------------------|-------------|-------------|
| Total taxa | 27 | 12 | 21 | 18 | 25 | 16 | 16 | 20 | 8 | 18 |
| POET | 3 | 0 | 2 | 3 | 4 | 2 | 2 | 4 | 2 | 3 |
| Chironomidae taxa | 7 | 5 | 5 | 5 | 8 | 5 | 6 | 11 | 1 | 2 |
| Crustacea + Mollusca | 7 | 3 | 4 | 6 | 7 | 1 | 6 | 1 | 1 | 7 |
| % Chironomidae | 0.33636 | 0.18888 | 0.39285 | 0.57547 | 0.44329 | 0.55855 | 0.41666 | 0.84 | 0.09090 | 0.06087 |
| Orthoclaadiinae/Chir | 0.05405 | 0.35294 | 0.06818 | 0.36065 | 0.27907 | 0.69354 | 0.4 | 0.16666 | 0 | 0 |
| %Amphipoda | 0.03636 | 0 | 0.01785 | 0.05660 | 0.05154 | 0 | 0.00925 | 0 | 0 | 0 |
| %Crustacea + %Mollusca | 0.31818 | 0.73333 | 0.05357 | 0.12264 | 0.18556 | 0.03603 | 0.36111 | 0.01 | 0.09090 | 0.73913 |
| HBI | 7.97169 | 7.88888 | 8.36363 | 8.15789 | 7.61855 | 7.19090 | 7.32291 | 4.84 | 6 | 6.92173 |
| %Dominant taxon | 0.2 | 0.57777 | 0.23214 | 0.25471 | 0.23711 | 0.38738 | 0.13888 | 0.38 | 0.27272 | 0.37391 |
| %Collector-Gatherers | 0.40909 | 0.75555 | 0.51785 | 0.62264 | 0.78350 | 0.05405 | 0.67592 | 0.74 | 0.18181 | 0.29565 |
| %Filterers | 0.12727 | 0 | 0 | 0 | 0.01030 | 0.15315 | 0.09259 | 0.17 | 0 | 0.06087 |
| | | | | | | | | | | |
| Total taxa | 5 | 1 | 5 | 3 | 5 | 3 | 3 | 3 | 1 | 3 |
| POET | 3 | 1 | 1 | 3 | 5 | 1 | 1 | 5 | 1 | 3 |
| Chironomidae taxa | 5 | 3 | 3 | 3 | 5 | 3 | 3 | 5 | 1 | 1 |
| Crustacea + Mollusca | 5 | 1 | 3 | 5 | 5 | 1 | 5 | 1 | 1 | 5 |
| % Chironomidae | 3 | 3 | 3 | 1 | 1 | 1 | 1 | 1 | 5 | 5 |
| Orthoclaadiinae/Chir | 1 | 3 | 1 | 3 | 3 | 5 | 3 | 1 | 1 | 1 |
| %Amphipoda | 5 | 5 | 5 | 3 | 3 | 5 | 5 | 5 | 5 | 5 |
| %Crustacea + %Mollusca | 5 | 1 | 5 | 5 | 5 | 5 | 3 | 5 | 5 | 1 |
| HBI | 1 | 1 | 1 | 1 | 1 | 3 | 3 | 5 | 5 | 3 |
| %Dominant taxon | 5 | 1 | 5 | 5 | 5 | 3 | 5 | 3 | 5 | 3 |
| %Collector-Gatherers | 1 | 3 | 3 | 3 | 3 | 1 | 3 | 3 | 1 | 1 |
| %Filterers | 1 | 3 | 3 | 3 | 3 | 1 | 1 | 1 | 3 | 1 |
| | | | | | | | | | | |
| | 40 | 26 | 38 | 38 | 44 | 32 | 36 | 38 | 34 | 32 |
| | 0.666667 | 0.433333 | 0.633333 | 0.633333 | 0.733333 | 0.533333 | 0.6 | 0.633333 | 0.566667 | 0.533333 |
| | sub-optimal | poor | sub-optimal | sub-optimal | optimal | sub-optimal | sub-optimal | sub-optimal | sub-optimal | sub-optimal |

Table 3b.

| | FOURCHETTE CREEK ALBATROSS RESERVOIR | FOURCHETTE CREEK FLASHLIGHT RESERVOIR | FOURCHETTE CREEK PENGUIN RESERVOIR | FOURCHETTE CREEK PUFFIN RESERVOIR | JACK CREEK | MDT CAMP CREEK | MDT HOSKINS LANDING | MDT KLEINSCHMIDT CREEK | MDT KLEINSCHMIDT POND |
|------------------------|---|--|---|--|---------------|----------------------|---------------------------|------------------------------|-----------------------------|
| Total taxa | 18 | 23 | 19 | 22 | 23 | 35 | 25 | 19 | 19 |
| POET | 3 | 5 | 4 | 3 | 5 | 12 | 4 | 4 | 6 |
| Chironomidae taxa | 6 | 9 | 6 | 4 | 8 | 14 | 4 | 6 | 4 |
| Crustacea + Mollusca | 3 | 4 | 5 | 8 | 7 | 1 | 6 | 2 | 4 |
| % Chironomidae | 0.135135 | 0.265306 | 0.066116 | 0.247934 | 0.352113 | 0.37963 | 0.036697 | 0.438776 | 0.047619 |
| Orthoclaadiinae/Chir | 0.2 | 0.346154 | 0.625 | 0.3 | 0.52 | 0.585366 | 0.5 | 0.627907 | 0.8 |
| %Amphipoda | 0.126126 | 0.336735 | 0.578512 | 0.041322 | 0.028169 | 0 | 0.018349 | 0.010204 | 0.009524 |
| %Crustacea + %Mollusca | 0.684685 | 0.387755 | 0.77686 | 0.371901 | 0.380282 | 0.111111 | 0.541284 | 0.061224 | 0.190476 |
| HBI | 7.972973 | 7.216495 | 7.7 | 6.950413 | 7.647059 | 4.570093 | 6.59633 | 6.561224 | 6.67619 |
| %Dominant taxon | 0.495495 | 0.336735 | 0.561983 | 0.140496 | 0.15493 | 0.111111 | 0.366972 | 0.316327 | 0.552381 |
| %Collector-Gatherers | 0.873874 | 0.816327 | 0.702479 | 0.38843 | 0.394366 | 0.416667 | 0.091743 | 0.683673 | 0.114286 |
| %Filterers | 0 | 0.010204 | 0.132231 | 0.008264 | 0.042254 | 0.12037 | 0.018349 | 0.153061 | 0.047619 |
| Total taxa | | | | | | | | | |
| POET | 3 | 5 | 3 | 5 | 5 | 5 | 5 | 3 | 3 |
| Chironomidae taxa | 3 | 5 | 5 | 3 | 5 | 5 | 5 | 5 | 5 |
| Crustacea + Mollusca | 3 | 5 | 3 | 3 | 5 | 5 | 3 | 3 | 3 |
| % Chironomidae | 1 | 3 | 3 | 5 | 5 | 1 | 5 | 1 | 3 |
| Orthoclaadiinae/Chir | 5 | 3 | 5 | 3 | 3 | 3 | 5 | 1 | 5 |
| %Amphipoda | 3 | 3 | 5 | 3 | 5 | 5 | 5 | 5 | 5 |
| %Crustacea + %Mollusca | 3 | 1 | 1 | 3 | 5 | 5 | 5 | 5 | 5 |
| HBI | 1 | 3 | 1 | 3 | 3 | 5 | 3 | 5 | 5 |
| %Dominant taxon | 1 | 3 | 1 | 3 | 1 | 5 | 5 | 5 | 5 |
| %Collector-Gatherers | 1 | 5 | 1 | 5 | 5 | 5 | 3 | 5 | 1 |
| %Filterers | 5 | 5 | 3 | 1 | 1 | 1 | 1 | 3 | 1 |
| | 3 | 3 | 1 | 3 | 3 | 1 | 3 | 1 | 3 |
| | | | | | | | | | |
| | 32 | 44 | 32 | 40 | 46 | 46 | 48 | 42 | 44 |
| | 0.533333 | 0.733333 | 0.533333 | 0.666667 | 0.766667 | 0.766667 | 0.8 | 0.7 | 0.733333 |
| | sub-optimal | optimal | sub-optimal | optimal | optimal | optimal | optimal | optimal | optimal |

Table 3d.

| | ROUNDUP | SOUTH FORK SMITH RIVER | STILLWATER | WIGEON |
|------------------------|----------|---------------------------------|------------|-------------|
| Total taxa | 9 | 20 | 23 | 16 |
| POET | 0 | 5 | 4 | 3 |
| Chironomidae taxa | 4 | 7 | 9 | 5 |
| Crustacea + Mollusca | 3 | 3 | 4 | 3 |
| % Chironomidae | 0.55 | 0.482143 | 0.466667 | 0.314815 |
| Orthoclaadiinae/Chir | 0.072727 | 0.055556 | 0.244898 | 0.647059 |
| %Amphipoda | 0 | 0.071429 | 0.12381 | 0.481481 |
| %Crustacea + %Mollusca | 0.42 | 0.116071 | 0.180952 | 0.574074 |
| HBI | 8.89 | 6.589286 | 6.47619 | 7.534653 |
| %Dominant taxon | 0.28 | 0.294643 | 0.133333 | 0.481481 |
| %Collector-Gatherers | 0.56 | 0.839286 | 0.628571 | 0.657407 |
| %Filterers | 0.14 | 0 | 0 | 0.083333 |
| | | | | |
| Total taxa | | | | |
| POET | 1 | 3 | 5 | 3 |
| Chironomidae taxa | 1 | 5 | 5 | 3 |
| Crustacea + Mollusca | 3 | 5 | 5 | 3 |
| % Chironomidae | 1 | 1 | 3 | 1 |
| Orthoclaadiinae/Chir | 1 | 1 | 1 | 3 |
| %Amphipoda | 1 | 1 | 3 | 5 |
| %Crustacea + %Mollusca | 5 | 3 | 3 | 1 |
| HBI | 3 | 5 | 5 | 3 |
| %Dominant taxon | 1 | 5 | 5 | 3 |
| %Collector-Gatherers | 5 | 5 | 5 | 3 |
| %Filterers | 3 | 5 | 3 | 3 |
| | 1 | 3 | 3 | 1 |
| | | | | |
| | 26 | 42 | 46 | 32 |
| | 0.433333 | 0.7 | 0.766667 | 0.533333 |
| | poor | optimal | optimal | Sub-optimal |

Aquatic Invertebrate Taxonomic Data

Site Name WIGEON

| Order | Family | Taxon | Date Collected | | | | |
|-----------------------|----------------|--------------------------------|----------------|---------|--------|----|-----|
| | | | Count | Percent | Unique | BI | FFG |
| | | Nematoda | 1 | 0.93% | Yes | 5 | PA |
| | | Ostracoda | 2 | 1.85% | Yes | 8 | CG |
| Amphipoda | Talitridae | | | | | | |
| | | <i>Hyaletta</i> | 52 | 48.15% | Yes | 8 | CG |
| Coleoptera | Haliplidae | | | | | | |
| | | <i>Haliphus</i> | 1 | 0.93% | Yes | 5 | PH |
| | Hydrophilidae | | | | | | |
| | | <i>Berosus</i> | 1 | 0.93% | Yes | 5 | PR |
| Diplostraca | | | | | | | |
| | | Cladocera | 8 | 7.41% | Yes | 8 | CF |
| Diptera | Chironomidae | | | | | | |
| | | <i>Apedilum</i> | 7 | 6.48% | Yes | 11 | CG |
| | | <i>Cladotanytarsus</i> | 1 | 0.93% | Yes | 7 | CG |
| | | <i>Cricotopus (Isocladius)</i> | 22 | 20.37% | Yes | 7 | SH |
| | | <i>Pseudochironomus</i> | 3 | 2.78% | Yes | 5 | CG |
| | | <i>Tanytarsus</i> | 1 | 0.93% | Yes | 6 | CF |
| Ephemeroptera | Baetidae | | | | | | |
| | | <i>Callibaetis</i> | 2 | 1.85% | Yes | 9 | CG |
| | Caenidae | | | | | | |
| | | <i>Caenis</i> | 4 | 3.70% | Yes | 7 | CG |
| Heteroptera | Corixidae | | | | | | |
| | | Corixidae | 1 | 0.93% | Yes | 10 | PH |
| Odonata | Coenagrionidae | | | | | | |
| | | Coenagrionidae | 1 | 0.93% | Yes | 7 | PR |
| Trombidiformes | | | | | | | |
| | | Acari | 1 | 0.93% | Yes | 5 | PR |
| Grand Total | | | 108 | | | | |

Aquatic Invertebrate Data Summary

Project ID: MDT04LW
STORET Station ID:
Station Name: WIGEON

Activity ID:

Sample Date:

| | |
|------------------------------------|--------|
| Sample type | |
| SUBSAMPLE TOTAL ORGANISMS | 108 |
| Portion of sample used | 34.17% |
| Estimated number in total sample | 316 |
| Conversion factor | 3.937 |
| Estimated number in 1 square meter | 425 |
| Sampling effort | |
| Habitat type | |
| EPT abundance | 6 |
| Taxa richness | 16 |
| Number EPT taxa | 2 |
| Percent EPT | 5.56% |

| | | |
|-------------------------|-----------|---------|
| DOMINANCE | | |
| TAXON | ABUNDANCE | PERCENT |
| Hyalella | 52 | 48.15% |
| Cricotopus (Isocladius) | 22 | 20.37% |
| Cladocera | 8 | 7.41% |
| Apedium | 7 | 6.48% |
| Caenis | 4 | 3.70% |
| SUBTOTAL 5 DOMINANTS | | |
| | 93 | 86.11% |
| Pseudochironomus | 3 | 2.78% |
| Ostracoda | 2 | 1.85% |
| Callibaetis | 2 | 1.85% |
| Nematoda | 1 | 0.93% |
| Acari | 1 | 0.93% |
| TOTAL DOMINANTS | 102 | 94.44% |

| | | | |
|------------------------------|---------|-----------|-------|
| TAXONOMIC COMPOSITION | | | |
| GROUP | PERCENT | ABUNDANCE | #TAXA |
| Non-insect taxa | 59.26% | 64 | 5 |
| Odonata | 0.93% | 1 | 1 |
| Ephemeroptera | 5.56% | 6 | 2 |
| Plecoptera | 0.00% | 0 | 0 |
| Heteroptera | 0.93% | 1 | 1 |
| Megaloptera | 0.00% | 0 | 0 |
| Trichoptera | 0.00% | 0 | 0 |
| Lepidoptera | 0.00% | 0 | 0 |
| Coleoptera | 1.85% | 2 | 2 |
| Diptera | 0.00% | 0 | 0 |
| Chironomidae | 31.48% | 34 | 5 |


| | |
|-------------------------|---------|
| TAXONOMIC RATIOS | |
| METRIC | VALUE |
| EPT/Chironomidae | 0.18 |
| Baetidae/Ephemeroptera | 0.33 |
| Hydrosychidae/Trichopt | #DIV/0! |

| | |
|-------------------------------------|-------|
| TOLERANCE/CONDITION INDICES | |
| METRIC | VALUE |
| Community Tolerance Quotient (CTQa) | 93.27 |
| Hilsenhoff Biotic Index | 7.53 |

| | |
|------------------|-------|
| DIVERSITY | |
| METRIC | VALUE |
| Shannon H (loge) | 2.22 |
| Shannon H (log2) | 1.54 |
| Margalef D | 3.20 |
| Simpson D | 0.28 |
| P evenness | 0.10 |

| | | | |
|--------------------|-----------|--------|---------|
| VOLUNTINISM | | | |
| TYPE | ABUNDANCE | # TAXA | PERCENT |
| Multivoltine | 48 | 10 | 44.44% |
| Univoltine | 58 | 4 | 53.70% |
| Semivoltine | 2 | 2 | 1.85% |

| | | |
|------------------------|---------|--------|
| TAXA CHARACTERS | | |
| #TAXA | PERCENT | |
| Tolerant | 6 | 9.26% |
| Sensitive | 0 | 0.00% |
| Clinger | 2 | 21.30% |





| | | | |
|-------------------------------|---------|-----------|-------|
| FUNCTIONAL COMPOSITION | | | |
| GROUP | PERCENT | ABUNDANCE | #TAXA |
| Predator | 2.78% | 3 | 3 |
| Parasite | 0.93% | 1 | 1 |
| Gatherer | 65.74% | 71 | 7 |
| Filterer | 8.33% | 9 | 2 |
| Herbivore | 0.00% | 0 | 0 |
| Piercer | 1.85% | 2 | 2 |
| Scraper | 0.00% | 0 | 0 |
| Shredder | 20.37% | 22 | 1 |
| Omnivore | 0.00% | 0 | 0 |
| Unknown | 0.00% | 0 | 0 |

| | |
|----------------------------|-------|
| FUNCTIONAL RATIOS | |
| METRIC | VALUE |
| Scraper/Filterer | 0.00 |
| Scraper/Scraper + Filterer | 0.00 |

| | | |
|------------------------------|--------|-------|
| BIOASSESSMENT INDICES | | |
| B-IBI (Karr et al.) | | |
| METRIC | VALUE | SCORE |
| Taxa richness | 16 | 1 |
| E richness | 2 | 1 |
| P richness | 0 | 1 |
| T richness | 0 | 1 |
| Long-lived | 2 | 1 |
| Sensitive richness | 0 | 1 |
| %tolerant | 9.26% | 5 |
| %predators | 2.78% | 1 |
| Clinger richness | 2 | 1 |
| %dominance (3) | 75.93% | 1 |
| TOTAL SCORE | 14 | 28% |

| | | | | |
|--|----------|-------------------|----------------------------------|---------------------|
| MONTANA DEQ INDICES (Bukantis 1998) | | | | |
| METRIC | VALUE | Plains Ecoregions | Valleys and Foothills Ecoregions | Mountain Ecoregions |
| Taxa richness | 16 | 1 | 1 | 0 |
| EPT richness | 2 | 0 | 0 | 0 |
| Biotic Index | 7.53 | 0 | 0 | 0 |
| %Dominant taxon | 48.15% | 1 | 1 | 0 |
| %Collectors | 74.07% | 2 | 2 | 1 |
| %EPT | 5.56% | 0 | 0 | 0 |
| Shannon Diversity | 1.54 | 0 | 0 | 0 |
| %Scrapers + Shredder | 20.37% | 2 | 2 | 0 |
| Predator taxa | 3 | 1 | 1 | 0 |
| %Multivoltine | 44.44% | 2 | 2 | 0 |
| %H of T | #DIV/0! | #DIV/0! | #DIV/0! | #DIV/0! |
| TOTAL SCORES | 9 | #DIV/0! | #DIV/0! | 1 |
| PERCENT OF MAXIMUM | 30.00 | #DIV/0! | #DIV/0! | 4.76 |
| IMPAIRMENT CLASS | MODERATE | #DIV/0! | #DIV/0! | SEVERE |



| | |
|----------------------------------|-------|
| COMMUNITY TOLERANCES | |
| Sediment tolerant taxa | 0 |
| Percent sediment tolerant | 0.00% |
| Sediment sensitive taxa | 0 |
| Percent sediment sensitive | 0.00% |
| Metals tolerance index (McGuire) | 3.07 |
| Cold stenotherm taxa | 0 |
| Percent cold stenotherms | 0.00% |

| | | | |
|---|--------|-----------------------|----------|
| Montana Valleys and Foothills revised index (Bollman 1998) | | | |
| Percent max. | 27.78% | Impairment class | MODERATE |
| Montana Plains ecoregions metrics (Bramblett and Johnson 2002) | | | |
| Riffle | | Pool | |
| EPT richness | 2 | E richness | 2 |
| Percent EPT | 5.56% | T richness | 0 |
| Percent Oligochaetes and Leeches | 0.00% | Percent EPT | 5.56% |
| Percent 2 dominants | 68.52% | Percent non-insect | 59.26% |
| Filterer richness | 2 | Filterer richness | 2 |
| Percent intolerant | 0.00% | Univoltine richness | 4 |
| Univoltine richness | 4 | Percent supertolerant | 66.67% |
| Percent clingers | 21.30% | | |
| Swimmer richness | 4 | | |

| | |
|----------------------------|-------|
| HABITUS MEASURES | |
| Hemoglobin bearer richness | 1 |
| Percent hemoglobin bearers | 2.78% |
| Air-breather richness | 1 |
| Percent air-breathers | 0.93% |
| Burrower richness | 1 |
| Percent burrowers | 2.78% |
| Swimmer richness | 4 |
| Percent swimmers | 4.63% |